

2023 ESC Guidelines for the management of acute coronary syndromes

Supplementary data

Developed by the task force on the management of acute coronary syndromes of the European Society of Cardiology (ESC)

Authors/Task Force Members: Robert A. Byrne ^{*†}, (Chairperson) (Ireland), Xavier Rossello [‡], (Task Force Co-ordinator) (Spain), J.J. Coughlan [‡], (Task Force Co-ordinator) (Ireland), Emanuele Barbato  (Italy), Colin Berry  (United Kingdom), Alaide Chieffo  (Italy), Marc J. Claeys  (Belgium), Gheorghe-Andrei Dan  (Romania), Marc R. Dweck  (United Kingdom), Mary Galbraith  (United Kingdom), Martine Gilard (France), Lynne Hinterbuchner  (Austria), Ewa A. Jankowska  (Poland), Peter Jüni (United Kingdom), Takeshi Kimura (Japan), Vijay Kunadian  (United Kingdom), Margret Leosdottir  (Sweden), Roberto Lorusso  (Netherlands), Roberto F.E. Pedretti  (Italy), Angelos G. Rigopoulos  (Greece), Maria Rubini Gimenez  (Germany), Holger Thiele (Germany), Pascal Vranckx (Belgium), Sven Wassmann (Germany), Nanette Kass Wenger (United States of America), Borja Ibanez ^{*†}, (Chairperson) (Spain), and ESC Scientific Document Group

* Corresponding authors: Robert A. Byrne, Department of Cardiology and Cardiovascular Research Institute (CVRI) Dublin, Mater Private Network, Dublin, Ireland, and School of Pharmacy and Biomolecular Sciences, RCSI University of Medicine and Health Sciences, Dublin, Ireland. Tel: +353-1-2483190, E-mail: robertabyrne@rcsi.ie; and Borja Ibanez, Clinical Research Department, Centro Nacional de Investigaciones Cardiovasculares Carlos III (CNIC), Madrid, Spain, and Cardiology Department, IIS-Fundación Jiménez Díaz University Hospital, Madrid, Spain, CIBERCV, ISCIII, Madrid, Spain. Tel: +3491 4531200, E-mail: bibanez@cnic.es.

[†] The two Chairpersons contributed equally to the document and are joint corresponding authors.

[‡] The two Task Force Co-ordinators contributed equally to the document.

Author/Task Force Member affiliations are listed in author information in the full text.

ESC Clinical Practice Guidelines (CPG) Committee: listed in the Appendix in the full text.

ESC subspecialty communities having participated in the development of this document:

Associations: Association of Cardiovascular Nursing & Allied Professions (ACNAP), Association for Acute CardioVascular Care (ACVVC), European Association of Cardiovascular Imaging (EACVI), European Association of Preventive Cardiology (EAPC), European Association of Percutaneous Cardiovascular Interventions (EAPCI), European Heart Rhythm Association (EHRA), and Heart Failure Association (HFA).

Working Groups: Cardiovascular Pharmacotherapy, Cardiovascular Surgery, E-Cardiology, Myocardial and Pericardial Diseases, Thrombosis.

Patient Forum

The content of these European Society of Cardiology (ESC) Guidelines has been published for personal and educational use only. No commercial use is authorized. No part of the ESC Guidelines may be translated or reproduced in any form without written permission from the ESC. Permission can be obtained upon submission of a written request to Oxford University Press, the publisher of the *European Heart Journal*, and the party authorized to handle such permissions on behalf of the ESC (journals.permissions@oup.com).

Disclaimer. The ESC Guidelines represent the views of the ESC and were produced after careful consideration of the scientific and medical knowledge and the evidence available at the time of their publication. The ESC is not responsible in the event of any contradiction, discrepancy, and/or ambiguity between the ESC Guidelines and any other official recommendations or guidelines issued by the relevant public health authorities, in particular in relation to good use of healthcare or therapeutic strategies. Health professionals are encouraged to take the ESC Guidelines fully into account when exercising their clinical judgment, as well as in the determination and the implementation of preventive, diagnostic or therapeutic medical strategies; however, the ESC Guidelines do not override, in any way whatsoever, the individual responsibility of health professionals to make appropriate and accurate decisions in consideration of each patient's health condition and in consultation with that patient and, where appropriate and/or necessary, the patient's caregiver. Nor do the ESC Guidelines exempt health professionals from taking into full and careful consideration the relevant official updated recommendations or guidelines issued by the competent public health authorities, in order to manage each patient's case in light of the scientifically accepted data pursuant to their respective ethical and professional obligations. It is also the health professional's responsibility to verify the applicable rules and regulations relating to drugs and medical devices at the time of prescription.

Document Reviewers: Sigrun Halvorsen, (Clinical Practice Guidelines Review Co-ordinator) (Norway), Stefan James, (Clinical Practice Guidelines Review Co-ordinator) (Sweden), Magdy Abdelhamid (Egypt), Victor Aboyans (France), Nina Ajmone Marsan (Netherlands), Sotiris Antoniou (United Kingdom), Riccardo Asteggiano (Italy), Maria Bäck (Sweden), Davide Capodanno (Italy), Ruben Casado-Arroyo (Belgium), Salvatore Cassese (Germany), Jelena Čelutkienė (Lithuania), Maja Cikes (Croatia), Jean-Philippe Collet (France), Gregory Ducrocq (France), Volkmar Falk (Germany), Laurent Fauchier (France), Tobias Geisler (Germany), Diana A. Gorog (United Kingdom), Lene Holmvang (Denmark), Tiny Jaarsma (Sweden), Hywel Wynne Jones (United Kingdom), Lars Køber (Denmark), Konstantinos C. Koskinas (Switzerland), Dipak Kotecha (United Kingdom), Konstantin A. Krychtiuk (Austria), Ulf Landmesser (Germany), George Lazaros (Greece), Basil S. Lewis (Israel), Bertil Lindahl (Sweden), Ales Linhart (Czech Republic), Maja-Lisa Løchen (Norway), Mamas A. Mamas (United Kingdom), John William McEvoy (Ireland), Borislava Mihaylova (United Kingdom), Richard Mindham (United Kingdom), Christian Mueller (Switzerland), Lis Neubeck (United Kingdom), Josef Niebauer (Austria), Jens Cosedis Nielsen (Denmark), Alexander Niessner (Austria), Valeria Paradies (Netherlands), Agnes A. Pasquet (Belgium), Steffen E. Petersen (United Kingdom), Eva Prescott (Denmark), Amina Rakisheva (Kazakhstan), Bianca Rocca (Italy), Giuseppe M.C. Rosano (Italy), Leyla E. Sade (United States of America), François Schiele (France), Jolanta M. Siller-Matula (Austria), Christian Sticherling (Switzerland), Robert F. Storey (United Kingdom), Matthias Thielmann (Germany), Christiaan Vrints (Belgium), Stephan Windecker (Switzerland), Rune Wiseth (Norway), and Adam Witkowski (Poland)

All experts involved in the development of these guidelines have submitted declarations of interest. These have been compiled in a report and simultaneously published in a supplementary document to the guidelines. The report is also available on the ESC website www.escardio.org/Guidelines

Keywords

Guidelines • Acute cardiac care • Acute coronary syndrome • Antithrombotic therapy • Fibrinolysis • High-sensitivity troponin • Invasive strategy • MINOCA • Myocardial infarction • Non-ST-elevation myocardial infarction • Patient-centred care • Percutaneous coronary intervention • Recommendations • Reperfusion therapy • Revascularization • Secondary prevention • ST-segment elevation myocardial infarction • Unstable angina

Table of contents

1. Preamble	6	3.3.2.2. Caveats of using rapid algorithms	12
2. Introduction	6	3.3.2.3. Practical guidance on how to implement the	
2.1. Definitions Acute coronary syndrome and myocardial		European Society of Cardiology 0 h/1 h algorithm	13
infarction	6	3.3.3. Other biomarkers	13
2.2. Epidemiology of acute coronary syndromes	6	3.4. Diagnostic tools Non-invasive imaging	13
3. Triage and diagnosis	6	3.4.1. Cardiac magnetic resonance imaging with or without	
3.1. Clinical presentation and physical examination	6	stress testing	13
3.1.1. Clinical presentation	6	3.4.2. Single-photon emission computerized tomography	
3.1.2. Physical examination	7	perfusion imaging and stress echocardiography	13
3.2. Diagnostic tools Electrocardiogram	7	3.5. Differential diagnosis for acute chest pain	13
3.2.1. Acute coronary syndrome with persistent ST-segment		4. Initial measures for patients presenting with suspected acute	
elevation or other signs of acute vessel occlusion	7	coronary syndrome Initial treatment	14
3.2.2. Acute coronary syndrome without persistent ST-		4.1. Pre-hospital logistics of care	14
segment elevation or other signs of acute vessel occlusion	10	4.1.1. Organization of ST-elevation myocardial infarction	
3.3. Diagnostic tools Biomarkers	11	treatment in networks	14
3.3.1. High-sensitivity cardiac troponin	11	5. Acute-phase management of patients with acute coronary	
3.3.1.1 Reasons for high-sensitivity cardiac troponin		syndrome	14
elevations other than Type 1 myocardial infarction	11	5.1. Acute coronary syndrome managed with invasive strategy	14
3.3.2. Rapid 'rule-in' and 'rule-out' algorithms	12	5.2. Patients not undergoing reperfusion	15
3.3.2.1. European Society of Cardiology 0 h/1 h and		5.2.1. Patients who are not candidates for invasive coronary	
European Society of Cardiology 0 h/2 h algorithms	12	angiography	15
		5.2.2. Patients with coronary artery disease not amenable to	
		revascularization	15

6. Antithrombotic therapy	15	12.1.3.4. Bleeding events related to coronary artery bypass surgery	28
6.1. Long-term treatment	15	12.1.3.5. Transfusion therapy	28
6.1.1. Prolonging antithrombotic therapy beyond 12 months	15	12.2. Comorbid conditions	28
6.2. Antiplatelet therapy in patients requiring oral anticoagulation	17	12.2.1. Patients at high bleeding risk and with blood disorders (anaemia and thrombocytopenia)	28
6.2.1. Acute coronary syndrome patients with atrial fibrillation	17	12.2.1.1. Thrombocytopenia following GP IIb/IIIa inhibitor therapy	28
6.3. Fibrinolysis and pharmaco-invasive strategy	19	12.2.1.2. Heparin-induced thrombocytopenia	28
6.3.1. Comparison of fibrinolytic agents	20	12.2.2. Chronic kidney disease	29
6.3.2. Hazards of fibrinolysis and contraindications	20	12.2.3. Older adults with frailty and multimorbidity	29
6.4. Antithrombotic therapy in patients not undergoing reperfusion	20	12.2.3.1. The older person	29
7. Acute coronary syndrome with unstable presentation	20	12.2.3.2. Frailty and multimorbidity	29
7.1. Out-of-hospital cardiac arrest in acute coronary syndrome	20	12.2.3.3. Pharmacotherapy in older and frail patients	29
7.1.1. Healthcare systems and systems of care	20	12.2.4. Pregnancy	30
8. Management of acute coronary syndrome during hospitalization	20	12.2.5. Drug abuse	31
8.1. Coronary care unit/intensive cardiac care unit	20	12.2.5.1. Acute coronary syndrome associated with alcohol dependence and illicit drug use	31
8.2. In-hospital care	20	12.2.5.2. Acute coronary syndrome associated with illicit drug use	31
8.2.1. Duration of hospital stay	20	12.2.6. Patients with cancer	31
8.2.2. Risk assessment	21	12.2.6.1. Pathophysiology	32
8.2.2.1. Clinical risk assessment	21	12.2.6.2. Clinical presentation	32
8.2.2.2. Biomarkers for risk assessment	21	12.2.6.3. Initial management and acute multidisciplinary approach	32
8.2.2.3. Bleeding risk assessment	21	12.2.6.4. Invasive strategy	32
8.2.2.4. Integrating ischaemic and bleeding risks	22	12.2.6.5. Antithrombotic treatment	32
8.2.2.4. Evaluation of long-term risk before discharge	22	12.2.6.6. Thrombocytopenia and cancer	32
9. Technical aspects of invasive strategies	22	12.2.6.7. Cancer treatment	32
9.1. Percutaneous coronary intervention	22	12.2.7. Coronavirus disease (COVID-19)	32
9.1.1. Intravascular imaging/physiology of the infarct-related artery	22	13. Long-term treatment	33
9.1.1.1. Intravascular imaging	22	13.1. Lifestyle management	33
9.1.1.2. Intravascular physiology	23	13.1.1. Nutrition and alcohol	33
9.1.2. Embolic protection and microvascular salvage strategies	23	13.1.2. Resumption of activities	33
9.1.2.1. Interventions to protect the microcirculation	23	13.1.2.1. Sexual activity	33
9.2. Ongoing major trials	24	13.1.2.2. Environmental factors	33
10. Management of patients with multivessel disease	24	13.2. Pharmacological treatment	33
11. Myocardial infarction with non-obstructive coronary arteries	24	13.2.1. Lipid-lowering therapy	33
11.1. Characteristics, prognosis, and symptoms at presentation	24	13.2.2. Hormone replacement therapy	35
11.2. Invasive coronary angiography	24	14. Patient perspectives	35
11.3. Functional coronary angiography	24	14.1. Patient-centred care	35
11.4. Intravascular imaging (intravascular ultrasound/optical coherence tomography)	24	14.2. Informed consent	35
11.5. Left ventricular angiography, pressure, and function	24	14.3. Research participation and consent in the acute setting	37
11.6. Non-invasive evaluation	24	14.4. Patient-reported outcome measures and patient-reported experience measures	37
11.7. Management of myocardial infarction with non-obstructive coronary arteries	25	14.5. Preparation for discharge	38
12. Special situations	26	15. Gaps in evidence	40
12.1. Complications	26	16. References	41
12.1.1. Heart failure	26		
12.1.1.1. Mechanical complications	26		
12.1.2. Post-acute coronary syndrome pericarditis	27		
12.1.3. Bleeding	27		
12.1.3.1. Management of bleeding	27		
12.1.3.1.1. Bleeding events on antiplatelet agents	27		
12.1.3.1.2. Bleeding events on vitamin K antagonists	27		
12.1.3.2. Bleeding events on non-vitamin K antagonist oral anticoagulants	27		
12.1.3.3. Bleeding events related to percutaneous coronary intervention	27		
		List of tables	
		Table S1 Fourth universal definition of myocardial infarction	6
		Table S2 Clinical implications of high-sensitivity cardiac troponin assays	11
		Table S3 Conditions other than acute Type 1 myocardial infarction associated with cardiomyocyte injury (i.e. cardiac troponin elevation)	11

Table S4 Assay specific cut-off levels in ng/L within the 0 h/1 h and 0 h/2 h algorithms	12
Table S5 Differential diagnoses of acute coronary syndrome in the setting of acute chest pain	13
Table S6 Key features of networks of care for the pre-hospital management of STEMI	14
Table S7 Treatment options for extended dual antithrombotic or antiplatelet therapies	16
Table S8 Risk criteria for extended treatment with a second antithrombotic agent	17
Table S9 High-risk features of stent-driven recurrent ischaemic events	17
Table S10 Doses of fibrinolytic agents and antithrombotic co-therapies	19
Table S11 Contraindications to fibrinolytic therapy	20
Table S12 Major and minor criteria for high bleeding risk according to the Academic Research Consortium for High Bleeding Risk at the time of percutaneous coronary intervention	22
Table S13 Common manifestations of myocardial ischaemia symptoms in patients with MINOCA	25
Table S14 Mechanical complications	26
Table S15 Recommended doses of antithrombotic agents in the acute care of patients with chronic kidney disease	29
Table S16 Cancer treatment-related acute coronary syndrome	31
Table S17 Characteristics of a healthy diet	33
Table S18 Components needed for informed consent	37
Table S19 Patient expectations and clear communication for patients with acute coronary syndrome	38
Table S20 Preparation for discharge	39

List of figures

Figure S1 Symptoms at presentation in acute coronary syndrome in women and men	8
Figure S2 Electrocardiographic abnormalities in patients with STEMI and ECG findings that, if present, may prompt triage for immediate reperfusion therapy	9
Figure S3 Electrocardiographic abnormalities in patients with non-ST-segment elevation acute coronary syndrome	10
Figure S4 Antithrombotic strategies beyond the first 12 months after ACS	16
Figure S5 Information for patients on how to optimize their 'heart health' after an acute coronary syndrome	34
Figure S6 Informed consent process using the 'teach back' technique	36
Figure S7 Patient concerns and educational needs throughout their ACS journey	40

Abbreviations and acronyms

ACS	Acute coronary syndrome
ACUITY	Acute Catheterization and Urgent Intervention Triage strategy
AF	Atrial fibrillation
AGRI	Australian GRACE Risk score Intervention Study
AMI	Acute myocardial infarction
APPRAISE-2	Apixaban for Prevention of Acute Ischemic Events 2
aPTT	Activated partial thromboplastin time

ARC-HBR	Academic Research Consortium for High Bleeding Risk
ASCVD	Atherosclerotic cardiovascular disease
ATLAS ACS 2-TIMI 51	Anti-Xa Therapy to Lower Cardiovascular Events in Addition to Standard Therapy in Subjects With Acute Coronary Syndrome ACS 2-Thrombolysis In Myocardial Infarction 51
AUGUSTUS	An Open-Label, 2x2 Factorial, Randomized Controlled, Clinical Trial to Evaluate the Safety of Apixaban Versus Vitamin K Antagonist and Aspirin Versus Aspirin Placebo in Patients With Atrial Fibrillation and Acute Coronary Syndrome or Percutaneous Coronary Intervention
b.i.d.	<i>Bis in die</i> (twice a day)
BNP	Brain natriuretic peptide
CAD	Coronary artery disease
CAGB	Coronary artery bypass graft(ing)
CARMEN	CARDiovascular Magnetic rEsonance imaging and computed Tomography Angiography
CCS	Chronic coronary syndrome
CCTA	Coronary computed tomography angiography
CI	Confidence interval
CKD	Chronic kidney disease
CMR	Cardiac magnetic resonance
CODE-MI	hs-cTn-Optimizing the Diagnosis of Acute Myocardial Infarction/Injury in Women
COMPASS	Cardiovascular Outcomes for People Using Anticoagulation Strategies
COVADIS	Coronary Vasomotion Disorders International Study Group
COVID-19	Coronavirus disease 2019
CRUSADE	Can Rapid risk stratification of Unstable angina patients Suppress Adverse outcomes with Early implementation of the American College of Cardiology/American Heart Association guidelines
CT	Computed tomography
cTn	Cardiac troponin
CV	Cardiovascular
CVD	Cardiovascular disease
DAPT	Dual antiplatelet therapy
DAT	Dual antithrombotic therapy
DES	Drug-eluting stent
DOCTORS	Does Optical Coherence Tomography Optimize Results of Stenting
DTU-STEMI	Primary Unloading and Delayed Reperfusion in ST-Elevation Myocardial Infarction: The STEMI-DTU Trial
ECG	Electrocardiogram
ED	Emergency department
EMS	Emergency medical services
ENTRUST-AF PCI	Edoxaban Treatment Versus Vitamin K Antagonist in Patients With Atrial Fibrillation Undergoing Percutaneous Coronary Intervention
EURO-ICE	European Intracoronary Cooling Evaluation in Patients With ST-Elevation Myocardial Infarction
FOURIER	Further Cardiovascular Outcomes Research with PCSK9 Inhibition in Subjects with Elevated Risk
FFR	Fractional flow reserve

GP	Glycoprotein	PCI	Percutaneous coronary intervention
GRACE	Global Registry of Acute Coronary Events	PCSK9	Proprotein convertase subtilisin/kexin type 9
HBR	High bleeding risk	PIONEER	A Study Exploring Two Strategies of Rivaroxaban and One of Oral Vitamin K Antagonist in Patients With Atrial Fibrillation Who Undergo Percutaneous Coronary Intervention
HDL-C	High-density lipoprotein-cholesterol	AF-PCI	PLATElet inhibition and patient Outcomes
HF	Heart failure		Ticagrelor or Prasugrel Versus Clopidogrel in Elderly Patients With an Acute Coronary Syndrome and a High Bleeding Risk: Optimization of Antiplatelet Treatment in High-risk Elderly
HIT	Heparin-induced thrombocytopenia	PLATO	Primary percutaneous coronary intervention
HR	Hazard ratio	POPular-AGE	PREdicting bleeding Complications In patients undergoing Stent implantation and subsequent Dual Anti Platelet Therapy
HRT	Hormone replacement therapy		Patient-reported experience measures
hs-cTn	High-sensitivity cardiac troponin		Patient-reported outcome measures
ICA	Invasive coronary angiography		Right bundle branch block
ICCU	Intensive cardiac care unit		Randomized controlled trial
ICH	Intracranial haemorrhage		Randomized Evaluation of Dual Antithrombotic Therapy with Dabigatran versus Triple Therapy with Warfarin in Patients with Nonvalvular Atrial Fibrillation Undergoing Percutaneous Coronary Intervention
ILUMIEN III: OPTIMIZE PCI	Optical Coherence Tomography compared to Intravascular Ultrasound and Angiography to Guide Coronary Stent Implantation: a Multicenter Randomized Trial in PCI	PPCI	Reduction of Cardiovascular Events With Icosapent Ethyl—Intervention Trial
ILUMIEN IV: OPTIMAL PCI	Optical Coherence Tomography Guided Coronary Stent Implantation Compared to Angiography: a Multicenter Randomized Trial in PCI	PRECISE-DAPT	Systolic blood pressure
IMR	Index of microvascular resistance		Spontaneous coronary artery dissection
INR	International normalized ratio		Short Duration of Dual antiplatelet Therapy With Synergy II Stent in Patients Older Than 75 Years Undergoing Percutaneous Coronary Revascularization
IRA	Infarct-related artery		Single-photon emission computerized tomography
ISAR-TRIPLE	Triple Therapy in Patients on Oral Anticoagulation After Drug Eluting Stent Implantation		ST-elevation acute coronary syndrome
i.v.	Intravenous		ST-elevation myocardial infarction
IVUS	Intravascular ultrasound		Strategic Reperfusion Early After Myocardial Infarction
LAD	Left anterior descending		Swedish Web-System for Enhancement and Development of Evidence-Based Care in Heart Disease Evaluated According to Recommended Therapies
LBBB	Left bundle branch block		Triple antithrombotic therapy
LDL-C	Low-density lipoprotein-cholesterol		Thrombolysis In Myocardial Infarction
LV	Left ventricular		Tissue plasminogen activator
LVEDP	Left ventricular end-diastolic pressure		Target vessel failure
LVEF	Left ventricular ejection fraction		Target vessel revascularization
MACE	Major adverse cardiovascular events		Unstable angina
MACCE	Major adverse cardiovascular and cerebrovascular events		Unfractionated heparin
MATRIX	Minimizing Adverse Haemorrhagic Events by Transradial Access Site and Systemic Implementation of angioX		UK GRACE Risk score Intervention Study
MCS	Mechanical circulatory support		What is the Optimal antiplatelet & Anticoagulant Therapy in Patients With Oral Anticoagulation and Coronary Stenting
MI	Myocardial infarction		Investigation of Metformin in Pre-Diabetes on Atherosclerotic Cardiovascular Outcomes
MINOCA	Myocardial infarction with non-obstructive coronary arteries		Vitamin K antagonist
MVO	Microvascular obstruction		
NOAC	Non-vitamin K antagonist oral anticoagulant		
NPV	Negative predictive value		
NSTE-ACS	Non-ST-elevation acute coronary syndrome		
NSTEMI	Non-ST-elevation myocardial infarction		
NT-pro BNP	N-terminal pro B-type natriuretic peptide		
OAC	Oral anticoagulant/anticoagulation		
OCT	Optical coherence tomography		
OFDI	Optical frequency domain imaging		
OPINION	Optical frequency domain imaging vs. Intravascular ultrasound in percutaneous coronary intervention		
OR	Odds ratio		
PAD	Peripheral arterial disease		

1. Preamble

There is no supplementary material for this section.

2. Introduction

2.1. Definitions | Acute coronary syndrome and myocardial infarction

Table S1 Fourth universal definition of myocardial infarction

Universal definition of myocardial infarction

A combination of criteria is required to meet the diagnosis of acute myocardial infarction, namely the detection of an increase and/or decrease of a cardiac biomarker, preferably high-sensitivity cardiac troponin T or I, with at least one value above the 99th percentile of the upper reference limit and at least one of the following:

- (i) Symptoms of myocardial ischaemia.
- (ii) New ischaemic ECG changes.
- (iii) Development of pathological Q waves on ECG.
- (iv) Imaging evidence of loss of viable myocardium or new regional wall motion abnormality in a pattern consistent with an ischaemic aetiology (vascular bed distribution).
- (v) Intracoronary thrombus detected on angiography or autopsy.

Different types of MI have been described on the basis of different underlying pathological conditions.

Type 1 MI	Characterized by atherosclerotic plaque rupture, ulceration, fissure, or erosion with resulting intraluminal thrombus in one or more coronary arteries, leading to decreased myocardial blood flow and/or distal embolization and subsequent myocardial necrosis. Patients diagnosed with Type 1 MI will usually have underlying obstructive coronary artery disease (i.e. >50% diameter stenosis) but in ~5–10% of cases there may be non-obstructive coronary atherosclerosis, particularly in women. ^{1–5}
Type 2 MI	Myocardial necrosis in which a condition other than coronary plaque instability causes an imbalance between myocardial oxygen supply and demand. ^{1,5} Mechanisms include hypotension, hypertension, tachyarrhythmias, bradyarrhythmias, anaemia, hypoxaemia, coronary artery spasm, spontaneous coronary artery dissection, coronary embolism, and coronary microvascular dysfunction. ^{6–8}
Type 3 MI	MI resulting in cardiac death with symptoms suggestive of myocardial ischaemia when biomarkers are not available or MI is detected at autopsy.
Type 4 MI	MI caused by percutaneous coronary intervention. ⁵
Type 5 MI	MI caused by coronary artery bypass grafting. ⁵

ECG, electrocardiogram; MI, myocardial infarction.

2.2. Epidemiology of acute coronary syndromes

The incidence of acute coronary syndrome (ACS) increases with age. On average, ACS occurs 7–10 years earlier in men than in women.^{9,10} The risk of acute coronary events is increased with exposure to traditional cardiovascular (CV) risk factors and can be estimated using risk scores, such as the European Society of Cardiology Systemic Coronary Risk Evaluation (SCORE) system.^{10,11} The incidence of ST-elevation myocardial infarction (STEMI) has decreased relative to non-ST-elevation MI (NSTEMI) in recent years.^{12,13}

There are considerable differences within European and global regions in the incidence and prevalence of ACS, alongside differences in case fatality rates.^{14,15} The increase in the relative incidence of NSTEMI is multifactorial (e.g. due to changes in diagnostic criteria and the emergence of high-sensitivity troponin assays). In Europe, there has been an overall trend toward a reduction in mortality due to ACS over the past three decades.¹⁶ Recent studies have highlighted a fall in both acute and long-term mortality following STEMI, in parallel with increasing use of reperfusion therapy, primary percutaneous coronary intervention (PPCI), modern antithrombotic therapy, and secondary prevention therapies.^{14,17,18}

Women tend to receive reperfusion therapy less frequently than men, and for high-risk groups, women less frequently receive reperfusion therapy within the recommended timeframes.^{19–21} One potentially relevant contributor to this observation is that women with ACS tend to present later than men.^{22–24} Women also have a higher risk of bleeding complications with percutaneous coronary intervention (PCI), which may influence treatment decisions.²⁵ In patients who do not undergo invasive management, mortality is higher in women than in men. As such, it is important to maintain a high degree of awareness for MI in women with potential symptoms of ischaemia and to ensure that they do not receive lower rates of guideline-recommended investigations and therapies.

3. Triage and diagnosis

3.1. Clinical presentation and physical examination

3.1.1. Clinical presentation

The relief of symptoms after nitroglycerine (glycerine trinitrate) administration may increase the likelihood of ACS, but is not specific for ACS as it is also reported in other causes of non-cardiac chest pain, such as gastrointestinal disorders.²⁶ In patients with a working diagnosis of STEMI, the administration of nitroglycerine can be misleading and is not recommended as a diagnostic manoeuvre.^{27,28} However, if symptoms resolve after nitroglycerine administration, it is recommended to obtain another 12-lead electrocardiogram (ECG). Complete normalization of ST-segment elevation after nitroglycerine administration, along with complete relief of symptoms, is suggestive of coronary spasm, with or without associated myocardial infarction (MI).

Overall, the diagnostic performance of chest pain characteristics is limited in patients presenting to the emergency department (ED) with suspected ACS.²⁶ There also appear to be more similarities than

differences between males and females in relation to symptoms associated with an ACS presentation.^{26,29} Older age, male sex, family history of coronary artery disease (CAD), diabetes, hyperlipidaemia, smoking, hypertension, renal dysfunction, previous manifestation of CAD, and peripheral or carotid artery disease all increase the likelihood of ACS.^{30,31} In addition, some conditions may exacerbate or precipitate ACS, including: anaemia, infection, inflammation, fever, hypertensive crisis, emotional stress, and metabolic or endocrine (in particular thyroid) disorders.

It is important for clinicians to be aware that the presentation of symptoms, the ability of the patient to express these symptoms, and the explanation of how the symptoms actually affect the individual may differ between men and women. However, it is also important to note that any differences in the sex-specific diagnostic performance of chest pain characteristics appear to be relatively minor and do not support the use of sex-specific chest pain characteristics for the early diagnosis of MI.²² In general, while some sex differences in symptoms exist, the symptoms experienced by men and women with confirmed ACS show substantial overlap, as demonstrated in [Figure S1](#).²² What may be of more relevance to clinical presentation is that the interpretation of cardiac symptoms by physicians may be subject to gender bias and it is important for caregivers to be aware of this.³²

Over 80% of women and men with ACS present with chest pain or pressure. Other common symptoms, like diaphoresis, shoulder/arm pain, and indigestion/epigastric pain, occur relatively commonly in both women and men with ACS. While some of the less common symptoms at presentation may be more common in women with ACS, these differences are minor and do not support the use of women-specific chest pain characteristics for the early diagnosis of MI.

As mentioned in the main text, chest pain can be classified as cardiac, possible cardiac, or non-cardiac. The term cardiac is used to describe chest pain occurring because of an underlying cardiac aetiology. This includes classic chest discomfort based on quality, location, radiation, and provoking and relieving factors that make it more likely to be of cardiac ischaemic origin. The term 'possible cardiac' is used to refer to chest pain symptoms that suggest a cardiac origin. 'Non-cardiac' is a term used to refer to chest pain symptoms likely due to a non-cardiac cause in patients with persistent or recurring symptoms despite a negative stress test or anatomic cardiac evaluation, or a low-risk designation by a clinical decision pathway.³³

3.1.2. Physical examination

Cardiac auscultation may reveal a systolic murmur due to ischaemic mitral regurgitation, which is associated with a poor prognosis.³⁴ Alternatively, the murmur of aortic stenosis may be detected, which can mimic ACS presentations and may influence subsequent revascularization strategies. Rarely, a systolic murmur may indicate a mechanical complication (i.e. papillary muscle rupture or ventricular septal defect), particularly in patients who have presented late after MI and in whom revascularization has been delayed. A murmur consistent with aortic insufficiency should prompt consideration of acute aortic dissection associated with ACS. Physical examination may identify signs of non-coronary causes of chest pain (e.g. pulmonary embolism, acute aortic syndromes, myopericarditis, or aortic stenosis) or extracardiac pathologies (e.g. pneumothorax, pneumonia, or musculoskeletal diseases). In this setting, the presence of chest pain that can be reproduced by exerting pressure on the chest wall has a relatively high negative predictive value for ACS.²⁶ According to the clinical presentation,

abdominal disorders (e.g. reflux disease, oesophageal spasm, oesophagitis, gastric ulcer, cholecystitis, or pancreatitis) may also be considered in the differential diagnosis. Differences in blood pressure between the upper and lower limbs or between the arms, irregular pulse, jugular vein distension, heart murmurs, friction rub, and pain reproduced by chest or abdominal palpation are findings suggestive of alternative diagnoses. Pallor, sweating, or tremor are often signs of stress related to MI but may also point towards precipitating conditions (i.e. anaemia or thyrotoxicosis).²

3.2. Diagnostic tools | Electrocardiogram

ECG criteria are based on changes in electrical currents of the heart (measured in millivolts). Standard calibration of the ECG is 10 mm/mV—therefore 0.1 mV equals 1 mm square on the vertical axis. For simplicity, in this document ECG deviations are expressed in mm following the standard calibration.

3.2.1. Acute coronary syndrome with persistent ST-segment elevation or other signs of acute vessel occlusion

In patients with left bundle branch block (LBBB), specific ECG criteria (Sgarbossa's criteria) may help in the detection of candidates for immediate coronary angiography.^{35,36} It is important to recognize that the presence of LBBB is not suggestive of ongoing coronary artery occlusion in isolation. However, the presence of LBBB precludes the identification of underlying ECG alterations that may indicate coronary artery occlusion. Therefore, patients with signs/symptoms that are highly suspicious for ongoing myocardial ischaemia and who have LBBB on ECG (whether previously known or not) should undergo a reperfusion strategy. Patients with LBBB and other symptoms (e.g. non-persistent chest discomfort) who are haemodynamically stable only have a slightly higher risk of having MI than patients without LBBB. In these patients, the result of the high-sensitivity cardiac troponin T or I (hs-cTn T/I) measurement at presentation should be integrated into decision-making regarding triage for (and timing of) coronary angiography.^{36,37} It is also important to consider that more than 50% of patients presenting to the ED or chest pain unit with chest discomfort and LBBB will ultimately be found to have a diagnosis other than MI.³⁶ In addition to the Sgarbossa criteria, some other novel algorithms to improve the identification of acute MI in patients with LBBB have also been proposed.^{36,38–41}

In patients with right bundle branch block (RBBB), ST elevation is indicative of STEMI and ST-segment depression in leads I, aVL, and V5–6 is indicative of non-ST-elevation ACS (NSTEMI-ACS).⁴² For the same reasons previously outlined for LBBB, patients with signs/symptoms that are highly suspicious for ongoing myocardial ischaemia and RBBB should be triaged for reperfusion therapy. In patients with a lower degree of clinical suspicion, the use of hs-cTn at presentation is recommended. Less than 40% of patients presenting to the ED with chest discomfort and RBBB will be found to have a final diagnosis of MI.^{37,42}

Comparison of current and previous ECG tracings can be valuable in the setting of suspected ACS, particularly in patients with pre-existing ECG abnormalities. In cases of suspected ACS with persistent or recurrent symptoms or in cases where there is diagnostic uncertainty, it is recommended to obtain serial 12-lead ECGs. [Figure S2](#) demonstrates electrocardiographic abnormalities in patients with STEMI and some other ECG findings that may prompt triage for an immediate reperfusion therapy if present.




Figure S1 Symptoms at presentation in acute coronary syndrome in women and men. ACS, acute coronary syndrome.






ECG pattern	Criteria	Signifying	Figure
i STEMI	New ST-elevation at the J-point in ≥ 2 contiguous leads ^a ≥ 2.5 mm in men <40 years, ≥ 2 mm in men ≥ 40 years, or ≥ 1.5 mm in women regardless of age in leads V2–V3 and/or ≥ 1 mm in the other leads (in the absence of LV hypertrophy or left bundle branch block) ^a Including V3R and V4R	Ongoing acute coronary artery occlusion	
ii Posterior STEMI	ST-segment depression in leads VI–V3, especially when the terminal T-wave is positive (ST-segment elevation equivalent), and concomitant ST-segment elevation ≥ 0.5 mm recorded in leads V7–V9	Posterior STEMI	
iii LCx occlusion/ right ventricular MI	ST-segment elevation in V7–V9 and V3R and V4R, respectively	Left circumflex (LCX) artery occlusion or right ventricular MI	
iv Multivessel ischaemia/ left main obstruction	ST depression ≥ 1 mm in six or more surface leads (inferolateral ST depression), coupled with ST-segment elevation in aVR and/or VI	Multivessel ischaemia or left main coronary artery obstruction, particularly if the patient presents with haemodynamic compromise	
v Left bundle branch block/ paced rhythm	QRS duration greater than 120 ms Absence of Q wave in leads I, V5 and V6 Monomorphic R wave in I, V5 and V6 ST and T wave displacement opposite to the major deflection of the QRS complex	Patients with a high clinical suspicion of ongoing myocardial ischaemia should be managed in a similar way to STEMI patients	
vi Right bundle branch block	QRS duration greater than 120 ms rsR' "bunny ear" pattern in the anterior precordial leads (leads V1–V3) Slurred S waves in leads I, aVL and frequently V5 and V6	Patients with a high clinical suspicion of ongoing myocardial ischaemia should be managed in a similar way to STEMI patients	

Figure S2 Electrocardiographic abnormalities in patients with STEMI and ECG findings that, if present, may prompt triage for immediate reperfusion therapy. ECG, electrocardiogram; STE-ACS, ST-elevation acute coronary syndrome; STEMI, ST-elevation myocardial infarction.

3.2.2. Acute coronary syndrome without persistent ST-segment elevation or other signs of acute vessel occlusion

ST-segment depression is not only a qualitative, but also a quantitative marker of risk. Both the number of leads with ST-segment depression and the magnitude of ST-segment depression (either within a single lead or the sum of all leads) are indicative of the extent of ischaemia and correlate with prognosis.^{43–45} While the prognostic impact of ST-segment depression is indisputable, the evidence regarding the prognostic impact of isolated T wave inversion is conflicting. T wave inversion was only independently predictive for an adverse outcome when occurring in ≥ 5 –6 leads, with no correlation found for T wave inversion occurring in fewer leads.^{46–50} Another relevant issue in this regard is that the interpretation of the prognostic value of T wave inversion may be hindered by inconsistent definitions. Overall, the prognostic value of T wave inversion certainly appears to be less than that of ST-segment depression, and the presence of concomitant T wave inversion does not alter the prognostic value of associated ST-segment depression.⁵⁰ The presence of ST-segment depression >1 mm in ≥ 6 leads in conjunction with ST-segment elevation in aVR and/or V1 is suggestive of multivessel ischaemia or severe left main coronary artery stenosis, particularly if the patient presents with haemodynamic compromise.^{51–53}

Beyond ST-segment deviation and T wave inversion, additional ECG patterns that may signify severe stenosis or even occlusion of the proximal left anterior descending (LAD) coronary artery have been described. However, the majority of these ECG patterns were identified in old, small, single-centre series. As a result, their true frequency and diagnostic yield remains uncertain. Up to a quarter of patients presenting with NSTEMI-ACS may have a totally occluded vessel on angiography, which is associated with increased mortality.^{54,55} Therefore, the recognition of ECG patterns (in the absence of ST-segment elevation) that may be associated with a totally occluded vessel on angiography is potentially clinically important. An abnormal ST-segment and T wave morphology, now known as Wellens' syndrome, was described in the early 1980s (Figure S3).⁵⁶ In a series of 1260 patients hospitalized for unstable angina (UA) between July 1980 and December 1985, 204 (16%) had this ECG pattern.⁵⁷ After excluding patients with recent MI and missing data, 180 patients were further analysed. All of these patients had stenosis of $\geq 50\%$ in the proximal LAD and 18% had a total occlusion. The type A pattern was present in 25% and the type B pattern in 75% of patients. In 2008, de Winter *et al.* reported another abnormal ST-segment and T wave morphology signifying proximal LAD occlusion.⁵⁸ Figure S3 demonstrates some of the potential electrocardiographic abnormalities in patients with NSTEMI-ACS. Novel ECG algorithms using digital ECG data are in development and offer the potential to improve diagnosis and risk stratification.^{59–62}




Figure S3 Electrocardiographic abnormalities in patients with non-ST-segment elevation acute coronary syndrome. ECG, electrocardiogram; LAD, left anterior descending artery. This figure highlights some of the electrocardiographic abnormalities that may be present in patients with NSTEMI-ACS.

3.3. Diagnostic tools | Biomarkers

3.3.1. High-sensitivity cardiac troponin

Table S2 Clinical implications of high-sensitivity cardiac troponin assays

Compared with standard cardiac troponin assays, hs-cTn assays:
• Have a higher NPV for acute MI, especially for patients who present early. • Reduce the ‘troponin-blind’ interval, leading to earlier detection of MI. • Result in a ~4% absolute and ~20% relative increase in the detection of Type 1 MI and a corresponding decrease in the diagnosis of unstable angina. • Are associated with a two-fold increase in the detection of Type 2 MI.
Levels of hs-cTn should be interpreted as quantitative markers of cardiomyocyte damage (i.e. the higher the level, the greater the likelihood of MI):
• The terms positive and negative troponin levels should be avoided: instead, elevated and non-elevated troponin levels are preferred. • Elevations beyond five-fold the upper reference limit have high (>90%) PPV for acute Type 1 MI. • Elevations up to three-fold the upper reference limit have only limited (50–60%) PPV for MI and may be associated with a broad spectrum of conditions. • It is common to detect circulating levels of cTn in healthy individuals. • Due to their high sensitivity, cTn levels can be elevated due to acute and chronic conditions other than MI.
Rising and/or falling cTn levels differentiate acute MI from chronic but not acute myocardial injury

cTn, cardiac troponin; hs-cTn, high-sensitivity cardiac troponin; MI, myocardial infarction; NPV, negative predictive value; PPV, positive predictive value.

3.3.1.1 Reasons for high-sensitivity cardiac troponin elevations other than Type 1 myocardial infarction

Many cardiac pathologies other than Type 1 MI can also result in cardiomyocyte injury and therefore cardiac troponin (cTn) elevations (Table S3). In these cases, myocardial injury can be related not only to acute myocardial ischaemia due to oxygen supply/demand imbalance related to coronary spasm, dissection, or micro-embolism, but also to tachy-/brady-arrhythmias, hypotension, or shock and respiratory failure (Type 2 MI). In addition, myocardial injury can be related to causes other than acute myocardial ischaemia. These causes include both cardiac (heart failure [HF], hypertensive emergencies, myocarditis, takotsubo syndrome, and valvular heart disease) and non-cardiac conditions (critical illness, chronic kidney disease, stroke, pulmonary embolism) (Table S3).⁵ Patients with suspected ACS therefore require careful clinical evaluation and consideration of these differential diagnoses, which frequently require different treatments.

In patients with elevations in cTn without a specific diagnosis, these elevations should not be primarily attributed to impaired clearance and considered harmless. Cardiac conditions seem to be the most important contributors to cTn elevation in this setting, including stable CAD, HF, and hypertensive heart disease.^{31,63} Other life-threatening conditions can present with chest pain and may result in elevated cTn concentrations, including aortic dissection and pulmonary embolism.

Table S3 Conditions other than acute Type 1 myocardial infarction associated with cardiomyocyte injury (i.e. cardiac troponin elevation)

Myocardial injury related to acute myocardial ischaemia because of oxygen supply/demand imbalance (Type 2 MI)
Reduced myocardial perfusion, e.g.:
• Coronary artery spasm, microvascular dysfunction • Coronary embolism • Non-atherosclerotic coronary artery dissection • Sustained bradyarrhythmia • Hypotension or shock • Respiratory failure • Severe anaemia
Increased myocardial oxygen demand, e.g.:
• Sustained tachyarrhythmia • Severe hypertension with or without left ventricular hypertrophy
Other causes of myocardial injury
Cardiac conditions:
• Heart failure • Myocarditis^a • Cardiomyopathy (any type) • Takotsubo syndrome • Cardiac contusion or cardiac procedures (CABG, PCI, valvular interventions, ablation, pacing, cardioversion, or endomyocardial biopsy)
Systemic conditions:
• Sepsis, infectious disease • Chronic kidney disease • Stroke, subarachnoid haemorrhage • Pulmonary embolism, pulmonary hypertension • Infiltrative diseases (e.g. amyloidosis, sarcoidosis, haemochromatosis, scleroderma) • Myocardial drug toxicity or poisoning (e.g. doxorubicin, 5-fluorouracil, trastuzumab, snake venoms) • Critically ill patients • Hypo- and hyper-thyroidism • Strenuous exercise • Rhabdomyolysis

CABG, coronary artery bypass grafting; MI, myocardial infarction; PCI, percutaneous coronary intervention.

^aIncludes myocardial extension of endocarditis or pericarditis.

As such, these conditions should also be considered as differential diagnoses.

While Type 1 MI is due to atherosclerotic CAD with plaque rupture/erosion and either occlusive or non-occlusive thrombus, Type 2 MI is due to an oxygen supply/demand imbalance and can have multiple causes (see Section 12.1 in the main text). The observed incidence of Type 2 MI is increasing with the implementation of hs-cTn assays.⁸ Treatments differ substantially between Type 2 and Type 1 MI and their early and accurate non-invasive differentiation before angiography remains a gap in evidence. Recent studies have proposed different scores, including clinical parameters and biomarkers, to better differentiate these entities before angiography.^{7,8,64–66}

© ESC 2023

© ESC 2023

3.3.2. Rapid 'rule-in' and 'rule-out' algorithms

3.3.2.1. European Society of Cardiology 0 h/1 h and European Society of Cardiology 0 h/2 h algorithms

NSTEMI can be ruled out at presentation if the 0 h hs-cTn concentration is very low and the chest pain onset was >3 h prior to the 0 h hs-cTn measurement. NSTEMI can also be ruled out by the combination of low baseline levels of hs-cTn and the lack of a relevant increase within 1 h (no 1 hΔ). Patients have a high likelihood for NSTEMI if the hs-cTn concentration at presentation is at least moderately elevated or shows a clear rise within the first hour (1 hΔ).^{4,6–8,89–96} Cut-offs are assay-specific (see Table S4) and derived to meet pre-defined criteria for sensitivity and specificity for NSTEMI.

Recently, specific cut-offs for the patients assigned to the 'observe zone' using the hs-cTn T assay (combination of a 3 h hs-cTn T concentration <15 ng/L and a 0 h/3 h absolute change <4 ng/L) have been derived and validated as having acceptable safety and efficacy for further decision-making.⁹⁷ Specific cut-offs for other hs-cTn I assays in the observe zone are currently being developed.

3.3.2.2. Caveats of using rapid algorithms

When using any algorithm, six main caveats apply:

- (i) Algorithms should only be used in conjunction with all available clinical information, including detailed assessment of chest pain characteristics and ECGs, and should be applied only following exclusion of STEMI or other life-threatening conditions. Patients with a clear pattern of crescendo or UA should undergo further investigation.
- (ii) The rapid algorithms should be used only in patients presenting with suspected ACS and should not be applied in an unselected ED population (i.e. in patients with stroke or sepsis).
- (iii) The European Society of Cardiology (ESC) 0 h/1 h and 0 h/2 h algorithms apply to all patients irrespective of chest pain onset. This approach is very safe (negative predictive value [NPV] and sensitivity >99%), including in the subgroup of patients presenting very early (e.g. <2 h).⁷¹ However, due to the time dependency of cTn release and the moderate number of patients presenting <1 h after chest pain onset in previous studies, obtaining an additional cTn concentration at 3 h in early presenters triaged towards rule-out should be considered.
- (iv) As late increases in cTn have been described in ~1% of patients, serial cTn testing should also be pursued if clinical suspicion remains high or if the patient develops recurrent chest pain.^{30,31,68,71–74}
- (v) Time to decision = time of blood draw + turnaround time. The use of the ESC 0 h/1 h algorithms is irrespective of the local turnaround time (time from blood draw to blood results); 0 h and 1 h refer to the time points at which blood is taken. The second blood draw may need to be taken before the result from the first one is available (although the results should be available in most cases within 60 min of blood sampling), but this does not affect the interpretation of the algorithms. The clinical and economic benefit of the ESC 0 h/1 h algorithm compared with other algorithms where the second blood draw is later than 1 h is therefore independent of the local turnaround time.⁹⁸
- (vi) The ESC 0 h/1 h and 0 h/2 h algorithms are assay specific and can be used only for the suggested assays for which the algorithms have been validated. If none of these assays are available, an alternative strategy needs to be considered.

Table S4 Assay specific cut-off levels in ng/L within the 0 h/1 h and 0 h/2 h algorithms

0 h/1 h algorithm	Very low	Low	No 1 hΔ	High	1 hΔ
hs-cTnT (Elecys; Roche)	<5	<12	<3	≥52	≥5
hs-cTnI (Architect; Abbott)	<4	<5	<2	≥64	≥6
hs-cTnI (Centaur; Siemens)	<3	<6	<3	≥120	≥12
hs-cTnI (Access; Beckman Coulter)	<4	<5	<4	≥50	≥15
hs-cTnI (Clarity; Singulex)	<1	<2	<1	≥30	≥6
hs-cTnI (Vitros; Clinical Diagnostics)	<1	<2	<1	≥40	≥4
hs-cTnI (Pathfast; LSI Medience)	<3	<4	<3	≥90	≥20
hs-cTnI (TriageTrue; Quidel)	<4	<5	<3	≥60	≥8
hs-cTnI (Dimension EXL; Siemens)	<9	<9	<5	≥160	≥100
0 h/2 h algorithm	Very low	Low	No 2 hΔ	High	2 hΔ
hs-cTnT (Elecys; Roche)	<5	<14	<4	≥52	≥10
hs-cTnI (Architect; Abbott)	<4	<6	<2	≥64	≥15
hs-cTnI (Centaur; Siemens)	<3	<8	<7	≥120	≥20
hs-cTnI (Access; Beckman Coulter)	<4	<5	<5	≥50	≥20
hs-cTnI (Clarity; Singulex)	<1	TBD	TBD	≥30	TBD
hs-cTnI (Vitros; Clinical Diagnostics)	<1	TBD	TBD	≥40	TBD
hs-cTnI (Pathfast; LSI Medience)	<3	TBD	TBD	≥90	TBD
hs-cTnI (TriageTrue; Quidel)	<4	TBD	TBD	≥60	TBD

The cut-offs apply irrespective of age, sex, and renal function. Optimized cut-offs for patients above 75 years of age and patients with renal dysfunction have been evaluated, but not consistently shown to provide better balance between safety and efficacy as compared with these universal cut-offs.^{30,31} The algorithms for additional assays are in development: hs-cTn T on Elecys (Roche), hs-cTn I on Architect (Abbott), hs-cTn I on Centaur (Siemens), hs-cTn I on Access (Beckman Coulter), hs-cTn I on Clarity (Singulex), hs-cTn I on Vitros (Clinical Diagnostics), hs-cTn I on Pathfast (LSI Medience), and hs-cTn I on TriageTrue (Quidel).
hs-cTn, high-sensitivity cardiac troponin; TBD, to be determined.^{30,31,67–88}

3.3.2.3. Practical guidance on how to implement the European Society of Cardiology 0 h/1 h algorithm

Documentation of the time of the 0 h blood draw allows exact determination of the time window (± 10 min) of the 1 h blood draw. If for whatever reason the 1 h (± 10 min) blood draw was not feasible, then blood should be drawn at 2 h and the ESC 0 h/2 h algorithm applied (if validated for the assay in use, please see [Table S4](#) for further details).

3.3.3. Other biomarkers

Compared with cTn, the creatine kinase myocardial band isoenzyme shows a more rapid decline after MI and may provide added value for assessment of the timing of myocardial injury and for the detection of early re-infarction.⁴

Myosin-binding protein C is more abundant than cTn and may therefore provide clinical value as an alternative to or in combination with cTn.⁹⁹ Assessment of copeptin, the C-terminal part of the vasopressin pro-hormone, may quantify the endogenous stress level in multiple medical conditions, including MI. As the level of endogenous stress appears to be high at the onset of MI in most patients, the added value of copeptin to lower-sensitivity cTn assays is substantial.^{100–102} Therefore, the routine use of copeptin as an additional biomarker for the early rule-out of MI may be most valuable in the increasingly uncommon setting where hs-cTn assays are not available. However, copeptin does not have relevant added value for institutions using one of the well-validated hs-cTn-based rapid protocols for the early diagnosis of MI.^{103–111}

3.4. Diagnostic tools | Non-invasive imaging

3.4.1. Cardiac magnetic resonance imaging with or without stress testing

The CARMENTA (CARDiovascular Magnetic rEsoNance imaging and computed Tomography Angiography) trial compared a cardiac magnetic resonance (CMR)- or coronary computed tomography angiography (CCTA)-first strategy with a control strategy of routine clinical care in 207 patients (age 64 years; 62% male patients) with acute chest pain, elevated hs-cTn T levels (> 14 ng/L), and an inconclusive ECG.¹¹² Follow-up invasive coronary angiography (ICA) was recommended if the initial CMR or CCTA suggested myocardial ischaemia, infarction, or obstructive CAD ($\geq 70\%$ stenosis). The CMR- and CCTA-first strategies reduced ICA

compared with routine clinical care (87% [$P = 0.001$], 66% [$P < 0.001$], and 100%, respectively), with no differences in major adverse cardiac events (MACE) at 1 year. Obstructive CAD disclosed by ICA occurred in 61% of patients in the routine clinical care arm, in 69% in the CMR-first arm ($P = 0.308$ vs. routine), and in 85% in the CCTA-first arm ($P = 0.006$ vs. routine). In the non-CMR and non-CCTA arms, follow-up CMR and CCTA were performed in 67% and 13% of patients, leading to a new diagnosis in 33% and 3%, respectively ($P < 0.001$). CARMENTA found that implementing CMR or CCTA first in the diagnostic process for patients with NSTEMI safely reduces the need for ICA and is diagnostically useful for identifying alternative diagnoses as compared with management guided by CCTA or invasive angiography.

3.4.2. Single-photon emission computerized tomography perfusion imaging and stress echocardiography

In patients in the observe zone with a normal ECG and non-elevated hs-cTn levels and who have been pain free for several hours, stress echocardiography or single-photon emission computerized tomography (SPECT) imaging can be performed safely as an alternative to CCTA during hospitalization or shortly after discharge. A limitation of these techniques is that they do not detect non-obstructive plaque. Stress echocardiography or SPECT imaging are preferred over exercise ECG due to their greater diagnostic accuracy and prognostic value.^{113,114}

Stress echocardiography can detect regional wall motion abnormalities associated with acute myocardial ischaemia. If the acoustic windows are not adequate to assess regional wall motion abnormalities, the use of echocardiographic contrast is recommended to improve the detection of wall motion abnormalities. Various studies have shown that normal exercise, dobutamine or dipyridamole stress echocardiograms have high NPV for ischaemia and are associated with good patient outcomes.^{115,116}

3.5. Differential diagnosis for acute chest pain

Some conditions that should be considered in the differential diagnosis of ACS are shown in [Table S5](#). These include aortic dissection, pulmonary embolism, and tension pneumothorax. Takotsubo syndrome is an increasingly recognized differential diagnosis and usually requires ICA in order to rule out ACS.¹¹⁷

Table S5 Differential diagnoses of acute coronary syndrome in the setting of acute chest pain

Cardiac	Pulmonary	Vascular	Gastrointestinal	Orthopaedic	Other
Myocarditis/pericarditis, cardiomyopathies ^a	Pulmonary embolism	Aortic dissection	Oesophagitis, reflux, or spasm	Musculoskeletal disorders	Anxiety disorders
Tachyarrhythmias	(Tension) Pneumothorax	Symptomatic aortic aneurysm	Peptic ulcer, gastritis	Chest trauma	Herpes zoster
Acute heart failure	Bronchitis, pneumonia	Stroke	Pancreatitis	Muscle injury/inflammation	Anaemia
Hypertensive emergencies	Pleuritis		Cholecystitis	Costochondritis	
Aortic valve stenosis				Cervical spine pathologies	
Takotsubo syndrome					
Coronary spasm					
Cardiac trauma					

^aDilated, hypertrophic, and restrictive cardiomyopathies may cause angina or chest discomfort.

Chest X-ray is recommended in all patients in whom NSTEMI-ACS is considered unlikely in order to detect/exclude potential differential diagnoses, including pneumonia, pneumothorax, rib fractures, or other thoracic disorders. Stroke may be accompanied by ECG changes, myocardial wall motion abnormalities, and cardiomyocyte injury (resulting in an increase in cTn concentrations). The majority of patients presenting to the ED with acute chest pain have non-cardiac conditions causing the chest discomfort.^{30,31,68,70,71,118–123} In many instances the pain is musculoskeletal, and this is generally benign, self-limiting, and does not require hospitalization. Chest pain characteristics may help to some extent in the early identification of these patients.

4. Initial measures for patients presenting with suspected acute coronary syndrome | Initial treatment

4.1. Pre-hospital logistics of care

Some special considerations may be relevant when making decisions about invasive management. This includes decisions regarding invasive assessment in severely frail patients with comorbid conditions (e.g. advanced cognitive impairment or advanced cancer), in whom the invasive procedure may be more likely to lead to harm than benefit. Considerations for end-of-life care have crucial importance in the pre-hospital setting and should involve a multidisciplinary decision-making process, taking account of patient and carer preferences.

4.1.1. Organization of ST-elevation myocardial infarction treatment in networks

Table S6 Key features of networks of care for the pre-hospital management of STEMI

• Clear definition of geographic areas of responsibility.
• Shared protocols, based on risk stratification and transportation by a trained physician, nurse, or paramedic staff in appropriately equipped ambulances or helicopters.
• Pre-hospital triage of patients with a STEMI working diagnosis, according to risk stratification (ECG and symptoms), to the appropriate centre, bypassing non-PCI hospitals or hospitals without a 24/7 service.
• On arrival at the appropriate hospital, the patient with suspected STEMI should be immediately taken to the catheterization laboratory, bypassing the ED and other ward areas.
• Patients presenting to a non-PCI-capable hospital and awaiting transportation for primary/rescue PCI must be attended to in an appropriately monitored and staffed area.
• If the working diagnosis of STEMI has not been made by the ambulance crew and the ambulance arrives at a non-PCI-capable hospital, the ambulance should wait for the diagnosis and, if a working diagnosis of STEMI is made, should continue to a PCI-capable hospital.

ECG, electrocardiogram; ED, emergency department; PCI, percutaneous coronary intervention; STEMI, ST-elevation myocardial infarction.

To maximize staff experience, PCI centres should perform invasive management acutely on a 24/7 basis for all ACS patients. Other models, although not ideal, may include weekly or daily rotation of PPCI centres or multiple acute centres in the same region. Hospitals that cannot offer a 24/7 service for PCI should be allowed to perform urgent invasive management (angiography and PCI if needed) in patients already admitted for another reason who develop ACS during their hospital stay. All of these hospitals should comply with contemporary guidelines for providing emergency medical services (EMS) for STEMI and NSTEMI-ACS. Since urgent or emergency coronary artery bypass grafting (CABG) may be indicated, the organizational network should designate one or more centres for acute cardiac surgery.

Hospitals should be discouraged from initiating a service limited to daytime- or within-hours acute ACS PCI, as this may generate confusion with EMS operators and affect both the ACS diagnosis-to-revascularization time and the quality of intervention at a focused 24/7 PPCI centre.

5. Acute-phase management of patients with acute coronary syndrome

5.1. Acute coronary syndrome managed with invasive strategy

In several trials, a pre-specified subgroup analysis of patients with a Global Registry of Acute Coronary Events (GRACE) risk score >140 showed that they benefitted from an early invasive strategy as opposed to those with a GRACE risk score <140 (TIMACS [Timing of Intervention in Acute Coronary Syndromes] trial: hazard ratio (HR) 0.65, 95% confidence interval (CI), 0.48–0.89 vs. HR 1.12, 95% CI, 0.81–1.56; $P_{interaction}$ = 0.01; VERDICT [Very Early vs. Deferred Invasive evaluation using Computerized Tomography] trial: HR, 0.81, 95% CI, 0.67–1.00 vs. HR 1.21, 95% CI, 0.92–1.60; $P_{interaction}$ = 0.02).^{124,125} A significant interaction was found between timing of invasive angiography and GRACE score in terms of the risk of death: a trend toward decreased risk of all-cause mortality was seen with an early invasive strategy in patients with a GRACE score risk >140 (HR 0.83, 95% CI, 0.63–1.10), with a higher risk of all-cause mortality in patients with a GRACE risk score ≤140 (HR 2.04, 95% CI, 1.16–3.59).¹²⁶ In addition, an early invasive strategy reduced the risk of all-cause mortality in patients with dynamic ECG changes, higher heart rate, and lower systolic blood pressure (SBP).¹²⁶ It should be highlighted that both randomized controlled trials (RCTs) calculated the original GRACE risk score for in-hospital death.¹²⁷ Due to different weighting of variables, scores other than GRACE may be considerably different for the same patient, possibly leading to different treatment decisions.¹²⁶ Elevated cardiac biomarkers are part of the GRACE risk score, and therefore, there are some cases in which a patient can present with elevated troponin but a low GRACE score (i.e. <140). The evidence supporting the clinical benefit of routine early invasive strategy in these patients is less robust. Most recent trials in this area enrolled a large proportion of patients (around 80%) with elevated troponin, making the evaluation of this issue more complex. In a recent analysis of the VERDICT trial (which excluded patients with transient ST-segment elevation), a significant interaction between timing of ICA and GRACE score, with regard to the risk of death, was found but not with raised troponin.¹²⁶ Another important point is that the GRACE risk score may overestimate risk in ethnic minorities with NSTEMI.¹²⁸

5.2. Patients not undergoing reperfusion

ACS patients not receiving reperfusion/revascularization treatment represent a heterogeneous group, which includes patients not undergoing coronary angiography, patients with extensive CAD not amenable to revascularization, and patients without obstructive CAD.

5.2.1. Patients who are not candidates for invasive coronary angiography

These patients represent a small subgroup for which data indicating a hypothetical advantage of an invasive strategy are scarce. Advanced age, female sex, chronic kidney disease (CKD), diabetes mellitus, prior HF/revascularization, history of cancer, and frailty are commonly reported reasons for not performing diagnostic angiography in these patients.^{129–132} These features largely overlap with the predictors of bleeding and ischaemic adverse events and this may explain the poor prognosis of this population.

Medical management without invasive assessment should only be chosen after careful risk assessment, bearing in mind that coronary angiography using the radial approach is a relatively low-risk procedure, that impaired left ventricular (LV) function increases mortality risk, and that knowledge of the coronary anatomy may impact on both risk stratification and the choice of pharmacological therapy. Advanced age or female sex alone, in the absence of severe comorbidities or frailty, should not be considered sufficient reasons not to perform ICA and, likewise, ICA should not be denied for logistical reasons.¹³³

5.2.2. Patients with coronary artery disease not amenable to revascularization

Patients diagnosed with severe CAD who are not amenable to any type of revascularization are at very high risk of recurrent ischaemic events.¹³⁴ These patients are frequently female, elderly, and/or suffering from severe CKD, have multivessel CAD, and a history of prior MI/revascularization. The decision not to perform PCI is an independent predictor of increased CV mortality, both in hospital and long-term.^{131,135} Accordingly, the decision not to perform revascularization should only be made in very selected patients, when there is a consensus that the risk of revascularization outweighs the benefit for clinical or anatomical reasons. These patients should undergo an aggressive secondary prevention treatment with potent antiplatelet therapy and anti-anginal agents, taking their comorbidities into account.¹³⁶

6. Antithrombotic therapy

6.1. Long-term treatment

A strategy consisting of Factor Xa inhibition with a very low dose of rivaroxaban (2.5 mg b.i.d. [*bis in die*, twice a day]) plus dual antiplatelet therapy (DAPT; aspirin plus clopidogrel) was studied in the ATLAS ACS 2-TIMI 51 (Anti-Xa Therapy to Lower Cardiovascular Events in Addition to Standard Therapy in Subjects With Acute Coronary Syndrome ACS 2-Thrombolysis In Myocardial Infarction 51) trial on a background of clopidogrel treatment.¹³⁷ The study showed a reduction

of ischaemic events and CV mortality along with a higher risk of major and intracranial bleeding. However, data are lacking regarding this strategy on a background of ticagrelor or prasugrel treatment. Therefore, it is difficult to extrapolate these trial results to contemporary practice with the use of potent P2Y₁₂ receptor inhibitors. A similar approach was investigated in the APPRAISE-2 (Apixaban for Prevention of Acute Ischemic Events 2) trial in patients with recent ACS and at least two additional risk factors for recurrent ischaemic events. In this study, apixaban at a dose of 5 mg b.i.d. (mostly in combination with DAPT) was not associated with a lower risk of CV death, MI, or ischaemic stroke but was associated with a more than two-fold increase in major bleeding risk.¹³⁸

6.1.1. Prolonging antithrombotic therapy beyond 12 months

Prolonging DAPT: based on the results of the Dual Antiplatelet Therapy (DAPT) and PrEvention with TicaGrelor of SecondAry Thrombotic Events in High-RiSk Patients with Prior AcUte Coronary Syndrome—Thrombolysis In Myocardial Infarction Study Group (PEGASUS-TIMI) 54 trials, a prolonged DAPT course >12 months should be considered in those with high thrombotic risk and without an increased risk of major or life-threatening bleeding, and may be considered in ACS patients with moderately elevated thrombotic risk who have tolerated DAPT without a bleeding complication (see [Figure S4](#); [Table S7](#) and [S8](#)).^{139,140} Of note, the 60 mg b.i.d. dose for ticagrelor was associated with reduced bleeding compared with the 90 mg b.i.d. dose and should be preferred for extended therapy >12 months.^{141,142}

Dual antithrombotic therapy: based on the Cardiovascular Outcomes for People Using Anticoagulation Strategies (COMPASS) trial in chronic coronary syndrome (CCS) patients (62% with a history of MI), a strategy of dual antithrombotic therapy (DAT)—consisting of Factor Xa inhibition with a very low dose of rivaroxaban (2.5 mg b.i.d.) in combination with aspirin—should be considered as an option for maintenance treatment beyond 12 months post-ACS in patients at high thrombotic risk and without an increased risk of major or life-threatening bleeding, and may be considered in patients with moderately elevated thrombotic risk.^{143,144}

In the COMPASS trial, 27 395 stable CAD patients were randomized (62% with a history of MI, 7.1 years before enrolment) to one of three arms: a very low dose of rivaroxaban (2.5 mg b.i.d.) plus aspirin, rivaroxaban (5 mg b.i.d.) plus an aspirin-matched placebo, or aspirin alone. The combination of rivaroxaban 2.5 mg b.i.d. plus aspirin resulted in a significant reduction in the risk of the combined ischaemic endpoint, overall mortality (without reaching the threshold *P*-value according to the Hochberg procedure), and CV mortality alone, but increased the risk of major bleeding complications without a significant increase in the risk of fatal, intracranial, or critical organ bleeding events. Rivaroxaban 5 mg b.i.d. did not result in improved CV outcomes in comparison to aspirin monotherapy but did result in an increase in bleeding events.

The available evidence for DAT in ACS-CABG patients comes from subgroup analyses of ACS trials that had results consistent with the overall findings.^{130,145,146}




Figure S4 Antithrombotic strategies beyond the first 12 months after ACS. ACS, acute coronary syndrome; DAPT, dual antiplatelet therapy; DAT, dual antithrombotic therapy; HBR, high bleeding risk; MD, maintenance dose; o.d., once a day.

Table S7 Treatment options for extended dual antithrombotic or antiplatelet therapies

Drug	Dose	Indication	NNT (ischaemic outcomes)	NNH (bleeding outcomes)
<i>DAT regimens for extended treatment (including aspirin 75–100 mg o.d.)</i>				
Rivaroxaban (COMPASS trial)	2.5 mg b.i.d.	Patients with CAD or symptomatic PAD at high risk of ischaemic events	77	84
<i>DAPT regimens for extended treatment (including aspirin 75–100 mg o.d.)</i>				
Clopidogrel (DAPT trial)	75 mg/d	Post-MI in patients who have tolerated DAPT for 1 year	63	105
Prasugrel (DAPT trial)	10 mg/d (5 mg/d if body weight <60 kg or age >75 years)	Post-PCI for MI in patients who have tolerated DAPT for 1 year	63	105
Ticagrelor (PEGASUS-TIMI 54)	60/90 mg b.i.d. ^a	Post-MI in patients who have tolerated DAPT for 1 year	84	81

Drugs (in addition to aspirin 75–100 mg/d) for extended DAPT treatment options are in alphabetical order. NNT refers to the primary ischaemic endpoints of the respective trials and NNH refers to the key safety (bleeding) endpoints. NNT and NNH numbers from the DAPT trial are pooled numbers for clopidogrel and prasugrel.

b.i.d., *bis in die* (twice a day); CAD, coronary artery disease; COMPASS, Cardiovascular Outcomes for People using Anticoagulation Strategies; DAPT, dual antiplatelet therapy; DAT, dual antithrombotic therapy; MI, myocardial infarction; NNH, number needed to harm; NNT, number needed to treat; o.d., once a day; PAD, peripheral arterial disease; PCI, percutaneous coronary intervention; PEGASUS-TIMI 54, Prevention of Cardiovascular Events in Patients With Prior Heart Attack Using Ticagrelor Compared to Placebo on a Background of Aspirin-Thrombolysis In Myocardial Infarction 54.

^aThe 60 mg b.i.d. dose for ticagrelor was associated with reduced bleeding compared with the 90 mg b.i.d. dose and should be preferred for extended therapy >12 months.

Table S8 Risk criteria for extended treatment with a second antithrombotic agent

High thrombotic risk (Class IIa)	Moderate thrombotic risk (Class IIb)
Complex CAD and at least one criterion	Non-complex CAD and at least one criterion
Risk enhancers	
Diabetes mellitus requiring medication History of recurrent MI Any multivessel CAD Premature (<45 years) or accelerated (new lesion within a 2-year timeframe) CAD Concomitant systemic inflammatory disease (e.g. human immunodeficiency virus, systemic lupus erythematosus, chronic arthritis) Polyvascular disease (CAD plus PAD) CKD with eGFR 15–59 mL/min/1.73 m ²	Diabetes mellitus requiring medication History of recurrent MI Polyvascular disease (CAD plus PAD) CKD with eGFR 15–59 mL/min/1.73 m ²
Technical aspects	
At least three stents implanted At least three lesions treated Total stent length >60 mm History of complex revascularization (left main, bifurcation stenting with ≥2 stents implanted, chronic total occlusion, stenting of last patent vessel) History of stent thrombosis on antiplatelet treatment	

CAD, coronary artery disease; CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; MI, myocardial infarction; PAD, peripheral arterial disease.

In line with guideline recommendations, CAD patients are stratified into two different risk groups (high vs. moderately increased thrombotic or ischaemic risk). Stratification of patients towards complex vs. non-complex CAD is based on individual clinical judgment with knowledge of the patient's cardiovascular history and/or coronary anatomy. Selection and composition of risk-enhancing factors are based on the combined evidence of clinical trials on extended antithrombotic treatment in CAD patients and on data from related registries.¹⁴¹

Table S9 High-risk features of stent-driven recurrent ischaemic events

Prior stent thrombosis on adequate antiplatelet therapy
Stenting of the last remaining patent coronary artery
Diffuse multivessel disease, especially in patients with diabetes
Chronic kidney disease (i.e. creatinine clearance <60 mL/min)
At least three stents implanted
At least three lesions treated
Bifurcation with two stents implanted
Total stent length >60 mm
Treatment of a chronic total occlusion

6.2. Antiplatelet therapy in patients requiring oral anticoagulation

Dual therapy with an oral anticoagulant (OAC) and one antiplatelet agent (aspirin or clopidogrel) may be considered beyond 1 year in patients at very high risk of coronary events, as defined in [Table S9](#).

6.2.1. Acute coronary syndrome patients with atrial fibrillation

Single antiplatelet therapy with clopidogrel was first evaluated in the WOEST (What is the Optimal antiplatelet and Anticoagulant Therapy in Patients With Oral Anticoagulation and Coronary StenTing) trial, where 573 patients were randomized to DAT with an oral anticoagulant (OAC) and clopidogrel or to TAT with an OAC, clopidogrel, and aspirin 80–100 mg/day.¹⁴⁷ Treatment was continued for 1 month after bare-metal stenting (35% of patients) and for 1 year after drug-eluting stent (DES) placement (65% of patients). PCI was performed while on vitamin K antagonist (VKA) in half of the patients and one-third of patients presented with NSTEMI-ACS. Femoral access was used in the majority of patients (74%). The primary endpoint of any TIMI (Thrombolysis In Myocardial Infarction) bleeding was significantly reduced in the DAT arm compared with the TAT arm, while no significant differences were observed in major bleeding. The rates of MI, stroke, target vessel revascularization (TVR), and stent thrombosis were comparable between the two groups, but all-cause mortality was lower in the DAT group at 1 year.

In the ISAR-TRIPLE (Triple Therapy in Patients on Oral Anticoagulation After Drug Eluting Stent Implantation) trial, 614 patients (one-third of whom presented with ACS) undergoing stenting and requiring OAC were randomly assigned to either 6 weeks or 6 months of clopidogrel therapy in addition to aspirin and a VKA.¹⁴⁸ The primary endpoint of death, MI, stent thrombosis, ischaemic stroke, or TIMI major bleeding at 9 months did not differ between the 6-week and 6-month TAT groups. The same was true for the combined incidence of death, MI, stent thrombosis, and ischaemic stroke. Furthermore, no significant difference in TIMI major bleeding was observed. Of note, 10% of patients in the WOEST trial and 7% of patients in the ISAR-TRIPLE trial had prosthetic heart valves. The subgroup analysis of WOEST showed that patients with prosthetic heart valves on DAT appeared to derive a similar benefit to the general population.

In PIONEER AF-PCI (A Study Exploring Two Strategies of Rivaroxaban and One of Oral Vitamin K Antagonist in Patients With Atrial Fibrillation Who Undergo Percutaneous Coronary Intervention), 2124 patients with atrial fibrillation (AF; [50% with ACS]) recently treated with stenting were randomized to rivaroxaban 15 mg once a day plus a P2Y₁₂ receptor inhibitor for 12 months

(group 1), rivaroxaban 2.5 mg twice a day plus DAPT for 1, 6, or 12 months (group 2), or standard therapy, consisting of a VKA plus DAPT for 1, 6, or 12 months (group 3).¹⁴⁹ The P2Y₁₂ receptor inhibitor used most frequently was clopidogrel and DAPT was continued for up to 12 months in 49% of patients. The primary endpoint of clinically significant bleeding events was significantly lower in the two groups receiving rivaroxaban than in the group receiving standard therapy. In ACS patients, the trend towards a reduced rate of clinically significant bleeding events was stronger in patients in group 2 than in group 1. The rates of death from CV causes, MI, or stroke were similar in the three groups. All-cause death or re-hospitalization was significantly reduced at 1 year in the two groups who received rivaroxaban compared with the group who received standard therapy.

The RE-DUAL PCI (Randomized Evaluation of Dual Antithrombotic Therapy with Dabigatran versus Triple Therapy with Warfarin in Patients with Nonvalvular Atrial Fibrillation Undergoing Percutaneous Coronary Intervention) trial studied 2725 patients (50% with ACS) recently treated with stenting.¹⁵⁰ Patients were randomized to TAT with VKA plus a P2Y₁₂ receptor inhibitor and aspirin (for 1–3 months) or DAT with dabigatran (110 mg or 150 mg b.i.d.) plus a P2Y₁₂ receptor inhibitor (no aspirin). The P2Y₁₂ receptor inhibitors used were primarily clopidogrel and ticagrelor (in 87% and 12% of patients, respectively). The primary endpoint of major or clinically relevant non-major bleeding occurred in 15.4% of the 110 mg DAT group compared with 26.9% of the TAT group, and 20.2% of the 150 mg DAT group compared with 25.7% of the corresponding TAT group. The trial also tested for the non-inferiority of DAT with dabigatran (both doses combined) to TAT with respect to the incidence of a composite efficacy endpoint of thrombo-embolic events (MI, stroke, or systemic embolism), death, or unplanned re-vascularization. The incidence of the composite efficacy endpoint was 13.7% in the two DAT groups combined compared with 13.4% in the TAT group. However, RE-DUAL PCI was underpowered for individual ischaemic endpoints, such as stent thrombosis, which occurred twice as often in the 110 mg DAT group compared with the TAT group, albeit at low absolute incidences.

AUGUSTUS (An Open-Label, 2 × 2 Factorial, Randomized Controlled, Clinical Trial to Evaluate the Safety of Apixaban Versus Vitamin K Antagonist and Aspirin Versus Aspirin Placebo in Patients With Atrial Fibrillation and Acute Coronary Syndrome or Percutaneous Coronary Intervention) randomized 4614 patients with AF recently treated with PCI or presenting with ACS to either apixaban (5 mg b.i.d.) or a VKA.¹⁵¹ The trial had a two-by-two factorial design, with a P2Y₁₂ receptor inhibitor administered to all patients for up to 6 months, while patients allotted to the apixaban or VKA groups were further randomized to either aspirin or placebo. The primary outcome of major or clinically relevant non-major bleeding occurred in 10.5% of the patients receiving apixaban, compared with 14.7% of those receiving a VKA, and in 16.1% of patients receiving aspirin, compared with 9.0% of those receiving placebo. Secondary outcomes included death or hospitalization and a composite of ischaemic events. Patients in the apixaban group had a lower incidence of death or hospitalization than those in the VKA group (23.5 vs. 27.4%), which was mostly driven by reduced hospitalization, and a similar incidence of death or ischaemic events. Patients in the aspirin group had a similar incidence of death or hospitalization and of death or ischaemic events in comparison to the placebo group. However, there was a trend towards lower rates of MI and stent thrombosis in the aspirin group (2.9% vs. 3.6% and 0.5% vs. 0.9%, respectively).

The ENTRUST-AF PCI (Edoxaban Treatment Versus Vitamin K Antagonist in Patients With Atrial Fibrillation Undergoing Percutaneous Coronary Intervention) trial randomized 1506 patients with AF successfully treated with PCI (50% presenting with ACS) to either edoxaban 60 mg daily plus a P2Y₁₂ receptor inhibitor or a VKA plus DAPT with a P2Y₁₂ receptor inhibitor and aspirin (for 1–12 months).¹⁵² The primary endpoint of major or clinically relevant non-major bleeding occurred in 17% of patients in the DAT group vs. 20% in the TAT group. The study showed non-inferiority for the primary endpoint but, in contrast to the other trials, the DAT strategy with edoxaban did not meet the criteria for superiority. Combined ischaemic events (CV death, stroke, systemic embolism, MI, or stent thrombosis) were not significantly different between the groups (7% vs. 6%).

6.3. Fibrinolysis and pharmaco-invasive strategy

Table S10 Doses of fibrinolytic agents and antithrombotic co-therapies

Drug	Initial treatment	Specific contraindications
Streptokinase	1.5 million units over 30–60 min i.v.	Previous treatment with streptokinase or anistreplase
Alteplase (tPA)	15 mg i.v. bolus 0.75 mg/kg i.v. over 30 min (up to 50 mg) then 0.5 mg/kg i.v. over 60 min (up to 35 mg)	
Reteplase (rPA)	10 units + 10 units i.v. bolus given 30 min apart	
Tenecteplase (TNK-tPA)	Single i.v. bolus: 30 mg (6000 U) if <60 kg 35 mg (7000 U) if 60 to <70 kg 40 mg (8000 U) if 70 to <80 kg 45 mg (9000 U) if 80 to <90 kg 50 mg (10 000 U) if ≥90 kg It is recommended to reduce to half dose in patients ≥75 years of age. ¹⁵³	
Doses of antiplatelet co-therapies		
Aspirin	Starting dose of 150–300 mg orally (or 75–250 mg i.v. if oral ingestion is not possible), followed by a maintenance dose of 75–100 mg/day	
Clopidogrel	Loading dose of 300 mg orally, followed by a maintenance dose of 75 mg/day. In patients >75 years of age: loading dose of 75 mg, followed by a maintenance dose of 75 mg/day.	
Doses of anti-thrombin binding anticoagulant co-therapies		
Enoxaparin	In patients <75 years of age: 30 mg i.v. bolus followed 15 min later by 1 mg/kg s.c. every 12 h until revascularization or hospital discharge for a maximum of 8 days. The first two s.c. doses should not exceed 100 mg per injection. In patients >75 years of age: no i.v. bolus; start with first s.c. dose of 0.75 mg/kg with a maximum of 75 mg per injection for the first two s.c. doses. In patients with eGFR <30 mL/min, regardless of age, the s.c. doses are given once every 24 h.	
Unfractionated heparin	60 U/kg i.v. bolus with a maximum of 4000 U followed by an i.v. infusion of 12 U/kg with a maximum of 1000 U/h for 24–48 h. Target aPTT: 50–70 s or 1.5 to 2.0 times that of control to be monitored at 3, 6, 12, and 24 h.	
Fondaparinux (only with streptokinase)	2.5 mg i.v. bolus followed by an s.c. dose of 2.5 mg once daily for up to 8 days or until hospital discharge.	

aPTT, activated partial thromboplastin time; eGFR, estimated glomerular filtration rate; i.v., intravenous; rPA, recombinant plasminogen activator; s.c., subcutaneous; tPA, tissue plasminogen activator.

Table adapted from 2017 Guidelines on management of acute myocardial infarction in patients presenting with ST-segment elevation.

Table S11 Contraindications to fibrinolytic therapy

Absolute
Previous intracranial haemorrhage or stroke of unknown origin at any time
Ischaemic stroke in the preceding 6 months
Central nervous system damage or neoplasms, or arteriovenous malformation
Recent major trauma/surgery/head injury (within the preceding month)
Gastrointestinal bleeding within the past month
Known bleeding disorder (excluding menstrual)
Aortic dissection
Non-compressible punctures in the past 24 h (e.g. liver biopsy, lumbar puncture)
Relative
Transient ischaemic attack in the preceding 6 months
Oral anticoagulant therapy
Pregnancy or within 1-week post-partum
Refractory hypertension (systolic blood pressure >180 mmHg and/or diastolic blood pressure >110 mmHg)
Advanced liver disease
Infective endocarditis
Active peptic ulcer
Prolonged or traumatic resuscitation

Table adapted from 2017 STEMI Guidelines.

© ESC 2023

6.3.1. Comparison of fibrinolytic agents

A fibrin-specific agent is preferred.¹⁵⁴ A single bolus of weight-adjusted tenecteplase tissue plasminogen activator (tPA) is equivalent to accelerated tPA in reducing 30-day mortality, but is safer in preventing non-cerebral bleeds and blood transfusion, in addition to being easier to use in the pre-hospital setting.¹⁵⁵

6.3.2. Hazards of fibrinolysis and contraindications

Fibrinolytic therapy is associated with a small but significant excess of strokes, largely attributable to cerebral haemorrhage, with the excess hazard appearing on the first day after treatment.¹⁵⁶ Advanced age, lower weight, female sex, previous cerebrovascular disease, being from a Black population, prior history of stroke, and systolic and diastolic hypertension on admission are significant predictors of intracranial haemorrhage.¹⁵⁷ Advanced age is also associated with rupture of the LV free wall. In the latest trials, intracranial bleeding occurred in 0.9–1.0% of the total population studied.^{155,158,159} In the STREAM (Strategic Reperfusion Early After Myocardial Infarction) trial, the initial excess of intracranial haemorrhage in patients 75 years and older was reduced after a protocol amendment to reduce the dose of tenecteplase by 50%. Data from a number of studies suggest that major non-cerebral bleeds occurred in 4–13% of the patients treated.^{154,155,158,159} The most common site of spontaneous bleeding is the gastrointestinal tract. The risk of moderate to severe bleeding appears to be greater in women than in men. Administration of streptokinase may be associated with hypotension, but severe allergic reactions are rare. Re-administration of streptokinase should be avoided, both because antibodies can impair its activity and due to the risk of allergic reactions. Short, successful resuscitation does not contraindicate fibrinolytic therapy. In patients with refractory cardiac arrest, lytic therapy is not effective, increases the risk of bleeding, and is not recommended. Prolonged/traumatic but successful resuscitation increases bleeding risk and is a relative contraindication to fibrinolysis.¹⁶⁰

6.4. Antithrombotic therapy in patients not undergoing reperfusion

Among ACS patients who are medically managed without revascularization, the combination of aspirin and clopidogrel reduces the risk of CV death, MI, or stroke but increases the risk of major bleeding compared with aspirin alone.¹⁴⁵ The combination of ticagrelor and aspirin has been associated with a lower risk of CV death, MI, or stroke compared with aspirin and clopidogrel in the overall PLATO (PLATElet inhibition and patient Outcomes) trial, which was consistent in the medically managed population and across all ages.^{129,161,162} Overall bleeding risks did not differ between ticagrelor and clopidogrel in medically managed ACS patients, although observational data have suggested that ticagrelor should be used cautiously in patients aged ≥80.^{161,163} In ACS patients ≥70 years, especially if higher bleeding risk, clopidogrel may provide a favourable alternative to ticagrelor.¹⁶⁴

7. Acute coronary syndrome with unstable presentation

7.1. Out-of-hospital cardiac arrest in acute coronary syndrome

7.1.1. Healthcare systems and systems of care

Cardiac arrest centres should offer immediate multispecialty primary patient survey and stabilization with airway control and ventilatory optimization, early targeted temperature management, urgent invasive cardiology for angiography and possible haemodynamic support, streamlined delivery of critical care services, cardiac electrophysiology, 24-hour radiology including computed tomography and magnetic resonance imaging, and specialist neurology services for neurological prognostication, as well as allied health specialties and access to rehabilitation services.¹⁶⁵ Systematic reviews suggest that there is a lack of randomized clinical trial data supporting the use of cardiac arrest centres, though a large number of observational studies have reported improved outcomes with care in specialized centres.^{112,166} Some analyses of registry data have reported that provision of care in specialized centres with capability for PCI for patients with resuscitated out-of-hospital cardiac arrest is associated with improved survival to hospital discharge and lower mortality at 1 year, although the evidence in this regard is mixed.^{167–170}

8. Management of acute coronary syndrome during hospitalization

8.1. Coronary care unit/intensive cardiac care unit

Intensive cardiac care units (ICCU) provide a higher degree of care in relation to other cardiology units, up to a telemetry cardiovascular ward. ACS presentations are heterogeneous and associated with different levels of acuity and requirements for care.¹⁷¹ ACS and HF are common principal diagnoses in patients admitted to modern critical care or equivalent units, where continuous monitoring and specialized care can be provided.^{172,173}

8.2. In-hospital care

8.2.1. Duration of hospital stay

A short hospital stay may result in limited time for proper patient education and up-titration of secondary prevention treatments.

Consequently, these patients should have early post-discharge consultations with a cardiologist, primary care physician, or specialized nurse scheduled and be rapidly enrolled into a formal rehabilitation programme, either in hospital or on an outpatient basis.¹⁷⁴ Shorter lengths of stay do not appear to adversely affect adherence to discharge quality of care measures.¹⁷⁵

8.2.2. Risk assessment

8.2.2.1. Clinical risk assessment

It is important to recognize that there are several GRACE risk scores, and each refers to different patient groups and predicts different outcomes.^{127,176–178} The GRACE risk score models have been externally validated using observational data.¹⁷⁹ Online risk calculators are available for other GRACE risk scores (i.e. www.outcomes-umassmed.org/risk_models_grace_orig.aspx for the GRACE risk score 1.0 and www.outcomes-umassmed.org/grace/acs_risk2/index.html for the GRACE risk score 2.0).

Given that the GRACE risk score predicts clinical outcomes, it enables stratification of patients according to their estimated risk of future ischaemic events. A GRACE risk score-based risk assessment has been found to be superior to subjective physician assessment for the occurrence of death or MI.^{180,181} Moreover, it is well recognized that the delivery of guideline-directed care is inversely related to the estimated risk of the patient with NSTEMI-ACS—the so-called risk–treatment paradox.^{182–184} This means that patients at highest risk often receive guideline-directed care less frequently. Guideline-directed care is associated with proportionally greater survival gains among those with higher baseline risk, therefore objective risk assessment may help to identify ACS patients who would most benefit from risk-determined care interventions.^{183,184} The Australian GRACE Risk score Intervention Study (AGRIS) and the ongoing UK GRACE Risk score Intervention Study (UKGRIS) have—or are for the first time—investigating the impact of the utilization of the GRACE risk score on outcomes of patients with NSTEMI-ACS in a randomized manner.^{185,186} The AGRIS cluster-randomized trial failed to demonstrate any add-on value, especially for the guideline-directed treatments, with routine implementation of the GRACE risk score.¹⁸⁷ This was largely explained by better than expected performance of the control hospitals. Given temporal improvements in early mortality from ACS, the prediction of long-term risk is important.¹⁸⁸ Deaths in the early phase following NSTEMI-ACS are more attributable to ischaemia/thrombosis-related events, whereas in the later phase they are more likely to be associated with the progression of atherosclerosis and non-CV causes.^{189–192}

Originally, the GRACE risk score was developed to estimate the risk of in-hospital death.¹²⁷ In essence, all GRACE risk score models calculated at hospital presentation use the same eight variables (four continuous variables: age, SBP, pulse rate, and serum creatinine; three binary variables: cardiac arrest at admission, elevated cardiac biomarkers, and ST-segment deviation; and one categorical variable: Killip class at presentation) for risk prediction. The weighting of these variables, however, differs according to the model version. Continuous variables have to be entered as a range rather than exact numerical values in GRACE risk score calculators (i.e. printable charts, web calculators, and mobile phone applications). GRACE risk score calculators then use midpoints of the selected ranges for risk estimation. For the GRACE risk score 2.0, a modified score can be calculated by

substituting renal failure and use of diuretics for Killip class or serum creatinine values, respectively, if these are not available.¹⁹³ Notably, the variables used by the GRACE risk score to predict post-discharge risk are different.¹⁷⁶ The initially developed GRACE risk score for predicting the risk of in-hospital death can be calculated using a paper sheet.¹²⁷ Based on the results of a small study, utilization of an hs-cTn T assay—compared with a conventional assay—does not alter the discriminatory ability of the GRACE risk score.¹⁹⁴ Notably, the GRACE risk score model versions 1.0 and 2.0 (each derived from populations enrolled more than 10 years ago) likely overestimate risk, but discrimination into low and high risk remains good.^{181,195,196} The GRACE 3.0 score has recently been developed and is suggested to reduce sex inequalities in risk stratification.¹⁹⁷

8.2.2.2. Biomarkers for risk assessment

Quantifying the presence and severity of haemodynamic stress and HF using brain natriuretic peptide (BNP) or N-terminal pro B-type natriuretic peptide (NT-pro BNP) concentrations in patients with left main CAD or three-vessel CAD without ACS may help the Heart Team decide whether to use PCI or CABG as the revascularization strategy of choice.^{198–201} However, this needs confirmation in RCTs and has not been tested in ACS patients so far. Similarly, natriuretic peptides provide prognostic information in addition to cTn.^{202–204} Other biomarkers, such as high-sensitivity C-reactive protein, mid-regional pro-adrenomedullin, growth differentiation factor 15, heart-type fatty acid-binding protein, and copeptin may also have some prognostic value.^{101,205–210} However, the assessment of these markers has not been shown to improve patient management, and their added value in risk assessment on top of the GRACE risk calculation and/or BNP/NT-pro BNP levels appears to be marginal. Therefore, the routine use of these biomarkers for prognostic purposes is not recommended at present.

8.2.2.3. Bleeding risk assessment

In order to estimate bleeding risk, scores such as the Can Rapid risk stratification of Unstable angina patients Suppress ADverse outcomes with Early implementation of the American College of Cardiology/American Heart Association guidelines (CRUSADE [in hospital]; www.mdcalc.com/crusade-score-post-mi-bleeding-risk) and the Acute Catheterization and Urgent Intervention Triage strategY (ACUITY) (30 day) bleeding risk scores have been developed. Both of these scores have reasonable predictive value for short-term major bleeding in ACS patients undergoing coronary angiography, with CRUSADE being the most discriminatory.^{211–213} Changes in interventional practice, such as the use of radial access for coronary angiography and PCI, as well as in antithrombotic treatment, may modify the predictive value of risk scores. In addition, the predictive value of these scores has not been established in medically treated patients or in patients on OACs. Given these limitations, the use of the CRUSADE bleeding risk score may be considered in patients undergoing coronary angiography to quantify in-hospital bleeding risk.

The assessment of bleeding risk according to the Academic Research Consortium for High Bleeding Risk (ARC-HBR) criteria is a valuable alternative to these risk scores.^{214–216} This consensus definition of patients at high bleeding risk (HBR) was developed to provide consistency for clinical trials evaluating the safety and effectiveness of devices and drug regimens for patients undergoing

PCI.²¹⁷ The ARC-HBR assessment represents a pragmatic approach that includes the most recent trials performed in HBR patients, who were previously excluded from clinical trials of DAPT duration or intensity.^{218–220} However, bleeding risk assessment based on ARC-HBR criteria may be difficult to apply in routine clinical practice as several of the criteria are quite detailed (Table S12).

Table S12 Major and minor criteria for high bleeding risk according to the Academic Research Consortium for High Bleeding Risk at the time of percutaneous coronary intervention

Major criteria	Minor criteria
	Age >75 years
Anticipated use of long-term oral anticoagulation ^a	
Severe or end-stage CKD (eGFR <30 mL/min)	Moderate CKD (eGFR 30–59 mL/min)
Haemoglobin <11 g/dL	Haemoglobin 11–12.9 g/dL for men and 11–11.9 g/dL for women
Spontaneous bleeding requiring hospitalization or transfusion in the past 6 months or at any time, if recurrent	Spontaneous bleeding requiring hospitalization or transfusion within the past 12 months not meeting the major criterion
Moderate or severe baseline thrombocytopenia ^b (platelet count <100 × 10 ⁹ /L)	
Chronic bleeding diathesis	
Liver cirrhosis with portal hypertension	
	Long-term use of oral non-steroidal anti-inflammatory drugs or steroids
Active malignancy ^c (excluding non-melanoma skin cancer) within the past 12 months	
Previous spontaneous ICH (at any time)	Any ischaemic stroke at any time not meeting the major criterion
Previous traumatic ICH within the past 12 months	
Presence of a brain arteriovenous malformation	
Moderate or severe ischaemic stroke ^d within the past 6 months	
Non-deferrable major surgery on dual antiplatelet therapy	
Recent major surgery or major trauma within 30 days before percutaneous coronary intervention	

CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; ICH, intracranial haemorrhage.

Bleeding risk is high if at least one major criterion or two minor criteria are met.

^aThis excludes vascular protection doses.¹⁴³

^bBaseline thrombocytopenia is defined as thrombocytopenia before PCI.

^cActive malignancy is defined as diagnosis within 12 months and/or ongoing requirement for treatment (including surgery, chemotherapy, or radiotherapy).

^dNational Institutes of Health Stroke Scale score >5.

8.2.2.4. Integrating ischaemic and bleeding risks

The DAPT and PREdicting bleeding Complications In patients undergoing Stent implantation and subSequent Dual Anti Platelet Therapy (PRECISE-DAPT) scores have been designed to guide and inform decision-making on DAPT duration.^{221,222} The applicability of the PRECISE-DAPT score is at patient discharge, while the DAPT score is a bleeding risk estimation that is calculated at 1 year from the index event. The usefulness of the PRECISE-DAPT score was retrospectively assessed within patients randomized to different DAPT durations ($n = 10\,081$) to identify the effect on bleeding and ischaemia of a long (12–24 months) or short (3–6 months) treatment duration in relation to baseline bleeding risk.²²² Among HBR patients based on PRECISE-DAPT (i.e. PRECISE-DAPT score ≥ 25), prolonged DAPT was associated with no ischaemic benefit but an increase in the risk of bleeding events.²²² Conversely, longer treatment in patients without HBR (i.e. PRECISE-DAPT score <25) was not associated with an increase in bleeding but was associated with a significant reduction in the composite ischaemic endpoint of MI, definite stent thrombosis, stroke, and TVR. The findings remained valid in analyses restricted to ACS. However, for the majority of patients in the study, DAPT consisted of aspirin and clopidogrel. An external validation of the PRECISE-DAPT score in 4424 ACS patients undergoing PCI and treated with prasugrel or ticagrelor showed a modest predictive value for major bleeding at a median follow-up of 14 months (c-statistic = 0.65).²²³ The value of these risk prediction models in improving patient outcomes has not been established. The DAPT bleeding score has been less well validated, with a retrospective analysis in 1970 patients and a score calculation at a different time point (6 vs. 12 months) than in the derivation cohort used to generate the score.²²⁴

8.2.2.4. Evaluation of long-term risk before discharge

All patients should also have an evaluation of their long-term risk before discharge. This should include left ventricular ejection fraction (LVEF), complexity of CAD, completeness of coronary revascularization, residual ischaemia, occurrence of complications during hospitalization, and levels of metabolic risk markers (including total cholesterol, low-density lipoprotein-cholesterol [LDL-C], high-density lipoprotein-cholesterol [HDL-C], fasting triglycerides, and plasma glucose, as well as renal function). As LDL-C levels tend to decrease during the first days after MI, they should be measured as soon as possible after admission. Patients who do not undergo successful reperfusion are at higher risk of early complications and death.²²⁵ These patients should have an assessment of the presence of residual ischaemia and, if appropriate, myocardial viability.

9. Technical aspects of invasive strategies

9.1. Percutaneous coronary intervention

9.1.1. Intravascular imaging/physiology of the infarct-related artery

9.1.1.1. Intravascular imaging

The largest RCTs comparing intravascular ultrasound (IVUS)-guided PCI vs. angiography-guided PCI are the Impact of Intravascular Ultrasound Guidance on Outcomes of Xience Prime Stents in Long Lesions (IVUS-XPL) trial and the Intravascular Ultrasound Guided Drug Eluting Stents Implantation in 'All-Comers' Coronary Lesions (ULTIMATE) trial.^{226,227} The IVUS-XPL trial demonstrated, in 1400 patients (49%

with ACS) with long coronary lesions, a lower rate of MACE at 1 year for IVUS-guided than angiography-guided stent implantation (HR 0.48, 95% CI, 0.28–0.83; $P = 0.007$). The difference was attributable to a lower risk of ischaemia-driven target lesion revascularization (HR 0.51, 95% CI, 0.28–0.91; $P = 0.02$). A pre-specified subgroup analysis confirmed the MACE reduction (HR 0.35, 95% CI, 0.16–0.75) in ACS patients, with no significant interaction compared with non-ACS patients ($P = 0.20$). In 1448 all-comer patients (66% with UA and 12% with MI), the ULTIMATE trial showed significantly lower 12-month target vessel failure (TVF) in the IVUS-guided group (HR 0.530, 95% CI, 0.312–0.901). A pre-specified subgroup analysis in ACS patients showed a significant TVR reduction (HR 0.56, 95% CI, 0.32–0.99), with no significant interaction compared with non-ACS patients ($P = 0.737$). These findings were confirmed at 3 years, with a sustained lower TVF rate in the IVUS-guided group, driven mainly by the decrease in clinically driven TVR (4.5% vs. 6.9%; $P = 0.05$), especially in patients achieving IVUS-defined optimal procedural criteria.²²⁸

Two small RCTs have investigated the impact of optical coherence tomography (OCT) guidance vs. angiography guidance on surrogate endpoints. The Does Optical Coherence Tomography Optimize Results of Stenting (DOCTORS) study, randomizing 240 NSTEMI-ACS patients to OCT-guided PCI vs. angiography-guided PCI, reported a significant PCI functional result as suggested by higher post-PCI fractional flow reserve (FFR) with OCT guidance.²²⁹ Kala *et al.* randomized 201 patients with suspected STEMI to either angiography-guided PPCI or PPCI with OCT guidance. OCT guidance led to post-PCI optimization in 29% of the cases (59% for malapposition and 41% for dissections); at 9 months, OCT analysis showed significantly lower in-segment area stenosis in the OCT-guided group, with no difference in MACE.²³⁰

Two RCTs showed non-inferiority of OCT-guided PCI in comparison to IVUS-guided PCI. ILUMIEN III: OPTIMIZE PCI (OCT compared to Intravascular Ultrasound and Angiography to Guide Coronary Stent Implantation: a Multicenter Randomized Trial in PCI) demonstrated similar post-PCI minimum stent area between the OCT- and IVUS-guided PCI arms in 450 patients.²³¹ At 12 months follow-up, target lesion failure and MACE were also not significantly different between the study arms.²³² The OPTical frequency domain imaging vs. INtravascular ultrasound in percutaneous coronary intervention (OPINION) trial demonstrated similar TVF rates at 12 months with optical frequency domain imaging (OFDI)-guided PCI and angiography-guided PCI.²³³

9.1.1.2. Intravascular physiology

In ACS, the infarct-related artery (IRA) is affected to a variable extent by microvascular obstruction. Intracoronary physiology has been used in small observational studies to assess the success of myocardial reperfusion by evaluating the degree of microcirculatory resistance. In ACS patients undergoing PPCI, the index of microvascular resistance (IMR) has been correlated with the extent of microvascular obstruction (MVO) and infarct size.²³⁴ IMR has also been correlated with LV remodelling at follow-up and the extent of myocardial salvage.²³⁵ An IMR >40 (odds ratio [OR] 4.36, 95% CI, 2.10–9.06; $P < 0.001$) was an independent predictor of all-cause death or HF.²³⁶ Also, in patients with NSTEMI-ACS, post-PCI IMR in the IRA was the only independent predictor of MACE (HR 1.03, 95% CI, 1.01–1.05; $P = 0.001$).²³⁷ IMR is increasingly being used as a tool for risk stratification and/or as a surrogate endpoint to assess myocardial perfusion in phase II RCTs and has the potential to inform approaches to target microvascular dysfunction and reperfusion injury in ACS.^{238–240}

9.1.2. Embolic protection and microvascular salvage strategies

9.1.2.1. Interventions to protect the microcirculation

Recent studies in animal models reported that mechanical unloading of the left ventricle before coronary reperfusion during acute MI (AMI) may reduce infarct size and activate a cardioprotective process.^{241–243} However, these findings were not confirmed in high-risk patients (without shock) presenting with STEMI.²⁴⁴ There are no data about the potential benefits of different types of mechanical circulatory support (MCS) in high-risk AMI patients. It remains unclear if a protective strategy would provide superior clinical benefits in patients with high-risk AMI compared with a door-to-balloon strategy.²⁴⁵

Experimental studies have reported that hypothermia induced prior to reperfusion significantly reduces infarct size and is more effective if initiated soon after acute coronary occlusion.²⁴⁶ However, RCTs using different methods of systemic hypothermia, such as cold saline infusion, endovascular cooling catheters, surface cooling, and peritoneal lavage alone or in different combinations, have so far failed to show significant reductions in infarct size beyond the salvage observed with PPCI.^{247–251} Selective intracoronary hypothermia is currently being investigated in the ongoing EURO-ICE (European Intracoronary Cooling Evaluation in Patients With ST-Elevation Myocardial Infarction) RCT (NCT03447834).

In pre-clinical and small-scale clinical trials, different strategies, such as coronary post-conditioning, remote ischaemic conditioning, early intravenous (i.v.) metoprolol, glycoprotein (GP) IIb/IIIa inhibitors, drugs targeting mitochondrial integrity or nitric oxide pathways, adenosine, glucose modulators, hypothermia, and others, have been shown to be beneficial.^{252,253} However, no therapeutic intervention designed to limit reperfusion injury has translated into improved clinical outcomes. Early administration of the beta-blocker metoprolol has been shown to reduce the presence and extent of MVO in patients with anterior STEMI in the METOCARD-CNIC (Effect of METOpolol in CARDioprotection during an acute myocardial Infarction) trial.²⁵⁴ In this small trial ($n = 270$), early i.v. metoprolol was associated with smaller myocardial infarct size and improved long-term LVEF.^{255,256} The larger EARLY-BAMI (Early-Beta blocker Administration before reperfusion primary PCI in patients with ST-elevation Myocardial Infarction) trial did not confirm the infarct-limiting effect of metoprolol.²⁵⁷ Metoprolol was well tolerated in both trials and reduced the incidence of ventricular fibrillation. Differences in the timing of administration of metoprolol may explain the differences between trials.²⁵⁸ In animal models, metoprolol seems to reduce the time-dependent progression of infarction and therefore, it may make sense for it to be administered immediately after STEMI diagnosis in order to exert its cardio-protective effects.²⁵⁹ Recent data show that the protective effect of metoprolol is not shared by all beta-blockers.²⁶⁰

Deferred stenting in PPCI has been investigated as an option to prevent MVO and preserve LVEF. Clinical trials have focused either on patients with risk factors for no-reflow, such as heavy thrombus burden within the IRA, or a less selective approach involving all-comers with STEMI.²⁶¹ In DEFER-STEMI (Deferred Stent Trial in STEMI), involving patients with risk factors for no-reflow, a strategy of minimal-touch balloon angioplasty, parenteral antithrombotic therapy and deferral of stenting for 12–48 hours reduced angiographic no-reflow and intraprocedural thrombotic events.²⁶² In a less selected population (consisting of 1215 STEMI patients), the DANish Study of Optimal Acute Treatment of Patients with ST-segment Elevation Myocardial Infarction—Deferred versus conventional stent implantation in

patients with ST-segment elevation myocardial infarction (DANAMI-3–DEFER) reported that routine deferred stenting (at 48 h after the index procedure) had no effect on the primary clinical outcome (all-cause mortality, hospital admission for HF, recurrent MI, or unplanned revascularization of the IRA) or on microvascular function or infarct size at 3 months.^{263,264} These findings were consistent with other trials with a broadly similar design and a meta-analysis.^{261,265,266} Routine deferred stenting was also associated with a higher need for acute TVR due to re-occlusion of the IRA.²⁶⁴ There are no RCTs of CABG in this setting. Based on these findings, routine deferral of stenting in STEMI is not recommended. A deferred strategy may be useful in selected cases when the likelihood and clinical significance of no-reflow (e.g. heavy thrombus burden) are considered to be dominant.

9.2. Ongoing major trials

ILUMIEN IV: OPTIMAL PCI (OPTical Coherence Tomography Guided Coronary Stent Implantation Compared to Angiography: a Multicenter Randomized Trial in PCI) is a large, multicentre RCT that is currently enrolling. The sample size is approximately 3000 patients. The main objective is to assess whether OCT-guided PCI improves clinical outcomes in patients with high-risk clinical characteristics, compared with standard angiography-guided PCI.²⁶⁷

Primary Unloading and Delayed Reperfusion in ST-Elevation Myocardial Infarction: The STEMI-DTU Trial (DTU-STEMI) is currently enrolling patients (NCT03947619; DTU = Door to Unload).

Selective intracoronary hypothermia is a novel treatment designed to reduce myocardial reperfusion injury and is currently being investigated in the ongoing EURO-ICE trial (NCT03447834).

10. Management of patients with multivessel disease

There is no supplementary material for this section.

11. Myocardial infarction with non-obstructive coronary arteries

11.1. Characteristics, prognosis, and symptoms at presentation

Compared with patients with obstructive CAD, ACS patients diagnosed with MINOCA (myocardial infarction with non-obstructive coronary arteries) are more likely to be younger and female, and less likely to be diabetic, hypertensive, and dyslipidaemic. This suggests a predominant role of non-atherosclerotic-related aetiologies and a less prominent role of recognized CV risk factors.^{268–270}

Contemporary studies of MINOCA patients report a 12-month all-cause mortality of 4.7%.²⁷² Approximately 25% of patients with MINOCA will experience angina in the subsequent 12 months, which is similar to the frequency reported in patients with AMI and obstructive CAD.²⁷³

In order to correctly diagnose and manage a patient with MINOCA, patient symptoms should be investigated thoroughly, along with the patient's full history (past and present) of potentially related health issues and assessment of risk factors (Table S13).

11.2. Invasive coronary angiography

The coronary angiogram enables an immediate detailed assessment of coronary anatomy and helps to clarify a final diagnosis. Coronary angiography reveals CAD and visually enables plaque characterization (erosion, rupture), including distribution, length, calcification, dissection, and thrombosis. The angiogram also reveals coronary artery blood flow and perfusion, which can be evaluated using standard criteria such as TIMI flow grade, TIMI frame count, and TIMI myocardial perfusion grade.^{274,275} Intracoronary administration of vasoactive medications such as glyceryl trinitrate or verapamil with repeated angiography can be diagnostically useful in patients with coronary spasm, allowing direct visualization of the response to therapy. Coronary angiography is also the reference method for the diagnosis of spontaneous coronary artery dissection (SCAD).²⁷⁶ Myocardial bridges, a commonly overlooked cause of angina in patients without obstructive CAD, and which can also present as ACS, are also readily visualized with coronary angiography.²⁷⁷

11.3. Functional coronary angiography

If coronary angiography fails to clarify the aetiology in patients with a working diagnosis of MINOCA then adjunctive tests should be considered. Functional coronary angiography involves the combination of angiography with adjunctive diagnostic techniques, including intravascular imaging and physiological tests. The methodology for invasive testing of coronary vascular function has been extensively reviewed, notably by the Coronary Vasomotion Disorders International Study Group (COVADIS) and also in a recent consensus document in the context of CCS.^{278,279} Intracoronary acetylcholine or ergonovine testing may be performed when coronary or microvascular spasm is suspected.^{280,281}

11.4. Intravascular imaging (intravascular ultrasound/optical coherence tomography)

Intracoronary imaging with IVUS or OCT can also be valuable for the detection of unrecognized causes of ACS during coronary angiography, especially when thrombus, plaque rupture or erosion, or SCAD are suspected.^{282,283} In addition, intravascular imaging provides information on plaque composition, burden and outward re-modelling that is not possible to appreciate using coronary angiography and is particularly helpful to clarify ambiguous coronary lesions.²⁸⁴

11.5. Left ventricular angiography, pressure, and function

Based on the initial working diagnosis, initial assessment of LV wall motion should be promptly performed in the acute setting using LV angiography and/or echocardiography, depending on the renal function. Regional wall motion abnormalities may help confirm a final diagnosis of MI or indicate another specific underlying cause, such as takotsubo cardiomyopathy. A raised LV end-diastolic pressure (LVEDP; upper limit of normal is 12 mmHg) points to haemodynamic compromise due to impairment of systolic and/or diastolic dysfunction. Detecting a raised LVEDP may help to detect HF with preserved ejection fraction.

11.6. Non-invasive evaluation

Besides echocardiography and CMR, a computed tomography (CT) scan can be useful and can identify relevant findings in some cases (e.g. aortic dissection, pneumonia). Pulmonary embolism should also be considered as an alternative diagnosis in some cases and may be excluded with additional D-dimer/BNP testing and/or CT pulmonary angiography as appropriate.²⁸⁵

11.7. Management of myocardial infarction with non-obstructive coronary arteries

Table S13 Common manifestations of myocardial ischaemia symptoms in patients with MINOCA

Presentation of symptoms	MINOCA presenting as STEMI
Patients present more often with central chest pain	
Classically retrosternal, crushing, heavy, severe, and diffuse in nature	
Might be described by the patient as ‘pressing’ or ‘squeezing’	
May occur at rest or on activity	
May be constant or intermittent, or wax and wane in intensity	
Sometimes radiating to the left arm, neck, or jaw	
Associated symptoms	MINOCA presenting as STEMI
Nausea	
Vomiting	
Dyspnoea	
Light-headedness	
Palpitations	
Syncope	
Anxiety and or/ impending sense of doom	
Be aware of patient groups who are more likely to present with other symptoms	MINOCA presenting as STEMI
Women, older patients, and patients with diabetes are more likely to present with less common symptoms	
Less common presentations may include descriptions of chest pain as burning, throbbing, tight, or a feeling like trapped wind	
The patient may describe indigestion rather than chest pain	
In the absence of chest pain, there may be epigastric pain, back (interscapular) pain, neck or jaw pain, or arm pain (typically left-sided)	
Patients may present with breathlessness, sweating, palpitations, dizziness, nausea, or vomiting but no chest pain.	
MINOCA patients with common symptoms present more frequently with NSTEMI than STEMI	MINOCA presenting as NSTEMI
Prolonged (>20 min) chest discomfort at rest characterized by a retrosternal sensation of pain, pressure, or heaviness (‘angina’) radiating to the left arm, the right arm, both arms, the neck, or the jaw, which may be intermittent (usually lasting several minutes) or persistent	
New-onset (de novo) (<3 months) angina (Class II or III of the Canadian Cardiovascular Society classification)	
Recent destabilization of previously stable angina with at least Canadian Cardiovascular Society Class III angina characteristics (crescendo angina)	
Post-myocardial infarction angina	
Additional symptoms may be present	
Sweating	
Nausea/epigastric pain	
Dyspnoea	
Syncope	
Less common presentations of symptoms are more often observed in the older patient, in women, and in patients with diabetes, chronic renal disease, or dementia.	
Uncommon symptoms at presentation can be present	MINOCA presenting as NSTEMI
Isolated epigastric pain	
Fatigue	
Indigestion-like symptoms	
Dyspnoea	

MINOCA, myocardial infarction with non-obstructive coronary arteries; NSTEMI, non-ST elevation myocardial infarction; STEMI, ST-elevation myocardial infarction.

12. Special situations

12.1. Complications

12.1.1. Heart failure

12.1.1.1. Mechanical complications

Table S14 Mechanical complications

Aspects to consider	Ventricular septal rupture	Papillary muscle rupture	Free-wall rupture
Onset	Onset: 3–7 days from AMI	Onset: 3–7 days from AMI	Onset: 3–7 days from AMI
Clinical scenario	New acute chest pain episode, cardiogenic shock, pulmonary congestion, right heart failure signs and symptoms.	New acute chest pain episode, cardiogenic shock, acute pulmonary oedema.	New acute chest pain episode, cardiogenic shock, cardiac arrest, signs of cardiac tamponade.
Diagnosis/ workup	New cardiac murmur, echocardiographic evidence of ventricular septal defect, left-to-right ventricular shunt.	New cardiac murmur, echocardiographic evidence of mitral regurgitation with complete or partial rupture of PM (prolapse or flail of MV leaflet). Pulmonary hypertension. Hyperdynamic left ventricle.	Signs of cardiac tamponade, echocardiographic evidence of pericardial effusion, clots, contained rupture.
Management strategy	Haemodynamic stabilization (vasodilator, diuretics, IABP, inotropes). If possible, delayed surgical repair (beyond 7 days from diagnosis) with non-invasive or invasive systems like veno-arterial ECMO or other temporary percutaneous circulatory assist devices. Prompt surgery if refractory shock persistent or unresponsive right ventricular dysfunction develops.	Haemodynamic stabilization (vasodilator, diuretics, IABP, inotropes). If possible, delayed surgical repair (beyond 7 days from diagnosis) with non-invasive or invasive systems like veno-arterial ECMO or other temporary percutaneous circulatory assist devices. Prompt surgery if refractory shock persistent or unresponsive right ventricular dysfunction develops.	Immediate surgery.
Surgical treatment	Interventricular patch application. Prophylactic MCS in cases of left, right, or biventricular compromise and dysfunction. Heart transplant can be considered in patients with unsuccessful surgical repair and VSR recurrence with criteria for such a therapy and likely unsuccessful re-operation for VSR re-closure.	MV replacement (with rest of the native valve preserved besides resection of the involved PM and leaflet). In selected candidates (partial PM rupture), MV repair and PM reconstruction has been described.	Immediate surgery. Sutureless technique suggested if no large rupture present. Peri-operative support for reduced LVEDP
MCS as bridge to decision or surgery and peri-operative support	The use of MCS can improve compromised pre-operative patient conditions and/or help delay surgical repair to favour successful closure. The use of MCS, however, may have a substantial impact on underlying pathophysiology and septal shunt, potentially increasing the left-to-right shunt or even reversing the direction of blood flow, depending on the device and extent of circulatory support. MCS may also be prophylactically used to support the delicate peri-operative period and reduce the risk of patch dehiscence and VSR recurrence.	IABP to reduce MV regurgitant flow.	MCS might be useful in the pre-operative and peri-operative phases to reduce LVEDP after repair to reduce the risk of rupture recurrence.

Continued

Non-surgical treatment	Intraseptal closure device (case report evidence), particularly in the presence of prohibitive surgical risk.	Edge-to-edge MV repair in selected candidates (case report evidence) particularly in the presence of prohibitive surgical risk.	Very few cases of transventricular closure have been described. Extremely difficult procedure with risk of deterioration and sudden cardiac arrest.
Palliative care	Palliative care in the presence of very large VSR, not amenable to percutaneous closure attempt, and prohibitive risk of surgical approach.	Palliative care in the presence of prohibitive risk of surgical approach and unfeasibility of percutaneous procedure.	Palliative care (blood pressure control, aiming at chronicization and pseudoaneurysm formation) for delayed re-assessment.

AMI, acute myocardial infarction; ECMO, extracorporeal membrane oxygenation; IABP, intra-aortic balloon pump; LVEDP, left ventricular end-diastolic pressure; MCS, mechanical circulatory support; MV, mitral valve; PM, papillary muscle; VSR, ventricular septal rupture.

12.1.2. Post-acute coronary syndrome pericarditis

Early infarct-associated pericarditis (within the first 4 days after AMI, mostly transient), late pericarditis or post-cardiac injury (Dressler) syndrome (typically 1–2 weeks after AMI) and pericardial effusion have become infrequent in the era of PPCI and occur mostly with late/failed reperfusion or larger infarct size.²⁸⁶ Early pericarditis develops after AMI, with transmural necrosis causing inflammation of the adjacent pericardium. The pathophysiological mechanism in Dressler syndrome probably involves a hypersensitivity immune reaction in genetically predisposed individuals following the release of cardiac antigens during AMI.²⁸⁷

Diagnostic criteria do not differ from those for acute pericarditis and should include two of the following: (i) pleuritic chest pain (>80%); (ii) pericardial friction rub (>60%); (iii) suggestive ECG changes; and (iv) new or worsening pericardial effusion (>70%).²⁸⁸ The expected diffuse ST elevation and PR depression on ECG may be overshadowed by changes due to AMI, but persistent upright T waves and new-onset positive T waves may be seen.^{286,289,290} Inflammatory markers may rise (>80% of cases), while troponin levels may increase due to epicardial involvement. Echocardiography may show pericardial effusion while CMR can show pericardial inflammation and explore the possibility of subacute myocardial rupture in patients with significant pericardial effusion (>10 mm) post-MI.²⁸⁹

Early post-infarction pericarditis is generally self-limiting. Treatment includes aspirin 500 mg every 8–12 hours, according to the clinical case. Prolonged treatment beyond 5–7 days is usually not required.²⁸⁶ In late pericarditis, first-line therapy includes aspirin (500–1000 mg every 6–8 hours until symptomatic improvement and then taper by 250–500 mg every 2 weeks) and, as an adjunct, colchicine (0.5 mg every 12 hours) for 3 months. Although pericarditis is associated with a larger infarct size, it does not carry independent prognostic significance.²⁸⁶

The use of antithrombotics and/or anticoagulants (in the presence of LV thrombus, AF, or other indications) in patients with post-infarction pericarditis with or without pericardial effusion appears to be safe.²⁹¹

12.1.3. Bleeding

12.1.3.1. Management of bleeding

12.1.3.1.1. Bleeding events on antiplatelet agents. Treatment options in patients with ongoing bleeding while on antiplatelet therapy are limited. Platelet transfusion has been used extensively to improve platelet function in this setting, with different outcomes based on the type of antiplatelet therapy. Aspirin-inhibited platelet aggregation can be restored after transfusion of 2–5 units of platelets, whereas prasugrel- or clopidogrel-treated patients may need 4–6 h to restore platelet function after the last drug intake, and patients on ticagrelor may take ≥24 h to regain haemostatic competence.^{292–294}

12.1.3.1.2. Bleeding events on vitamin K antagonists. The risk of bleeding events for patients on VKAs increases markedly when the

international normalized ratio (INR) exceeds 4.5. Four RCTs have compared vitamin K1 with placebo in patients with an INR of 4.5–10 in the absence of ongoing bleeding, though they have only showed benefit on the surrogate outcome of reversing supratherapeutic INRs more rapidly, without evidence of benefit for hard clinical outcomes.^{295,296} Vitamin K1 administration can be used in the absence of ongoing haemorrhage in patients with an INR >10, as the risk of bleeds may be substantial. In the presence of a major or life-threatening bleed on a VKA, a combination of vitamin K1 with a rapid reversal agent (i.e. prothrombin complex concentrate, fresh frozen plasma, or recombinant activated Factor VII) should be considered.²⁹⁷

12.1.3.2. Bleeding events on non-vitamin K antagonist oral anticoagulants

After cessation of non-VKA oral anticoagulants (NOACs), improvement in haemostasis is to be expected within 12–24 h, unless patients have reduced renal function. Intracerebral haemorrhage or bleeding involving a critical organ, such as the eye, warrants immediate attempts to neutralize the anticoagulant effect of the NOAC. The first-line reversal agent to consider is the specific dabigatran antidote idarucizumab, which has been effectively tested in an uncontrolled phase III trial at a dose of 5 g i.v. in patients with uncontrollable overt bleeds or in patients requiring surgery.^{298,299} Prothrombin complex concentrates or activated prothrombin complex concentrates (i.e. with the addition of activated Factor VII) can be considered as second-line treatments when idarucizumab is not available.^{299,300} Based on studies with prothrombin complex concentrates in pre-clinical models and in healthy volunteers, an initial dose of 25 U/kg is suggested, with repeat dosing if clinically indicated. Activated prothrombin complex concentrates (50 IE/kg, with a maximum of 200 IE/kg/day) may be considered if available. Although product information for some of the NOACs mentions the use of fresh frozen plasma to help control bleeding, it seems unlikely that this would counteract drug effects.²⁹⁷ Thus, plasma should be administered only for major or life-threatening bleeds with additional dilutional coagulopathy. Neither vitamin K1 nor protamine have a role in the management of NOAC-associated bleeds.

With patients treated with Factor Xa (FXa) inhibitors (apixaban, edoxaban, rivaroxaban), prothrombin complex concentrate should be the first-line treatment.²⁹⁹ A specific antidote for FXa inhibitors, andexanet alfa, has been tested in patients with acute major bleeding associated with FXa inhibitors. At a dose of 400 mg bolus, followed by 480 mg infusion over 2 h, andexanet alfa significantly reduced anti-FXa activity, with effective haemostasis occurring in 79% of patients.^{301,302}

12.1.3.3. Bleeding events related to percutaneous coronary intervention

A pooled analysis of seven RCTs, including a total of 14 180 patients (with both stable CAD and NSTEMI-ACS), has shown that peri-procedural bleeds

are associated with a five-fold increase in 30-day mortality.³⁰³ Access site bleeding complications comprise ~40–60% of peri-procedural bleeds.^{304,305} In a pooled patient-level analysis of seven RCTs, 1-year mortality was significantly higher in patients with access site bleeds compared with those without peri-procedural bleeds.³⁰⁵ Modifications of the peri-procedural antithrombotic regimen have been efficacious in reducing peri-procedural bleeds.³⁰⁶ The radial approach for coronary angiography and PCI has been shown to be superior to the femoral one in patients with ACS. The Minimizing Adverse Haemorrhagic Events by TRansradial Access Site and Systemic Implementation of angioX (MATRIX) trial showed a significant reduction in major bleeds, as well as all-cause mortality, in patients allocated to the radial compared with the femoral approach.³⁰⁷ In the randomized Instrumental Sealing of ARterial puncture site CLOSURE device versus manual compression (ISAR-CLOSURE) trial in 4524 patients undergoing diagnostic catheterization, the incidence of vascular site complications including bleeds was 6.9% after the use of vascular closure devices and 7.9% after manual compression.³⁰⁸ Even in the context of intensified antithrombotic therapy in ACS, the use of vascular closure devices was not associated with a reduction in bleeding complications.³⁰⁴ Therefore, routine use of vascular closure devices with the goal of reducing peri-procedural bleeding complications cannot be recommended.

12.1.3.4. Bleeding events related to coronary artery bypass surgery

Bleeding events occur frequently during CABG in NSTEMI-ACS patients and are associated with the time elapsed between DAPT discontinuation and surgery.³⁰⁹ Bleeding events, as well as blood transfusions during CABG, have been associated with increased rates of morbidity and mortality.^{310,311} Severe CABG-associated bleeds in patients on DAPT should be managed with platelet concentrates. Recombinant Factor VIIa should only be used for rescue therapy in patients with uncontrollable bleeding events in whom other correctable causes have been managed (e.g. hypothermia, coagulation factor deficiencies, fibrinogen deficiency) because of concerns regarding an increased risk of graft thrombosis.³¹²

12.1.3.5. Transfusion therapy

Regardless of bleeding complications, the need for blood transfusion is associated with an approximately four-fold increase in early mortality and a three-fold increase in death or MI in ACS patients.^{313–315} The nadir haemoglobin cut-off value mandating transfusion is not standardized and varies across hospitals.^{314,316,317} In the majority of studies investigating different transfusion protocols, a liberal blood transfusion strategy has been defined as any red blood cell transfusion at a haemoglobin level <9.0 g/dL, while a restrictive blood transfusion strategy has been defined as any transfusion at a haemoglobin level <7.0 g/dL.^{316,318,319} A meta-analysis of 10 studies totalling 203 665 patients (nine observational studies and one RCT with 45 patients) with ACS (both STEMI and NSTEMI-ACS) has reported that blood transfusion or a liberal transfusion strategy was associated with increased risk of all-cause mortality (18.2 vs. 10.2%, relative risk [RR] 2.91, 95% CI, 2.46–3.44; $P < 0.001$) compared with no blood transfusion or a restrictive transfusion strategy.³¹⁸ However, a transfusion or liberal transfusion strategy seemed to be associated with a significantly higher risk of 30-day death only at a nadir haematocrit >25%.^{314,318} Observations from the CRUSADE initiative in 44 242 patients with NSTEMI-ACS reported that, among patients with haematocrit $\leq 24\%$, transfusions were associated with a trend towards a reduction in in-hospital mortality in comparison to no transfusion (11.8 vs. 15.0%, adjusted OR 0.68, 95% CI, 0.45–1.02). In patients with haematocrit between 25 and 30%, transfusions had a neutral

effect, while in those with haematocrit >30%, a significant increase in mortality was observed.³²⁰ A meta-analysis of 31, largely unblinded, RCTs totalling 9813 patients (only a minority with NSTEMI-ACS) found no significant difference in primary clinical outcomes for a liberal vs. a restrictive blood transfusion strategy.³²¹ An RCT (reported in 2015) was conducted in 2007 largely stable patients after cardiac surgery.³²² The study found no significant difference between a liberal vs. a restrictive transfusion strategy for the primary outcome of 90-day morbidity, whereas the secondary outcome of total mortality was significantly increased in the restrictive strategy arm. Based on inconsistent study results and the lack of adequately powered RCTs in the setting of NSTEMI-ACS, a restrictive policy of transfusion in anaemic patients may be considered. The effect of erythropoiesis-stimulating agents on the outcomes of ACS patients with anaemia has not been investigated. However, the accumulated evidence of these compounds in patients with congestive HF strongly suggested that they have no beneficial effects on mortality rates and may be harmful due to an increased risk of thrombo-embolism and hypertension.³¹⁶

12.2. Comorbid conditions

12.2.1. Patients at high bleeding risk and with blood disorders (anaemia and thrombocytopenia)

12.2.1.1. Thrombocytopenia following GP IIb/IIIa inhibitor therapy

Thrombocytopenia may occur during the course of therapy with GP IIb/IIIa inhibitors.³²³ Therefore, in patients treated with GP IIb/IIIa inhibitors, the platelet count should be assessed within 8–12 h of the first drug administration, at the time of any bleeding complications, and again after 24 h. GP IIb/IIIa inhibitor infusion must be discontinued if clinically relevant thrombocytopenia occurs. Platelet transfusions are recommended when there is active bleeding associated with profound thrombocytopenia, defined as a platelet count <20 000/mL.³²⁴ Platelet transfusion may be ineffective while reversibly binding GP IIb/IIIa inhibitors (eptifibatide or tirofiban) remain in circulation (half-life is 2 h for both drugs). In patients with ongoing major bleeding, fibrinogen supplementation with fresh frozen plasma or cryoprecipitate may be considered, and other supportive measures may include i.v. immunoglobulins and corticosteroids. Patients who have experienced thrombocytopenia while on GP IIb/IIIa inhibitors should avoid subsequent exposure.^{325–327}

12.2.1.2. Heparin-induced thrombocytopenia

Non-immune-mediated mild thrombocytopenia (platelet count <100 000/mL) presents within 48–72 h of the onset of unfractionated heparin (UFH) therapy, and generally resolves without complications despite continued UFH use. By contrast, immune-mediated heparin-induced thrombocytopenia (HIT) is a potentially fatal prothrombotic disorder occurring in 0.5–3% of patients who receive either UFH, a low-molecular-weight heparin, or other heparin products. HIT should be suspected when the platelet count drops to <100 000/mL (although it does not usually drop <10 000–20 000/mL). HIT usually occurs 5–10 days after a first UFH exposure, or within hours if a patient has previously received heparin. In the absence of heparin-dependent antibodies, re-exposure does not necessarily cause a relapse of HIT.^{328–330} Once HIT is suspected, all types of exposure to heparin (including flushes, coated catheters, etc.) must be discontinued. Prophylaxis against thrombosis with alternative antithrombotic therapy using non-heparin anticoagulants (e.g. argatroban, danaparoid) is indicated. Although not specifically approved for HIT, fondaparinux and bivalirudin are alternative antithrombotic agents. Bivalirudin is the recommended alternative to UFH for patients presenting with ACS who have a history of HIT. Platelet transfusions are not recommended.^{325–331}

12.2.2. Chronic kidney disease

Table S15 Recommended doses of antithrombotic agents in the acute care of patients with chronic kidney disease

Agent	Normal renal function and stage 1–3 CKD (eGFR ≥ 30 mL/min/1.73 m ²)	Stage 4 CKD (eGFR 15–29 mL/min/1.73 m ²)	Stage 5 CKD (eGFR < 15 mL/min/1.73 m ²)
Aspirin	Loading dose of 150–300 mg orally followed by a maintenance dose of 75–100 mg/day	No dose adjustment	No dose adjustment
Clopidogrel	Loading dose of 300–600 mg orally followed by 75 mg/day	No dose adjustment	No information available
Ticagrelor	Loading dose of 180 mg orally followed by 90 mg twice a day	No dose adjustment	Not recommended
Prasugrel	Loading dose of 60 mg orally followed by 10 mg/day	No dose adjustment	Not recommended
Enoxaparin	1 mg/kg s.c. twice a day 0.75 mg/kg s.c. twice daily in patients > 75 years old	1 mg/kg s.c. once a day	Not recommended
Unfractionated heparin	<i>Before coronary angiography:</i> bolus 60–70 IU/kg i.v. (maximum 5000 IU) and infusion (12–15 IU/kg/h, maximum 1000 IU/h), target activated partial thromboplastin time 1.5–2.5 $\times$ control <i>During PCI:</i> According to activated clotting time or 70–100 IU/kg i.v. in patients not anticoagulated (50–70 IU/kg if concomitant with GP IIb/IIIa inhibitors)	No dose adjustment	No dose adjustment
Fondaparinux	2.5 mg s.c. once a day	Not recommended if eGFR < 20 mL/min/1.73 m ² or patient on dialysis	Not recommended
Bivalirudin	Bolus 0.75 mg/kg i.v., infusion 1.75 mg/kg/h. <i>If eGFR ≥ 30 and ≤ 60 mL/min/1.73 m² reduce infusion dose to 1.4 mg/kg/h</i>	Not recommended	Not recommended
Eptifibatide	Bolus of 180 μ g/kg i.v. followed by an infusion of 2.0 μ g/kg/min for up to 18 h. <i>If eGFR < 50 mL/min/1.73 m² reduce infusion dose to 1.0 μg/kg/min</i>	Not recommended	Not recommended
Tirofiban	Bolus 25 μ g/kg i.v. followed by 0.15 μ g/kg/min	Reduce infusion rate to 50% if eGFR < 30 mL/min/1.73 m ²	Not recommended

© ESC 2023

Adapted from 2017 ESC Guidelines on management of acute myocardial infarction in patients presenting with ST-segment elevation.

CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; GP, glycoprotein; i.v., intravenous; PCI, percutaneous coronary intervention; s.c., subcutaneous.

12.2.3. Older adults with frailty and multimorbidity

12.2.3.1. The older person

The clinical characteristics of the older adult population are heterogeneous, with factors like frailty, comorbidity, cognitive function, and health-related quality of life playing important roles in guiding clinical care.^{332,333} Older age is also associated with a greater risk of ischaemic and bleeding events in ACS.³³⁴ The clinical presentation of ACS in the older person is more often atypical.³³⁵ Among the atypical presentations, dyspnoea is the leading symptom, while syncope, malaise, and confusion are less frequently encountered. Older patients with STEMI are more likely to experience delays between symptom onset and hospital admission, partly due to atypical presentations and delays in seeking help.³³⁶

In the context of ACS, electrocardiographic ST elevation is less frequently present in older than in younger patients.³³⁷ Hs-cTn assays have an excellent diagnostic performance for diagnosing early MI in the older person, but the specificity of the test is lower than in younger patients, and elevated troponin levels are more commonly associated with conditions other than ACS.³³⁸

12.2.3.2. Frailty and multimorbidity

Frailty and multimorbidity may impact on the degree of benefit derived from an invasive approach. Frailty assessment tools focus on either phenotype or cumulative deficit models with physicians' scaled judgment of activity, comorbidity impact, and dependency.

The number and severity of comorbidities are inversely related to rates of coronary angiography and PCI in patients with ACS. Comorbidity burden, as measured by the Charlson Comorbidity Index, predicts in-hospital and 1-year mortality in patients with ACS and is independently associated with adverse short-, medium-, and long-term outcomes after PCI.^{339,340}

12.2.3.3. Pharmacotherapy in older and frail patients

There is no reason to withhold standard medical treatment strategies in older patients. Older patients have been enrolled in the study populations of many contemporary RCTs investigating CV pharmacotherapy, in which sub-analyses showed comparable results, to varying degrees, for older and

younger patients. Therefore, in general, it is recommended that the same medical treatment strategies are applied in older ACS patients as in younger patients. Very old patients were excluded in the majority of RCTs, and therefore caution should be exercised when extrapolating trial results to this patient population. Pharmacotherapy in the acute setting and in secondary prevention should be adapted to renal function, comorbidities, comedications, frailty, and specific contraindications.

In the secondary prevention of ACS, lipid-lowering strategies play an important role. There is evidence that lipid-lowering treatment in general (and especially with statins) leads to prognostic benefits in older patients.^{341,342} Therefore, high-intensity lipid-lowering treatment is indicated for secondary prevention in older patients with ACS.

Antithrombotic treatment is mandatory in ACS patients, regardless of whether they undergo invasive management. Older patients are at particular risk of bleeding and other complications from acute as well as long-term antithrombotic therapies. This may partly be explained by the fact that renal function decreases with age and the prevalence of comorbidities is increased. Peri-procedural antithrombotic treatment is not different in older patients undergoing PCI but particular attention to proper dosing of antithrombotic therapies in this setting should be applied.³⁴³ Observational studies have reported frequent excess dosing of antithrombotic therapies in older patients.³⁴⁴

For older patients who are at HBR, different strategies with regard to antiplatelet treatment can be applied to reduce bleeding events. DAPT duration can be shortened (<12 months) or treatment intensity modified by de-escalating DAPT. Several RCTs have specifically enrolled older patients. The SENIOR (Short Duration of Dual antiplatelet Therapy With Synergy II Stent in Patients Older Than 75 Years Undergoing Percutaneous Coronary Revascularization) trial (mean age 81 years, 47% ACS) reported that 6 months of DAPT after DES implantation was safe with respect to ischaemic events in the ACS patient cohort.²²⁰ In another cohort of HBR patients (mean age 76 years, 49% ACS), a recent trial showed no negative impact on ischaemic events and a reduction in bleeding risk with 1 month vs. up to 6 months of DAPT (the P2Y₁₂ receptor inhibitor used was mostly clopidogrel) after implantation of a specific biodegradable-polymer sirolimus-eluting stent.³⁴⁵ Two ACS trials have investigated the benefit of a reduced prasugrel dose. In medically managed NSTEMI-ACS patients >75 years and in ACS patients treated with PCI (mean age 80 years), no reduction of ischaemic endpoints was achieved with prasugrel 5 mg per day vs. clopidogrel, and the bleeding risk was numerically higher in ACS-PCI patients treated with prasugrel.^{346,347} Of note, the current Summary of Product Characteristics of prasugrel recommends a dose reduction from 10 mg to 5 mg per day in patients ≥75 years. An aspirin-free concept with ticagrelor monotherapy after 3 months of DAPT with aspirin and ticagrelor demonstrated a reduced bleeding risk without a signal of increased ischaemic risk compared with continued DAPT regardless of age.^{348,349} Importantly, the same treatment effects were seen in a subgroup of patients meeting the ARC-HBR criteria, which included patients who were older and had a higher burden of comorbidities.³⁵⁰ In a sub-analysis of 2878 ACS patients aged over 75 years enrolled in the PLATO trial, the reduction of ischaemic events and overall safety of ticagrelor compared with clopidogrel was not found to be age dependent. Recent data from the SWEDEHEART (Swedish Web-System for Enhancement and Development of Evidence-Based Care in Heart Disease Evaluated According to Recommended Therapies) registry of 14 005 patients with MI aged over 80 years discharged on DAPT with aspirin and clopidogrel or ticagrelor showed that ticagrelor treatment was associated with a higher risk of bleeding and death than clopidogrel.^{162,163} In the recent POPular AGE (Ticagrelor or Prasugrel

Versus Clopidogrel in Elderly Patients With an Acute Coronary Syndrome and a High Bleeding Risk: Optimization of Antiplatelet Treatment in High-risk Elderly) trial with 1002 NSTEMI-ACS patients older than 70 years (mean age 77 years, 90% invasive management, 47% PCI, 17% CABG), DAPT with aspirin and clopidogrel vs. DAPT with aspirin and a potent P2Y₁₂ receptor inhibitor (ticagrelor 95%, prasugrel 5%) for 12 months was compared.¹⁶⁴ DAPT with clopidogrel led to fewer major and minor bleeding events and a trend towards a reduced rate of the combined net clinical benefit endpoint of all-cause death, MI, stroke, and bleeding, without an increase in MACE (CV death, MI, stroke). The discontinuation rate was twice as high with ticagrelor than with clopidogrel.

In patients with AF and ACS/PCI, DAT with a NOAC and a single antiplatelet drug reduces the risk of major and clinically relevant non-major bleeding events compared with TAT, especially with a VKA, without significantly increasing MACE.^{149–152,351} Therefore, DAT with a NOAC at the recommended dose for stroke prevention and single antiplatelet therapy (preferably with clopidogrel) for up to 12 months after a short period (up to 1 week) of TAT (with NOAC and DAPT with aspirin and clopidogrel) is recommended as the default strategy. In older patients with HBR, DAT should be shortened to 6 months by withdrawing the antiplatelet therapy and may be shortened further in older patients with very HBR, by clinical judgment; in older patients with high coronary ischaemic risk, TAT should be prolonged for up to 1 month, followed by DAT for up to 12 months. In ACS patients managed medically, available RCT data support DAT over TAT, with a single antiplatelet agent (most commonly clopidogrel) for at least 6 months.¹⁵¹

To summarize, the choice of antithrombotic agents and treatment strategy in the older ACS patient needs to be individualized, according to the patient's specific ischaemic and bleeding risk, the occurrence of adverse events, presence of comorbidities, frailty, cognitive function, and co-medications. Appropriate dosing, particularly in relation to renal function and other pharmacological factors, needs to be ensured. Proton pump inhibitors should be used in patients on DAPT, DAT, or TAT who are at increased risk of gastrointestinal bleeding. There are no specific RCTs regarding ACS-related pharmacotherapy, especially antithrombotic treatment, in frail patients.

12.2.4. Pregnancy

Cardiovascular disease in pregnancy is an increasingly important cause of maternal morbidity and mortality.^{352–355} AMI in pregnancy is associated with poor maternal and foetal outcomes, and the mortality rate is twice as high in cases where AMI occurred during the peripartum period.³⁵⁶ Overall, the incidence of AMI is higher in multigravidae and during the third trimester. In addition to traditional CV risk factors, other risk factors specific to pregnancy include pre-eclampsia, the presence of prosthetic valves, anaemia, and thrombophilia. Despite the increased risk of AMI in pregnancy, only 45% of cases are reported to undergo cardiac catheterization.^{355,357}

Pregnant women with SCAD can have a more severe clinical presentation (i.e. acute HF and multivessel dissections) in comparison to patients with non-pregnancy-associated SCAD. Additionally, SCAD has high rates of recurrence (~10% at 3-year follow-up) and MACE.^{358,359} Therefore, women of childbearing age with a history of SCAD should be carefully counselled regarding the risk of recurrent events. Pregnant women with SCAD can present with STEMI, and may also present with more serious conditions, such as cardiogenic shock. Clinical symptoms of SCAD include chest pain, dyspnoea, diaphoresis, nausea, vomiting, and a 'popping' or 'clicking' sensation in

the chest. Coronary angiography is the first-line diagnostic imaging method in SCAD due to its wide availability. When there is diagnostic uncertainty, intracoronary imaging using OCT and IVUS can be used to allow detailed visualization of the arterial wall. In the majority of cases, arteries affected by SCAD heal spontaneously, and studies have suggested that revascularization is associated with high rates of failure.³⁶⁰ In patients with ongoing or recurrent ischaemia, haemodynamic instability or isolated left main dissection, it has been suggested that PCI should be performed if the anatomy is suitable.

Coronary vasospasm in pregnancy may present as ACS as a result of enhanced vascular reactivity to angiotensin II and noradrenaline, renin release and angiotensin production due to decreased uterine perfusion in the supine position, and use of ergot derivatives to control pregnancy-related haemorrhage. The treatment is pharmacological, with vasodilators such as calcium channel blockers and nitroglycerine.³⁶¹

The highest risks of ionizing radiation exposure to the foetus occur during organogenesis and the early foetal period, while the risk decreases as pregnancy progresses from the second trimester. The recommended mean radiation exposure to the unshielded abdomen is 1.5 mGy, where <20% reaches the foetus, and procedures should follow the 'as low as reasonably achievable' radiation dose principle. Iodinated contrast material has not been reported to cause teratogenic effects. With regard to the pharmacological treatment of ACS, low-dose aspirin appears to be safe, but clopidogrel should only be used when essential and for the shortest duration of time.³⁶²

12.2.5. Drug abuse

Drug use in society is more prevalent in younger adults. Correspondingly, ACS occurring in an individual under the influence of psychotropic drugs is more common in younger individuals.³⁶³ Alcohol toxicity may occur with both acute and chronic excess intake. In the US National Inpatient Sample of patients hospitalized with AMI between 2005 and 2017, the proportion of cases associated with alcohol and illicit drug use exhibited generally increasing trends, with the exception of cocaine use.³⁶⁴

12.2.5.1. Acute coronary syndrome associated with alcohol dependence and illicit drug use

An ACS presentation can be caused by the CV effects of a toxic substance or, alternatively, reflect the natural history of ACS in an individual with atherosclerosis risk factors who at the time of the ACS is under the influence of alcohol and/or drug misuse. The complicating drug effects may manifest in the nature of the ACS presentation and also through the behaviour of the individual. This may be especially relevant in relation to non-compliance with an invasive procedure.

Alcohol has cardio-depressant effects, which can lead to hypotension. Alcohol toxicity may also cause arrhythmias, in part due to associated electrolyte disturbances. Alcohol and drug misuse can also be associated with non-cardiac, multisystem problems complicating ACS, including subarachnoid haemorrhage, pancreatitis, trauma, and infection.

12.2.5.2. Acute coronary syndrome associated with illicit drug use

Amphetamines, cocaine, and 3,4-methylenedioxymethamphetamine (MDMA) have chronotropic and vasopressor effects, while benzodiazepines, codeine, and opiates have respiratory depressant effects. Cannabis use is associated with a rising trend for CV hospitalization, in particular, for arrhythmias and, to a lesser extent, acute MI.³⁶⁵

Drugs that have vasopressor effects can be associated with Type 1 MI secondary to coronary artery plaque rupture. In addition, the use of illicit drugs with vasopressor effects may lead to atherothrombosis

through systemic hypertension, haemodynamic stress, and shear effects on coronary artery plaque. Acute myocardial injury and Type 2 MI may occur as a consequence of tachyarrhythmias, leading to a blood supply/demand mismatch. Illicit drugs with pressor effects can also induce coronary vasospasm. Sudden cardiac death is a recognized association.³⁶⁶ In a contemporary hospital registry with data from 2001 to 2015, 6.8% of ACS patients ≤50 years old presented with ACS associated with cocaine consumption.³⁶⁷ Compared with ACS patients without cocaine consumption, ACS patients with cocaine consumption exhibited a higher incidence of in-hospital ventricular tachycardia and a higher risk of recurrent MI and CV death during follow-up of up to 5 years.³⁶⁷

Patients who have presented with ACS associated with alcohol and drug misuse have an increased likelihood of re-hospitalization and an adverse prognosis in the longer term.³⁶⁷ Identifying and targeting atherosclerosis risk factors such as cigarette smoking is important. These individuals are likely to have sociodemographic factors that need to be addressed with social support and rehabilitation. Cardiac rehabilitation is particularly relevant to support compliance with secondary prevention. Clinical audit of social and cardiac rehabilitation is recommended. Systems for data collection should be in place in order to identify unwarranted variations in referral and adherence to rehabilitation programmes.

12.2.6. Patients with cancer

Cancer and its treatment are highly associated with CAD. Advances in therapies for cancer have resulted in a decline in mortality, thereby increasing life expectancy in cancer survivors. A substantial proportion of patients with active cancer or a history of cancer will present with CV disease, which is the leading cause of death in cancer survivors.³⁶⁸

Importantly, patients with active cancer have been excluded from RCTs for ACS. Cancer has additionally been omitted from all available risk stratification scores for the definition of ischaemic and bleeding risks. Moreover, most available studies have not differentiated between current vs. old cancer diagnoses, the type of cancer, and the presence/absence of metastases.

Due to the under-representation of cancer patients in prospective randomized trials, recommendations concerning patients with cancer presenting with ACS are mainly provided by expert consensus papers.^{369,370}

Table S16 Cancer treatment-related acute coronary syndrome

Accelerated atherosclerosis and plaque rupture	Androgen-deprivation therapies (GnRH agonists) Nilotinib, ponatinib, VEGF-TKIs, ICIs Radiation therapy
Vasospasm	Fluoropyrimidines, nitrogen mustards, taxanes, vinca alkaloids, bleomycin, VEGF inhibitors, ICIs
Coronary thrombosis	Alkylating agents (cisplatin, cyclophosphamide), VEGF-TKI, immunomodulatory drugs (lenalidomide, thalidomide), platinum chemotherapy, nilotinib, ponatinib, monoclonal antibodies (anti-VEGF, anti-CD20), proteasome inhibitors, ICIs

GnRH, gonadotropin-releasing hormone; ICI, immune checkpoint inhibitor; TKI, tyrosine kinase inhibitor; VEGF, vascular endothelial growth factor.

12.2.6.1. Pathophysiology

Cancer and CAD share several risk factors, with cancer patients being older and having traditional risk factors like smoking, obesity, diabetes, hypertension, hyperlipidaemia, and physical inactivity.^{368,369}

In addition to the known effects on cardiac function, cancer chemotherapy agents may affect the vascular system, including the coronary arteries. The mechanisms include coronary vasospasm, plaque rupture, or acute coronary thrombosis caused by endothelial damage (Table S16).³⁶⁹ Radiation therapy directly affects endothelial cells, leading to a variety of manifestations ranging from accelerated atherosclerosis to fibro-intimal thickening and thrombotic occlusion.^{370–372} Furthermore, cancer itself induces a pro-inflammatory and pro-thrombotic state associated with increased platelet activity and aggregability.³⁶⁹ Rarely, primary cardiac tumours may cause coronary embolism or external coronary compression, leading to ACS.

12.2.6.2. Clinical presentation

Diagnosis and definitions of ACS in cancer patients are based, to date, on the same principles as in the general population. Clinical presentation may be similar to that in patients without cancer; however, symptoms of ischaemia may be atypical or masked by cancer or therapy-related side effects, and often a high index of suspicion is required in order to secure an early diagnosis.

12.2.6.3. Initial management and acute multidisciplinary approach

Cancer patients presenting with ACS may be unstable and require multidisciplinary care in a specialized acute cardiology department with level 2 care and monitoring. A rapid diagnosis and differentiation between primary ACS and a complication secondary to cancer treatment are essential for the sometimes urgent discussion and decision about appropriate management, which may involve possible de-escalation of cancer therapy. The patient should be involved in the decision-making process as much as possible, especially with regard to changes to or interruption of cancer therapy, cardiac interventions, and strategic decisions about future cancer therapy.

12.2.6.4. Invasive strategy

Evidence-based management strategies are limited as cancer patients have been excluded from randomized trials and prospective studies. Observational data have reported differences in survival between patients with active and previous cancer diagnoses as well as in patients with and without metastases. Moreover, no prospective RCT has evaluated the risks and benefits of conservative vs. invasive strategies in patients with cancer who present with ACS. Registry data show that patients with current cancer and ACS have at least a 50% increased risk of major adverse cardiovascular and cerebrovascular events (MACCE), bleeding complications, and in-hospital mortality in comparison to patients without cancer. On the other hand, patients with a history of cancer seem to bear no increased risk of adverse outcomes, with the exception of bleeding. Furthermore, current cancer is more often associated with a more conservative management strategy than those used in patients with no cancer or history of cancer.^{368,373}

Lung cancer is associated with the highest mortality and MACCE, while colon cancer is associated with the highest bleeding risk, with

breast and prostate cancer patients having fewer complications. In patients with metastatic cancer and ACS, PCI is not associated with a mortality benefit in comparison to conservative treatment.³⁷⁴

Patients with high-risk ACS should always be evaluated for invasive management since PCI is associated with lower mortality in such patients. On the contrary, a conservative approach may be appropriate in non-high-risk ACS patients who have a poor cancer prognosis and/or HBR.³⁶⁹

12.2.6.5. Antithrombotic treatment

Patients with a history of cancer should be treated in the same way as all other ACS patients. For patients with active cancer, treatment should be individualized according to the type and treatment of cancer, the type of ACS management, the need for further cancer treatment after ACS, the level of pro-coagulant state, anaemia, and thrombocytopenia.³⁶⁹

According to the ARC-HBR criteria, patients with an active cancer diagnosed in the past 12 months are regarded as having HBR. In such patients the combination of aspirin (300 mg loading dose [LD]/75–100 mg maintenance dose [MD]) and clopidogrel (300–600 mg LD /75 mg MD) can be recommended if the platelet count is >10 000/μL for aspirin and >30 000/μL for the combination. Ticagrelor and prasugrel should not be used because of the HBR and the paucity of data in patients with active cancer, but they could be discussed in patients with previous stent thrombosis while taking aspirin and clopidogrel, with strict surveillance of bleeding risk.³⁶⁹

12.2.6.6. Thrombocytopenia and cancer

About 10–25% of cancer patients have thrombocytopenia (platelet count <100 000/μL). Although bleeding is more frequent in ACS patients with thrombocytopenia, aspirin use is not associated with a higher bleeding risk and is associated with a survival advantage.^{375,376} It is therefore recommended by experts that aspirin should be given to all patients with a platelet count >10 000/μL.^{369,370} Thrombocytopenia should not exclude patients from an invasive approach, which is associated with improved outcome after ACS in cancer patients.^{377,378}

12.2.6.7. Cancer treatment

Interdisciplinary discussion is necessary for patients with a direct relation of cancer treatment to the generation of ACS, and in these patients an alternative cancer treatment should be considered. In cases of coronary vasospasm due to cancer therapy that cannot be changed, a re-challenge with the same treatment under close monitoring is negotiable when underlying CAD has been excluded or treated. The concomitant use of a calcium antagonist in combination with a long-acting nitrate and aspirin can be prescribed, although it is not unequivocally effective.³⁶⁹

12.2.7. Coronavirus disease (COVID-19)

Cardiovascular comorbidities are common in patients with coronavirus disease 2019 (COVID-19), and the presence of CVD is associated with severe COVID-19 and higher mortality. Cardiac manifestations are associated with worse outcomes of COVID-19.

Evidence of acute cardiac injury with raised troponin levels appears in COVID-19 patients several days after initiation of fever, suggesting myocardial damage associated with viral infection. Mechanisms of

SARS-CoV-2-induced myocardial injury remain elusive, but it may be secondary to a direct effect of the virus on the myocardium, hypoxia, and/or myocardial inflammation in the context of a systemic immune response.^{379,380}

Chest pain and breathlessness are frequent symptoms in patients diagnosed with COVID-19. The same ECG diagnostic criteria for cardiac conditions apply in patients affected by SARS-CoV-2 infection and in the general population.

In patients hospitalized with COVID-19, mild elevations in cTn are in general the result of pre-existing cardiac disease and/or the acute injury/stress related to COVID-19.

SARS-CoV-2 infection is associated with an increased thrombotic burden.^{381,382} However, COVID-19 does not change the management (including organized networks) of patients with ACS.³⁸³ A PPCI strategy remains the treatment of choice for STEMI patients.³⁸⁴ Fibrinolysis is indicated (always within the first 12 h of MI) only in those in whom PCI-mediated reperfusion is not anticipated to occur within 120 min of STEMI diagnosis.

Reports have highlighted the unfavourable prognosis of ACS patients with concomitant COVID-19 disease.^{385,386} Therefore, appropriate vaccination against COVID-19 is recommended for ACS patients.

13. Long-term treatment

13.1. Lifestyle management

13.1.1. Nutrition and alcohol

Table S17 Characteristics of a healthy diet

Adopt a more plant-based and less animal-based food pattern
Increase consumption of fruit to ≥ 200 g per day
Increase consumption of vegetables to ≥ 200 g per day
35–45 g of fibre per day, preferably from whole grains
30 g unsalted nuts daily
1–2 servings of fish per week (one to be oily fish)
Limited lean meat, low-fat dairy products, and liquid vegetable oils
Red meat should be reduced to a maximum of 300–500 g a week; processed meat should be minimized
Saturated fats to account for $<10\%$ of total energy intake; replace with polyunsaturated fats
Trans-unsaturated fats as low as possible; preferably no intake from processed food, and $<1\%$ of total energy intake
≤ 5 g of salt per day
If alcohol is consumed, limit intake to two glasses (20 g) daily for men and one glass for women (10 g), or a total of 100 g per week
Avoid energy-dense foods such as sugar and sweetened soft drinks ¹¹

© ESC 2023

13.1.2. Resumption of activities

Resuming prior activities, including returning to work, engaging in personal and social activities, driving and travelling are important components of recovery after ACS. Returning to work is an important indicator of recovery. A recent review of observational data showed that male, younger, educated, non-manual workers or those who owned their own business, patients who evaluated their general and mental health highly, and those with shorter hospitalization, and fewer comorbidities and complications were more likely to return to work

after an MI.³⁸⁷ Offering adequate psychosocial and vocational support as required is an important part of comprehensive cardiac rehabilitation.^{11,388}

In 2013, the European Union issued standards on driving and CVD. However, laws can vary between countries and local regulations should always be consulted before advising patients. An aircraft environment generally does not pose a significant threat to CV health and is safe in revascularized post-ACS patients with stable symptoms and no HF.³⁸⁹ Patients who have suffered ACS with complications, especially with accompanying low LVEF ($<40\%$) or who have undergone CABG, should postpone air travel until their condition is more stable.³⁸⁹ Further, patients should always be advised to consult their insurance company and airline as rules and regulations for air travel for patients with chronic conditions may vary considerably.

13.1.2.1. Sexual activity

Counselling and advice from healthcare providers about sexual activity should be offered. Resumption of sexual activity is important to patients, and they often worry if it is going to harm them. Information on when they can resume activity according to their physical ability should be given. Information should be provided regarding a low risk of sudden death or AMI when sexual activity is with a stable partner in a familiar environment, and that they should avoid large meals and alcohol beforehand. Reassure the patient that if they can climb two flights of stairs this is the equivalent to the amount of expended energy needed during sexual activities. People who are physically active have less of a risk of adverse events during sexual activity.² In patients with a recent intake of a phosphodiesterase 5 inhibitor (within 24 h for sildenafil or vardenafil and 48 h for tadalafil), nitrates should not be administered.

13.1.2.2. Environmental factors

Air pollutants have been estimated to be one of the 10 leading risk factors for global mortality.³⁹⁰ Exposure to air pollution increases the risk of MI as well as hospitalization and death from HF, stroke, and arrhythmia.^{391,392} Patients with CCS should avoid strenuous outdoor exercise in heavily polluted areas (for instance in heavily traffic-congested areas).³⁹³ Air purifiers with high-efficiency particulate air filters reduce indoor pollution, and wearing N95 respirator face masks in heavily polluted areas has been shown to be protective.^{394,395} Studies have also shown that environmental noise increases the risk of CVD.³⁹⁶ Policies and regulations that reduce air pollution and environmental noise should be supported, and patients should be advised regarding these risks.

13.2. Pharmacological treatment

13.2.1. Lipid-lowering therapy

Given the extensive RCT evidence of prognostic benefits, statins are the first choice for pharmacological LDL-C lowering.³⁹⁷ The higher the patient's CV risk and the greater the achieved LDL-C reduction, the greater the absolute risk reduction by statin-mediated LDL-C lowering.³⁹⁸ In the ACS setting, the routine early use of high-intensity statin therapy is associated with rapid and sustained clinical benefits.³⁹⁹ It is therefore recommended that high-intensity statin therapy (e.g. with atorvastatin or rosuvastatin) is initiated as early as possible, during the first 1–4 days of hospitalization for the index ACS, and prescribed up to the highest tolerated dose in order to reach the LDL-C goals.⁴⁰⁰ The intensity of statin therapy should be increased in patients receiving low- or moderate-intensity statin treatment at presentation. If the




Figure S5 Information for patients on how to optimize their 'heart health' after an acute coronary syndrome. BMI, body mass index; LDL, low-density lipoprotein.

LDL-C goals are not achieved with the maximum tolerated dose of a statin after 4–6 weeks following the ACS event, combination with ezetimibe is recommended.⁴⁰⁰ Efficacy in terms of CV event reduction and the safety of ezetimibe treatment have been demonstrated in post-ACS patients.⁴⁰¹

Proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitors (the monoclonal antibodies evolocumab and alirocumab) are very effective in reducing LDL-C, regardless of baseline LDL-C and baseline lipid-lowering therapy, with ~50–60% lowering of LDL-C. Of note, PCSK9 inhibitor treatment also reduces lipoprotein(a) levels by approximately 25%.⁴⁰² In large outcome trials, both evolocumab and alirocumab led to a significant reduction in CV events in patients with stable atherosclerotic cardiovascular disease (ASCVD) including CAD, and in post-ACS patients, respectively, with a good safety profile and no apparent negative effects on liver, muscles, glucose metabolism, kidneys, and cognitive function, but also no impact on mortality during trial follow-up.^{403–406} Patients with a higher absolute CV risk, such as those with recent or multiple ACS events, or concomitant peripheral arterial or polyvascular disease, experience greater absolute risk reductions with PCSK9 inhibitor treatment ('highest risk, highest benefit').^{407–409} Sub-analyses showed that patients who achieved the lowest LDL-C values with PCSK9 inhibitor treatment also had the lowest risk of future MACE.⁴¹⁰ Very low levels of LDL-C (below 40 mg/dL) were generally well tolerated in the outcome trials using ezetimibe and PCSK9 inhibitors, and LDL-C lowering to these levels was associated

with even lower CV event rates.^{410–413} Data from the FOURIER (Further Cardiovascular Outcomes Research with PCSK9 Inhibition in Subjects with Elevated Risk) trial with evolocumab showed a consistent linear relationship between the achieved LDL-C level and major CV outcomes down to LDL-C concentrations of <0.2 mmol/L, without safety concerns around these very low LDL-C levels.⁴¹⁰ For secondary prevention, in patients with ACS not achieving their LDL-C goal on a maximum tolerated dose of a statin and ezetimibe, combination therapy with a PCSK9 inhibitor is recommended.⁴⁰⁰ The optimal timing of PCSK9 inhibitor treatment initiation remains to be determined. *Post-hoc* analyses of the outcome trials indicate that early initiation may be beneficial. A recent trial investigating PCSK9 inhibitor treatment initiation in the acute phase of ACS showed that evolocumab added to high-intensity statin therapy was well tolerated and resulted in early substantial reduction of LDL-C levels, with >95% of patients achieving the currently recommended LDL-C goals within 4–8 weeks.⁴¹⁴ The initiation of PCSK9 inhibitor treatment is recommended in patients with ACS who do not reach their LDL-C goal after 4–6 weeks of maximum tolerated statin and ezetimibe therapy. In patients who present with ACS and whose LDL-C levels are not at goal despite already taking a maximally tolerated statin dose and ezetimibe prior to the event, the addition of a PCSK9 inhibitor early after the event (during hospitalization for the ACS event if possible) should be considered. In statin-intolerant patients, first ezetimibe and secondly PCSK9 inhibitors should be used to achieve LDL-C goals.

The recent REDUCE-IT (Reduction of Cardiovascular Events With Icosapent Ethyl–Intervention Trial) trial with 8179 patients (70% with established CV disease) on statin therapy and with elevated triglycerides demonstrated a beneficial effect of icosapent ethyl, a highly purified and stable eicosapentaenoic acid ethyl ester, given at a high dose of 2 g twice a day, on a composite of CV death, MI, stroke, coronary revascularization, or UA compared with placebo.⁴¹⁵ MACE (CV death, MI, or stroke) and CV death were significantly reduced as well. Therefore, icosapent ethyl, at a dose of 2 g b.i.d., may be considered in combination with a statin in patients with ACS and triglyceride levels of 1.5–5.6 mmol/L (135–499 mg/dL) despite statin treatment.⁴⁰⁰

Recently, new compounds have become available for the treatment of hypercholesterolaemia. Bempedoic acid is a novel, first-in-class, oral small molecule that inhibits cholesterol synthesis by inhibiting the action of ATP citrate lyase, an enzyme upstream of HMG-CoA reductase. In a phase III trial in 2230 ASCVD patients on maximally tolerated statin therapy and a mean LDL-C level of 103 mg/dL, treatment with bempedoic acid 180 mg per day lowered LDL-C by ~18% vs. placebo.⁴¹⁶ Interestingly, high-sensitivity C-reactive protein was also significantly lowered, and there was no effect on muscle-related symptoms. Bempedoic acid was also tested in statin-intolerant patients and in combination with ezetimibe.⁴¹⁷ An alternative approach targeting PCSK9 uses RNA interference. The small interfering RNA molecule inclisiran is an injectable compound with long-lasting effects on PCSK9 synthesis and is administered subcutaneously every 6 months. Inclisiran was tested in two phase III trials in patients with ASCVD on maximally tolerated statin therapy, with or without other LDL-C-lowering agents.⁴¹⁸ A meta-analysis combining these two RCTs with a trial in patients with heterozygous familial hypercholesterolaemia, including a total of 3660 patients, demonstrated that inclisiran treatment reduced LDL-C by ~50% over a time course of 18 months, with stable effects on LDL-C for ≥6 months after each injection. A high percentage of patients achieved LDL-C reductions ≥50% and an LDL-C threshold <50 mg/dL.⁴¹⁹ No specific serious adverse events were observed. There are no outcome data available yet; a large CV outcomes trial is currently comparing inclisiran against placebo in patients with prior MI or stroke. New therapeutic options with alternative targets, especially those using RNA-based technologies (among others targeting lipoprotein(a), ANGPTL3, and ApoCIII), are currently being investigated and are at various stages of clinical development.⁴²⁰

13.2.2. Hormone replacement therapy

The risk of ACS increases in post-menopausal women. With observational study data implying a potential cardio-protective benefit of hormone replacement therapy (HRT) with oestrogen and progestin, several large trials were initiated in the 1990s to further test the hypothesis.^{421,422} However, the results showed that HRT actually increased CVD risk. Therefore, HRT should not be prescribed for cardio-protective purposes in post-menopausal women. Whether ongoing HRT in women

presenting with ACS should be discontinued is less well established.^{423,424} Individual evaluation, taking the indication for HRT, the patient's symptoms, patient's preferences, and overall CVD risk into account, should precede any decision on HRT discontinuation post-ACS.

14. Patient perspectives

Increased patient participation in their healthcare improves patient self-efficacy, health outcomes, and quality of life.⁴²⁵ Collaborative management using the best available evidence leads to better treatment compliance, better self-care, improved health literacy, better medication adherence, decreased anxiety and depression, and fewer hospitalizations.⁴²⁶

14.1. Patient-centred care

Using a shared decision-making approach allows patients' needs, preferences, and decisions to be established.

Explicitly asking the question 'What matters most to you?', and especially 'Why does this matter most?', will provide a more holistic picture and assist with the design of a personalized ACS care plan. This provides crucial information, which is necessary to deliver patient-centred care.

Patient-centred care improves patients' ability to handle information and have a better understanding of shared documentation. This approach particularly benefits those patients with lower levels of education.^{427,428} Actively involving family and friends (according to the patient's wishes) in the care process means their contribution to patient recovery is augmented through person-centred care.⁴²⁷ These models allow patients to feel included and heard, which improves the quality of care received and positively affects patient satisfaction. This ultimately improves patient health outcomes and enhances their quality of life by delivering care more appropriate to their own health needs.

14.2. Informed consent

The 'teach back' method (Figure S6) confirms that the patient has understood the clinical information provided. The simple question 'Do you understand?' does not confirm actual understanding. Similarly, the closed question 'Do you understand what I have told you?' does not allow the clinician to assess if the patient/family member has understood what has been said to them. To assess understanding of informed consent an open-ended question such as 'Can you tell me what risks from the procedure make you more concerned?' is preferable.

Informed consent is an ethical and legal obligation for medical practitioners and is required before any invasive procedure. The patient must be competent and have the capacity to make a voluntary decision. Informed consent should preferably be part of a process and should include the components listed in Table S18.^{429,430}

The 'teach back' technique



Figure S6 Informed consent process using the 'teach back' technique.

Providing written information before the procedure, in an easy-to-understand format, prepares the patient and family for a structured conversation.

Patients with ACS may have difficulty adhering to healthy lifestyles and medication regimens, which may be caused by false or poor comprehension of the informed consent process. Informed consent is not just a signature—it also relates to patient education and patient understanding in relation to what is being done and why.⁴³¹ Information given to patients undergoing PCI is heterogeneous, varies in the amount and quality, risks are forgotten, benefits overestimated, and alternative treatments not always considered.⁴³⁰ It is essential to make sure the patient understands what has been discussed during the consenting process.

It has been reported that patients do not always remember or understand the information given to them.^{430,432} Using decision aides and educational interventions can improve recall of the decision-making process by improving patient knowledge. The use of decision aids when discussing risk can also improve risk perception.⁴³⁰

Information should be provided in a simple, clear, and unbiased format. Between 27% and 48% of patients in Europe have inadequate health literacy, which impacts on the ability of patients to manage their own care.⁴³³ Health literacy refers not only to the patient's knowledge of health information, but also their ability to understand, access, appraise, and apply information to make informed decisions after judging all available material.⁴³⁴

Table S18 Components needed for informed consent

Components of discussion	Needed information
The nature of the procedure	• What the procedure entails
	• Aim of procedure
	• Condition warranting the procedure
	• Procedure itself
	• X-ray exposure and the C-arm
	• Access site
	• Personnel in room
	• Medication given during procedure
	• Position on table
	• Possibility of discomfort or pain from the procedure
Risks and benefits of the procedure	• Complications that can occur
	• Outcome to expect
Reasonable alternatives	• Coronary artery bypass grafting vs. stenting
	• Medical therapy only
	• No treatment
Risk and benefits of alternatives	• What could happen if they decide to have no treatment
	• Complications that can occur from alternative treatment
	• Expected outcomes of the alternative treatment
Assessment of the patient's understanding of information given	• Use 'teach back' technique to evaluate patient's understanding

© ESC 2023

14.3. Research participation and consent in the acute setting

The main aspect of informed consent for a clinical trial is to allow the patient or their surrogate to decide to participate or decline enrolment, and to inform the patient of the key aspects of the study.⁴³² The Declaration of Helsinki permits waiving consent for clinical trials for acute conditions when patients are unconscious and surrogates cannot provide consent within an appropriate time-frame.⁴³⁵ When the patient is unconscious a surrogate can act on the patient's behalf for making the decisions for participation in trials. It has been reported that when patients were asked about which approach to informed consent was most appropriate, a majority were happy that they were asked to be in a study, whereas some would prefer that the cardiologist makes the decision and asks for consent later.⁴³²

Patients who are conscious and not mentally impaired should be involved in enrolment decisions. It has been reported that when witnessed verbal consent was used for STEMI patients, the majority of patients had recall of being included in the study and a positive opinion about being asked to participate. The majority of patients also stated

that they did not want more comprehensive information during the initial informed consent discussion.⁴³⁶ It is important to have contact with the patient after the intervention, as many patients and surrogates may have poor understanding or forget the information that was given to them, even when assessed shortly after consenting.⁴³² Patients view *post-hoc* consent negatively, therefore medical staff should help facilitate the patient's expectations and preferences, as most patients want to be involved in the enrolment process.^{432,436,437} It has been reported that decisional regret was low and patients felt they were given enough information for deciding to participate in studies (written consent).⁴³² To avoid the risk of including a patient unwillingly in a trial, communication between the physician and patient/surrogate, allowing the patient/surrogate the chance to decline participation, is crucial. Using a short verbal informed consent process, followed by complementary written information after the acute phase, is seen as adequate from the patient's point of view. Improvement in the written format of the consent form, making it short, simple, and clear in language, can improve the process for both research purposes and normal care.

The consent process for ACS patients and patients participating in trials can differ according to presentation and the need for acute intervention. The standard process may need to be shortened due to time constraints. It is recommended to use a shortened verbal informed consent process in patients undergoing emergency invasive angiography, and when including patients in trials in the emergency setting, written consent should be considered after the acute phase.

14.4. Patient-reported outcome measures and patient-reported experience measures

Patient-reported outcome measures (PROMs) are standardized, validated tools used to measure the patient's reported status of their health condition, coming directly from the patient without interpretation. PROMs can be used for generic purposes or for a specific condition, and when chosen should be a reliable and validated tool for the condition under evaluation. They measure the patient's perception of their quality of life, status of disease, and general health. PROMs are used in the pursuit of improving care and can help determine the cost-effectiveness of clinical interventions.⁴³⁸

Patient-reported experience measures (PREMs) report the patient's perception of care given to them by assessing the following domains that impact patients' perception of safe, effective, patient-centred, timely, efficient, and equitable care.^{439,440}

PROMs and PREMs can be used to assess the quality of care for ACS patients during their patient journey. The quality of care of ACS patients should be measured during the patient's journey from initial presentation until discharge.⁴⁴¹ PREMs are usually gathered in a survey format and differ from satisfaction surveys in that they do not allow subjective views or comments.⁴⁴⁰ Training of staff in the correct use of PROMs and PREMs, and how to apply and use the information correctly, is needed for proper evaluation.

Patient perception and expectations of care are built on interpersonal interactions, quality of clinical interactions, delivery of care, and the administrative management of care. Each ACS patient has individual ideas and perceptions about how they should be treated and what constitutes the best possible care. Therefore, patient-centred care is important as it recognizes and incorporates the values and wishes of the patient when providing the healthcare needed.

Table S19 Patient expectations and clear communication for patients with acute coronary syndrome

Components of clear communication for patients with ACS	
Clear communication is required in...	✓ Interpersonal interactions ✓ Clinical quality interactions ✓ Care delivery interactions ✓ Administrative interactions
Patient expectations	How patient expectations can be met
To be able to recognize my symptoms of ACS	✓ Awareness/understanding of risk factors: both 'traditional' risk factors and other, e.g. female-specific and ethnicity risk factors ✓ Know how/where/when to seek appropriate help
Right care at the right time	✓ All symptoms to be taken seriously ✓ Help in articulation of symptoms ✓ Appropriate questions asked and a detailed/full history taken ✓ Timely care to all-comers, irrespective of age, sex, ethnicity, body habitus, or social background
High-quality, effective and safe care delivered by professionals	✓ Healthcare staff to be highly skilled/trained ✓ Healthcare staff to have excellent interpersonal/communication skills ✓ Protocols and guidelines are adhered to ✓ Care without judgment/bias ✓ Information flow—keep patient updated
Clear, comprehensible information and health literacy taken into account	✓ Awareness of levels of patient health literacy ✓ Explanations/information given in simple terms ✓ Use every encounter as an opportunity for patient education ✓ Check for understanding using, e.g. teach back technique ✓ Check for understanding of medication regimen and side effects of medications ✓ Check for understanding of treatment/care plan ✓ Check for understanding of long-term lifestyle and risk management

Continued

	✓ Check for understanding of potential intervention plan in the event of re-occurrence of new/related symptoms
Consider not only physical, but mental and emotional well-being	✓ Emotional support, empathy, and respect ✓ Adequate psychological and emotional insights into the patient's priorities ✓ Awareness of cognitive impact in relation to presentation/circumstance ✓ Reassurance
Shared decision-making and respect for preferences	✓ Ask patients: 'What matters to you?' 'Why?' ✓ Personalized care plan that includes all dimensions ✓ Precision medicine with person-centred care
Consideration/involvement of, and support for, family and carers	✓ Take a biopsychosocial perspective
Readiness for discharge	✓ Support for self-care ✓ Realistic conversations ✓ Information on disease, treatment plan, medication, pain control, secondary prevention ✓ Information provided orally and in written form ✓ Check for understanding ✓ Copies of patient discharge letter and other relevant documents
Continuity of care	✓ Within hospital ✓ Transition to primary care/onward care/referral ✓ Cardiac rehabilitation

© ESC 2023

14.5. Preparation for discharge

There is a need for a different approach to communicating with ACS patients at the time of discharge. Patient-centred care that allows personal narratives to emerge may enable healthcare professionals to offer an individualized care plan, with guidance for ACS patients that will help them cope with the everyday challenges they experience after discharge.⁴⁴²

Upon hospital discharge the patient should receive a copy of their discharge letter/document, including therapeutic targets and information on follow-up care.^{443–446} In addition, patient information leaflets are also helpful, but should use plain language, avoid medical terms and abbreviations, highlight important information, and use pictures and a large font size.^{430,443–447}

All aspects of self-care should be reviewed with the patient before discharge to help the patient's understanding. This may help improve

Table S20 Preparation for discharge

	Patient discharge plan Instructions/information	Primary care provider Discharge summary
	Discharge information (verbal and written) should include:	Discharge letter to primary care provider should include:
1	- Educate the patient: lifestyle changes (tailored to patient profile) - Diet - Smoking cessation - Body weight - Exercise	Risk factors and history Cardiovascular risk factors and history
2	Educate the patient: - Reason for hospitalization - Diagnosis - Medications - Procedures and test results - Warning symptoms, and what to do and who to call if problems arise - Knowing risks with delaying treatment	Reason for admission (symptoms, electrocardiogram changes, troponin) - Main diagnosis (Dx) (ST-elevation myocardial infarction [STEMI], non-ST-elevation myocardial infarction [NSTEMI], unstable angina [UA]) - Additional Dx (heart failure, diabetes mellitus, arrhythmia)
3		Invasive Angiography extent of disease, culprit lesion, was complete revascularization performed or is further revascularization planned?
4		Left ventricular ejection fraction at discharge (%)
5	Results of laboratory values and need to follow up pending results	Main lab values - Estimated glomerular filtration rate - Peak troponin - Glycosylated haemoglobin (HbA1c) - Low-density lipoprotein-cholesterol (LDL-C) (admission)
6	Medications: - Confirm correct medications at discharge (emphasizing any changes from admission) - Inform and educate patient regarding the purpose of the medications, how to take them correctly,	Discharge treatment: Aspirin(dose) P2Y ₁₂ inhibitor (reason for choice) Beta-blocker Angiotensin-converting enzyme (ACE) inhibitor Lipid-lowering drug Proton pump inhibitor Specific: Hypertension

Continued

	potential adverse side effects, and doses - Check for understanding of the medicine regimen - Provide an up-to-date written medication list - Instruct patient how to obtain medications	- Additional lipid lowering - Heart failure - Anticoagulant - Anti-arrhythmic treatment - Implanted cardioverter defibrillator or life vest
7	Need for cardiac rehab and follow-up	Therapeutic targets (for patient profile) - LDL-C (reason for adding ezetimibe) - HbA1c reason for increase/change in treatment - Duration of dual antiplatelet therapy: - Type and duration - Mention of high bleeding or high ischaemic risk - Strategy if using chronic anticoagulation Other risk factors of lifestyle (tailored to patient profile) - Diet - Smoking cessation - Body weight - Exercise
8	Contact information of primary care provider Appointments for follow-up care Outpatient services and medical equipment	Structured follow-up: - Time of first cardiology visit - Time of first cardiac rehabilitation visit - Time of other tests

© ESC 2023

medication adherence and decrease the risk of complications, including re-admission.^{430,448}

The patient's priorities about what they want to know are not always correctly perceived by the healthcare professional, and the priority of the patient's needs with regard to information change from the time of admission through to, and after, discharge.

When teaching patients, it is important to appreciate that the level of health literacy for each individual is different: many patients have low health literacy, so it may be useful to provide information in portions and check for understanding after each piece is provided.⁴⁴⁷

There are many new medications and recommendations for changes in lifestyle that a patient receives upon discharge—using the teach back method or motivational interviewing is recommended as an approach to educating the patient and/or surrogate.

Patient concerns and educational needs throughout their ACS journey are summarized in [Figure S7](#).




Figure S7 Patient concerns and educational needs throughout their ACS journey. ACS, acute coronary syndrome; CCU, cardiac care unit; OPD, out-patient department.

15. Gaps in evidence

Some of the gaps in evidence reported are addressed by the following ongoing trials.

DROP-Asian is an international stepped-wedge cluster randomized clinical trial investigating the implementation of the 0 h/1 h ESC algorithm in patients with suspected NSTEMI-ACS in Asia (Umin Id: UMIN000042461).

PRECISE1MI is a multicentre, prospective stepped-wedge cluster randomized clinical trial aiming to evaluate the clinical implementation of the ESC 0 h/1 h algorithm and its safety and efficacy for the diagnosis of AMI compared with the current algorithm, in nine different countries worldwide (NCT05649384).

CODE-MI (hs-cTn-Optimizing the Diagnosis of Acute Myocardial Infarction/Injury in Women) is a multicentre, stepped-wedge, cluster randomized trial that aims to assess the impact of using the women-specific 99th percentile cut-off for high hs-cTn, compared with uniform 99th percentile cut-offs, on the diagnosis, treatment, and outcomes of women presenting to the ED with suspected ACS.⁴⁴⁹

Results from the ongoing **SENIOR-RITA** (The older patients Randomised Interventional Trial in Acute non-ST elevation myocardial infarction) (NCT03052036) RCT comparing conservative and invasive management strategies in the context of NSTEMI are awaited.

16. References

- Thygesen K. 'Ten Commandments' for the Fourth Universal Definition of Myocardial Infarction 2018. *Eur Heart J* 2019;**40**:226. <https://doi.org/10.1093/eurheartj/ehy856>
- Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al. 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. The Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2021;**42**:1289–1367. <https://doi.org/10.1093/eurheartj/ehaa575>
- Roe MT, Harrington RA, Prosper DM, Pieper KS, Bhatt DL, Lincoff AM, et al. Clinical and therapeutic profile of patients presenting with acute coronary syndromes who do not have significant coronary artery disease. *Circulation* 2000;**102**:1101–1106. <https://doi.org/10.1161/01.CIR.102.10.1101>
- Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, et al. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2016;**37**:267–315. <https://doi.org/10.1093/eurheartj/ehv320>
- Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, et al. Fourth universal definition of myocardial infarction (2018). *Eur Heart J* 2019;**40**:237–269. <https://doi.org/10.1093/eurheartj/ehy462>
- Chapman AR, Shah ASV, Lee KK, Anand A, Francis O, Adamson P, et al. Long-term outcomes in patients with type 2 myocardial infarction and myocardial injury. *Circulation* 2018;**137**:1236–1245. <https://doi.org/10.1161/CIRCULATIONAHA.117.031806>
- Neumann JT, Sorensen NA, Rubsamen N, Ojeda F, Renné T, Qaderi V, et al. Discrimination of patients with type 2 myocardial infarction. *Eur Heart J* 2017;**38**:3514–3520. <https://doi.org/10.1093/eurheartj/ehx457>
- Nestelberger T, Boeddinghaus J, Badertscher P, Twerenbold R, Wildi K, Breitenbücher D, et al. Effect of definition on incidence and prognosis of type 2 myocardial infarction. *J Am Coll Cardiol* 2017;**70**:1558–1568. <https://doi.org/10.1016/j.jacc.2017.07.774>
- Kuznetsova T. Sex differences in epidemiology of cardiac and vascular disease. *Adv Exp Med Biol* 2018;**1065**:61–70. https://doi.org/10.1007/978-3-319-77932-4_4
- Virani SS, Alonso A, Aparicio HJ, Benjamin EJ, Bittencourt MS, Callaway CW, et al. Heart disease and stroke statistics-2021 update: a report from the American Heart Association. *Circulation* 2021;**143**:e254–e743. <https://doi.org/10.1161/cir.00000000000000950>
- Visseren FLJ, Mach F, Smulders YM, Carballo D, Koskinas KC, Bäck M, et al. 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J* 2021;**42**:3227–3337. <https://doi.org/10.1093/eurheartj/ehab484>
- Sugiyama T, Hasegawa K, Kobayashi Y, Takahashi O, Fukui T, Tsugawa Y. Differential time trends of outcomes and costs of care for acute myocardial infarction hospitalizations by ST elevation and type of intervention in the United States, 2001–2011. *J Am Heart Assoc* 2015;**4**:e001445. <https://doi.org/10.1161/jaha.114.001445>
- Puymirat E, Simon T, Cayla G, Cottin Y, Elbaz M, Coste P, et al. Acute myocardial infarction: changes in patient characteristics, management, and 6-month outcomes over a period of 20 years in the FAST-MI program (French registry of acute ST-elevation or non-ST-elevation myocardial infarction) 1995 to 2015. *Circulation* 2017;**136**:1908–1919. <https://doi.org/10.1161/circulationaha.117.030798>
- Townsend N, Wilson L, Bhatnagar P, Wickramasinghe K, Rayner M, Nichols M. Cardiovascular disease in Europe: epidemiological update 2016. *Eur Heart J* 2016;**37**:3232–3245. <https://doi.org/10.1093/eurheartj/ehw334>
- Timmis A, Vardas P, Townsend N, Torbica A, Katus H, De Smedt D, et al. European Society of Cardiology: cardiovascular disease statistics 2021. *Eur Heart J* 2022;**43**:716–799. <https://doi.org/10.1093/eurheartj/ehab892>
- García-García C, Oliveras T, Serra J, Vila J, Rueda F, Cedié G, et al. Trends in short- and long-term ST-segment-elevation myocardial infarction prognosis over 3 decades: a Mediterranean population-based ST-segment-elevation myocardial infarction registry. *J Am Heart Assoc* 2020;**9**:e017159. <https://doi.org/10.1161/jaha.120.017159>
- Gale CP, Allan V, Cattle BA, Hall AS, West RM, Timmis A, et al. Trends in hospital treatments, including revascularisation, following acute myocardial infarction, 2003–2010: a multilevel and relative survival analysis for the National Institute for Cardiovascular Outcomes Research (NICOR). *Heart* 2014;**100**:582–589. <https://doi.org/10.1136/heartjnl-2013-304517>
- Puymirat E, Simon T, Steg PG, Schiele F, Guérat P, Blanchard D, et al. Association of changes in clinical characteristics and management with improvement in survival among patients with ST-elevation myocardial infarction. *JAMA* 2012;**308**:998–1006. <https://doi.org/10.1001/2012.jama.11348>
- Regitz-Zagrosek V, Oertelt-Prigione S, Prescott E, Franconi F, Gerds E, Forst-Ludwig A, et al. Gender in cardiovascular diseases: impact on clinical manifestations, management, and outcomes. *Eur Heart J* 2016;**37**:24–34. <https://doi.org/10.1093/eurheartj/ehv598>
- Hvelplund A, Galatius S, Madsen M, Rasmussen JN, Rasmussen S, Madsen JK, et al. Women with acute coronary syndrome are less invasively examined and subsequently less treated than men. *Eur Heart J* 2010;**31**:684–690. <https://doi.org/10.1093/eurheartj/ehp493>
- Rashid M, Curzen N, Kinnaird T, Lawson CA, Myint PK, Kontopantelis E, et al. Baseline risk, timing of invasive strategy and guideline compliance in NSTEMI: nationwide analysis from MINAP. *Int J Cardiol* 2020;**301**:7–13. <https://doi.org/10.1016/j.ijcard.2019.11.146>
- van Oosterhout REM, de Boer AR, Maas A, Rutten FH, Bots ML, Peters SAE, et al. Sex differences in symptom presentation in acute coronary syndromes: a systematic review and meta-analysis. *J Am Heart Assoc* 2020;**9**:e014733. <https://doi.org/10.1161/jaha.119.014733>
- Araújo C, Laszczyńska O, Viana M, Melão F, Henriques A, Borges A, et al. Sex differences in presenting symptoms of acute coronary syndrome: the EPIHeart cohort study. *BMJ Open* 2018;**8**:e018798. <https://doi.org/10.1136/bmjopen-2017-018798>
- Hadid LAA, Al Barmawi M, Al Hmaitat NAA, Shohrat N. Factors associated with pre-hospital delay among men and women newly experiencing acute coronary syndrome: a qualitative inquiry. *Cardiol Res Pract* 2020;**2020**:3916361. <https://doi.org/10.1155/2020/3916361>
- Heer T, Hochadel M, Schmidt K, Mehili J, Zahn R, Kuck K-H, et al. Sex differences in percutaneous coronary intervention—insights from the coronary angiography and PCI registry of the German Society of Cardiology. *J Am Heart Assoc* 2017;**6**:e004972. <https://doi.org/10.1161/jaha.116.004972>
- Gimenez MR, Reiter M, Twerenbold R, Reichlin T, Wildi K, Haaf P, et al. Sex-specific chest pain characteristics in the early diagnosis of acute myocardial infarction. *JAMA Intern Med* 2014;**174**:241–249. <https://doi.org/10.1001/jamainternmed.2013.12199>
- Henrikson CA, Howell EE, Bush DE, Miles JS, Meiningner GR, Friedlander T, et al. Chest pain relief by nitroglycerin does not predict active coronary artery disease. *Ann Intern Med* 2003;**139**:979–986. <https://doi.org/10.7326/0003-4819-139-12-200312160-00007>
- Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, et al. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: the Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2018;**39**:119–177. <https://doi.org/10.1093/eurheartj/ehx393>
- Wouters L, Zwart DLM, Erkelens DCA, De Groot E, van Smeden M, Hoes AW, et al. Gender-stratified analyses of symptoms associated with acute coronary syndrome in telephone triage: a cross-sectional study. *BMJ Open* 2021;**11**:e042406. <https://doi.org/10.1136/bmjopen-2020-042406>
- Boeddinghaus J, Nestelberger T, Twerenbold R, Neumann JT, Lindahl B, Giannitsis E, et al. Impact of age on the performance of the ESC 0/1h-algorithms for early diagnosis of myocardial infarction. *Eur Heart J* 2018;**39**:3780–3794. <https://doi.org/10.1093/eurheartj/ehy514>
- Twerenbold R, Badertscher P, Boeddinghaus J, Nestelberger T, Wildi K, Puelacher C, et al. 0/1-hour triage algorithm for myocardial infarction in patients with renal dysfunction. *Circulation* 2018;**137**:436–451. <https://doi.org/10.1161/CIRCULATIONAHA.117.028901>
- Chiaromonte GR, Friend R. Medical students' and residents' gender bias in the diagnosis, treatment, and interpretation of coronary heart disease symptoms. *Health Psychol* 2006;**25**:255–266. <https://doi.org/10.1037/0278-6133.25.3.255>
- Anderson HVS, Masri SC, Abdallah MS, Chang AM, Cohen MG, Elgendy IY, et al. 2022 ACC/AHA key data elements and definitions for chest pain and acute myocardial infarction: a report of the American Heart Association/American College of Cardiology Joint Committee on clinical data standards. *J Am Coll Cardiol* 2022;**80**:1660–1700. <https://doi.org/10.1016/j.jacc.2022.05.012>
- Persson A, Hartford M, Herlitz J, Karlsson T, Omland T, Caidahl K. Long-term prognostic value of mitral regurgitation in acute coronary syndromes. *Heart* 2010;**96**:1803–1808. <https://doi.org/10.1136/hrt.2010.203059>
- Smith SW, Dodd KW, Henry TD, Dvorak DM, Pearce LA. Diagnosis of ST-elevation myocardial infarction in the presence of left bundle branch block with the ST-elevation to S-wave ratio in a modified Sgarbossa rule. *Ann Emerg Med* 2012;**60**:766–776. <https://doi.org/10.1016/j.annemergmed.2012.07.119>
- Nestelberger T, Cullen L, Lindahl B, Reichlin T, Greenslade JH, Giannitsis E, et al. Diagnosis of acute myocardial infarction in the presence of left bundle branch block. *Heart* 2019;**105**:1559–1567. <https://doi.org/10.1136/heartjnl-2018-314673>
- Meyer MR, Radovanovic D, Pedrazzini G, Rickli H, Roffi M, Rosemann T, et al. Differences in presentation and clinical outcomes between left or right bundle branch block and ST segment elevation in patients with acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:848–856. <https://doi.org/10.1177/2048872620905101>
- Lai YC, Chen YH, Wu KH, Chen YC. Validation of the diagnosis and triage algorithm for acute myocardial infarction in the setting of left bundle branch block. *Am J Emerg Med* 2020;**38**:2614–2619. <https://doi.org/10.1016/j.ajem.2020.03.024>
- Cai Q, Mehta N, Sgarbossa EB, Pinski SL, Wagner GS, Califf RM, et al. The left bundle-branch block puzzle in the 2013 ST-elevation myocardial infarction guideline: from falsely declaring emergency to denying reperfusion in a high-risk population. Are the Sgarbossa criteria ready for prime time? *Am Heart J* 2013;**166**:409–413. <https://doi.org/10.1016/j.ahj.2013.03.032>

40. Nestelberger T, Boeddinghaus J, Lopez Ayala P, Gehrke J, Twerenbold R, Cullen L, et al. Utility of echocardiography in patients with suspected acute myocardial infarction and left bundle-branch block. *J Am Heart Assoc* 2021;**10**:e021262. <https://doi.org/10.1161/JAHA.121.021262>
41. Pera VK, Larson DM, Sharkey SW, Garberich RF, Solie CJ, Wang YL, et al. New or presumed new left bundle branch block in patients with suspected ST-elevation myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2018;**7**:208–217. <https://doi.org/10.1177/2048872617691508>
42. Neumann JT, Sorensen NA, Rubsamen N, Ojeda F, Schäfer S, Keller T, et al. Right bundle branch block in patients with suspected myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2019;**8**:161–166. <https://doi.org/10.1177/2048872618809700>
43. Holmvang L, Clemmensen P, Lindahl B, Lagerqvist B, Venge P, Wagner G, et al. Quantitative analysis of the admission electrocardiogram identifies patients with unstable coronary artery disease who benefit the most from early invasive treatment. *J Am Coll Cardiol* 2003;**41**:905–915. [https://doi.org/10.1016/S0735-1097\(02\)02970-4](https://doi.org/10.1016/S0735-1097(02)02970-4)
44. Kaul P, Fu Y, Chang WC, Harrington RA, Wagner GS, Goodman SG, et al. Prognostic value of ST segment depression in acute coronary syndromes: insights from PARAGON-A applied to GUSTO-IIb. *J Am Coll Cardiol* 2001;**38**:64–71. [https://doi.org/10.1016/S0735-1097\(01\)01307-9](https://doi.org/10.1016/S0735-1097(01)01307-9)
45. Savonitto S, Cohen MG, Politi A, Hudson MP, Kong DF, Huang Y, et al. Extent of ST-segment depression and cardiac events in non-ST-segment elevation acute coronary syndromes. *Eur Heart J* 2005;**26**:2106–2113. <https://doi.org/10.1093/eurheartj/ehi395>
46. Damman P, Holmvang L, Tijssen JG, Lagerqvist B, Clayton TC, Pocock SJ, et al. Usefulness of the admission electrocardiogram to predict long-term outcomes after non-ST-elevation acute coronary syndrome (from the FRISC II, ICTUS, and RITA-3 [FIR] trials). *Am J Cardiol* 2012;**109**:6–12. <https://doi.org/10.1016/j.amjcard.2011.08.002>
47. Jacobsen MD, Wagner GS, Holmvang L, Kontny F, Wallentin L, Husted S, et al. Quantitative T-wave analysis predicts 1 year prognosis and benefit from early invasive treatment in the FRISC II study population. *Eur Heart J* 2005;**26**:112–118. <https://doi.org/10.1093/eurheartj/ehi026>
48. Mueller C, Neumann FJ, Perach WV, Perruchoud AP, Buettner HJ. Prognostic value of the admission electrocardiogram in patients with unstable angina/non-ST-segment elevation myocardial infarction treated with very early revascularization. *Am J Med* 2004;**117**:145–150. <https://doi.org/10.1016/j.amjmed.2004.02.034>
49. Boersma E, Pieper KS, Steyerberg EW, Wilcox RG, Chang W-C, Lee KL, et al. Predictors of outcome in patients with acute coronary syndromes without persistent ST-segment elevation. Results from an international trial of 9461 patients. *Circulation* 2000;**101**:2557–2567. <https://doi.org/10.1161/01.CIR.101.22.2557>
50. Tan NS, Goodman SG, Yan RT, Elbarouni B, Budaj A, Fox KAA, et al. Comparative prognostic value of T-wave inversion and ST-segment depression on the admission electrocardiogram in non-ST-segment elevation acute coronary syndromes. *Am Heart J* 2013;**166**:290–297. <https://doi.org/10.1016/j.ahj.2013.04.010>
51. Gorgels AP, Engelen DJ, Wellens HJ. Lead aVR, a mostly ignored but very valuable lead in clinical electrocardiography. *J Am Coll Cardiol* 2001;**38**:1355–1356. [https://doi.org/10.1016/S0735-1097\(01\)01564-9](https://doi.org/10.1016/S0735-1097(01)01564-9)
52. Yamaji H, Iwasaki K, Kusachi S, Murakami T, Hirami R, Hamamoto H, et al. Prediction of acute left main coronary artery obstruction by 12-lead electrocardiography: ST segment elevation in lead aVR with less ST segment elevation in lead V(1). *J Am Coll Cardiol* 2001;**38**:1348–1354. [https://doi.org/10.1016/S0735-1097\(01\)01563-7](https://doi.org/10.1016/S0735-1097(01)01563-7)
53. Yan AT, Yan RT, Kennelly BM, Anderson FA, Budaj A, López-Sendón J, et al. Relationship of ST elevation in lead aVR with angiographic findings and outcome in non-ST elevation acute coronary syndromes. *Am Heart J* 2007;**154**:71–78. <https://doi.org/10.1016/j.ahj.2007.03.037>
54. Khan AR, Golwala H, Tripathi A, Bin Abdulhak AA, Bavishi C, Riaz H, et al. Impact of total occlusion of culprit artery in acute non-ST elevation myocardial infarction: a systematic review and meta-analysis. *Eur Heart J* 2017;**38**:3082–3089. <https://doi.org/10.1093/eurheartj/ehx418>
55. Wang TY, Zhang M, Fu Y, Armstrong PW, Newby LK, Gibson CM, et al. Incidence, distribution, and prognostic impact of occluded culprit arteries among patients with non-ST-elevation acute coronary syndromes undergoing diagnostic angiography. *Am Heart J* 2009;**157**:716–723. <https://doi.org/10.1016/j.ahj.2009.01.004>
56. de Zwaan C, Bär FV, Wellens HJ. Characteristic electrocardiographic pattern indicating a critical stenosis high in left anterior descending coronary artery in patients admitted because of impending myocardial infarction. *Am Heart J* 1982;**103**:730–736. [https://doi.org/10.1016/0002-8703\(82\)90480-x](https://doi.org/10.1016/0002-8703(82)90480-x)
57. de Zwaan C, Bär FV, Janssen JH, Cheriex EC, Dassen WRM, Brugada P, et al. Angiographic and clinical characteristics of patients with unstable angina showing an ECG pattern indicating critical narrowing of the proximal LAD coronary artery. *Am Heart J* 1989;**117**:657–665. [https://doi.org/10.1016/0002-8703\(89\)90742-4](https://doi.org/10.1016/0002-8703(89)90742-4)
58. de Winter RJ, Verouden NJ, Wellens HJ, Wilde AA. A new ECG sign of proximal LAD occlusion. *N Engl J Med* 2008;**359**:2071–2073. <https://doi.org/10.1056/NEJMc0804737>
59. Abächerli R, Twerenbold R, Boeddinghaus J, Nestelberger T, Mächler P, Sassi R, et al. Diagnostic and prognostic values of the V-index, a novel ECG marker quantifying spatial heterogeneity of ventricular repolarization, in patients with symptoms suggestive of non-ST-elevation myocardial infarction. *Int J Cardiol* 2017;**236**:23–29. <https://doi.org/10.1016/j.ijcard.2017.01.151>
60. Strebel I, Twerenbold R, Boeddinghaus J, Abächerli R, Rubini Giménez M, Wildi K, et al. Diagnostic value of the cardiac electrical biomarker, a novel ECG marker indicating myocardial injury, in patients with symptoms suggestive of non-ST-elevation myocardial infarction. *Ann Noninvasive Electrocardiol* 2018;**23**:e12538. <https://doi.org/10.1111/anec.12538>
61. Strebel I, Twerenbold R, Wussler D, Boeddinghaus J, Nestelberger T, du Fay de Lavallaz J, et al. Incremental diagnostic and prognostic value of the QRS-T angle, a 12-lead ECG marker quantifying heterogeneity of depolarization and repolarization, in patients with suspected non-ST-elevation myocardial infarction. *Int J Cardiol* 2019;**277**:8–15. <https://doi.org/10.1016/j.ijcard.2018.09.040>
62. Grimm K, Twerenbold R, Abächerli R, Boeddinghaus J, Nestelberger T, Koehlin L, et al. Diagnostic and prognostic value of ST-segment deviation scores in suspected acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:857–868. <https://doi.org/10.1177/2048872619853579>
63. Eggers KM, Jernberg T, Lindahl B. Cardiac troponin elevation in patients without a specific diagnosis. *J Am Coll Cardiol* 2019;**73**:1–9. <https://doi.org/10.1016/j.jacc.2018.09.082>
64. Nestelberger T, Lopez-Ayala P, Boeddinghaus J, Strebel I, Rubini Gimenez M, Huber I, et al. External validation and extension of a clinical score for the discrimination of type 2 myocardial infarction. *J Clin Med* 2021;**10**:1264. <https://doi.org/10.3390/jcm10061264>
65. Nestelberger T, Boeddinghaus J, Lopez-Ayala P, Kaier TE, Marber M, Gysin V, et al. Cardiovascular biomarkers in the early discrimination of type 2 myocardial infarction. *JAMA Cardiol* 2021;**6**:771–780. <https://doi.org/10.1001/jamacardio.2021.0669>
66. Horiuchi Y, Wettersten N, Patel MP, Mueller C, Neath S-X, Christenson RH, et al. Biomarkers enhance discrimination and prognosis of type 2 myocardial infarction. *Circulation* 2020;**142**:1532–1544. <https://doi.org/10.1161/CIRCULATIONAHA.120.046682>
67. Boeddinghaus J, Twerenbold R, Nestelberger T, Koehlin L, Wussler D, Meier M, et al. Clinical use of a new high-sensitivity cardiac troponin I assay in patients with suspected myocardial infarction. *Clin Chem* 2019;**65**:1426–1436. <https://doi.org/10.1373/clinchem.2019.304725>
68. Boeddinghaus J, Twerenbold R, Nestelberger T, Badertscher P, Wildi K, Puelacher C, et al. Clinical validation of a novel high-sensitivity cardiac troponin I assay for early diagnosis of acute myocardial infarction. *Clin Chem* 2018;**64**:1347–1360. <https://doi.org/10.1373/clinchem.2018.286906>
69. Boeddinghaus J, Nestelberger T, Twerenbold R, Koehlin L, Meier M, Troester V, et al. High-sensitivity cardiac troponin I assay for early diagnosis of acute myocardial infarction. *Clin Chem* 2019;**65**:893–904. <https://doi.org/10.1373/clinchem.2018.300061>
70. Chapman AR, Lee KK, McAllister DA, Cullen L, Greenslade JH, Parsonage W, et al. Association of high-sensitivity cardiac troponin I concentration with cardiac outcomes in patients with suspected acute coronary syndrome. *JAMA* 2017;**318**:1913–1924. <https://doi.org/10.1001/jama.2017.17488>
71. Twerenbold R, Neumann JT, Sorensen NA, Ojeda F, Karakas M, Boeddinghaus J, et al. Prospective validation of the 0/1-h algorithm for early diagnosis of myocardial infarction. *J Am Coll Cardiol* 2018;**72**:620–632. <https://doi.org/10.1016/j.jacc.2018.05.040>
72. Neumann JT, Sorensen NA, Schwemer T, Ojeda F, Bourry R, Sciacca V, et al. Diagnosis of myocardial infarction using a high-sensitivity troponin I 1-hour algorithm. *JAMA Cardiol* 2016;**1**:397–404. <https://doi.org/10.1001/jamacardio.2016.0695>
73. Reichlin T, Irfan A, Twerenbold R, Reiter M, Hochholzer W, Burkhalter H, et al. Utility of absolute and relative changes in cardiac troponin concentrations in the early diagnosis of acute myocardial infarction. *Circulation* 2011;**124**:136–145. <https://doi.org/10.1161/CIRCULATIONAHA.111.023937>
74. Reichlin T, Schindler C, Drexler B, Twerenbold R, Reiter M, Zellweger C, et al. One-hour rule-out and rule-in of acute myocardial infarction using high-sensitivity cardiac troponin T. *Arch Intern Med* 2012;**172**:1211–1218. <https://doi.org/10.1001/archinternmed.2012.3698>
75. Boeddinghaus J, Reichlin T, Cullen L, Greenslade JH, Parsonage WA, Hammett C, et al. Two-hour algorithm for triage toward rule-out and rule-in of acute myocardial infarction by use of high-sensitivity cardiac troponin I. *Clin Chem* 2016;**62**:494–504. <https://doi.org/10.1373/clinchem.2015.249508>
76. Neumann JT, Sorensen NA, Rubsamen N, Ojeda F, Schock A, Seddighzadeh P, et al. Evaluation of a new ultra-sensitivity troponin I assay in patients with suspected myocardial infarction. *Int J Cardiol* 2019;**283**:35–40. <https://doi.org/10.1016/j.ijcard.2018.12.001>
77. Greenslade J, Cho E, Van Hise C, Hawkins T, Parsonage W, Ungerer J, et al. Evaluating rapid rule-out of acute myocardial infarction using a high-sensitivity cardiac troponin I assay at presentation. *Clin Chem* 2018;**64**:820–829. <https://doi.org/10.1373/clinchem.2017.283887>
78. Pickering JW, Than MP, Cullen L, Aldous S, ter Avest E, Body R, et al. Rapid rule-out of acute myocardial infarction with a single high-sensitivity cardiac troponin T measurement below the limit of detection: a collaborative meta-analysis. *Ann Intern Med* 2017;**166**:715–724. <https://doi.org/10.7326/M16-2562>
79. Shah AS, Anand A, Sandoval Y, Lee KK, Smith SW, Adamson PD, et al. High-sensitivity cardiac troponin I at presentation in patients with suspected acute coronary

- syndrome: a cohort study. *Lancet* 2015;**386**:2481–2488. [https://doi.org/10.1016/S0140-6736\(15\)00391-8](https://doi.org/10.1016/S0140-6736(15)00391-8)
80. Wildi K, Cullen L, Twerenbold R, Greenslade JH, Parsonage WV, Boeddinghaus J, et al. Direct comparison of 2 rule-out strategies for acute myocardial infarction: 2-h accelerated diagnostic protocol vs 2-h algorithm. *Clin Chem* 2017;**63**:1227–1236. <https://doi.org/10.1373/clinchem.2016.268359>
 81. Nestelberger T, Boeddinghaus J, Greenslade J, Parsonage WA, Than M, Wussler D, et al. Two-hour algorithm for rapid triage of suspected acute myocardial infarction using a high-sensitivity cardiac troponin I assay. *Clin Chem* 2019;**65**:1437–1447. <https://doi.org/10.1373/clinchem.2019.305193>
 82. Koechlin L, Boeddinghaus J, Nestelberger T, Lopez-Ayala P, Wussler D, Shrestha S, et al. Performance of the ESC 0/2h-algorithm using high-sensitivity cardiac troponin I in the early diagnosis of myocardial infarction. *Am Heart J* 2021;**242**:132–137. <https://doi.org/10.1016/j.ahj.2021.08.008>
 83. Boeddinghaus J, Lopez-Ayala P, Nestelberger T, Koechlin L, Münch T, Miro O, et al. Prospective validation of the ESC 0/1h-algorithm using high-sensitivity cardiac troponin I. *Am J Cardiol* 2021;**158**:152–153. <https://doi.org/10.1016/j.amjcard.2021.08.007>
 84. Boeddinghaus J, Nestelberger T, Lopez-Ayala P, Koechlin L, Buechi M, Miro O, et al. Diagnostic performance of the European Society of Cardiology 0/1-h algorithms in late presenters. *J Am Coll Cardiol* 2021;**77**:1264–1267. <https://doi.org/10.1016/j.jacc.2021.01.004>
 85. Sörensen NA, Neumann JT, Ojeda F, Giannitsis E, Spanuth E, Blankenberg S, et al. Diagnostic evaluation of a high-sensitivity troponin I point-of-care assay. *Clin Chem* 2019;**65**:1592–1601. <https://doi.org/10.1373/clinchem.2019.307405>
 86. Reichlin T, Cullen L, Parsonage WA, Greenslade J, Twerenbold R, Moehring B, et al. Two-hour algorithm for triage toward rule-out and rule-in of acute myocardial infarction using high-sensitivity cardiac troponin T. *Am J Med* 2015;**128**:369–379.e4. <https://doi.org/10.1016/j.amjmed.2014.10.032>
 87. Boeddinghaus J, Nestelberger T, Koechlin L, Wussler D, Lopez-Ayala P, Walter JE, et al. Early diagnosis of myocardial infarction with point-of-care high-sensitivity cardiac troponin I. *J Am Coll Cardiol* 2020;**75**:1111–1124. <https://doi.org/10.1016/j.jacc.2019.12.065>
 88. Koechlin L, Boeddinghaus J, Lopez-Ayala P, Nestelberger T, Wussler D, Mais F, et al. Diagnostic discrimination of a novel high-sensitivity cardiac troponin I assay and derivation/validation of an assay-specific 0/1h-algorithm. *Am Heart J* 2023;**255**:58–70. <https://doi.org/10.1016/j.ahj.2022.10.007>
 89. Mueller C. Biomarkers and acute coronary syndromes: an update. *Eur Heart J* 2014;**35**:552–556. <https://doi.org/10.1093/eurheartj/ehf530>
 90. Reichlin T, Twerenbold R, Maushart C, Reiter M, Moehring B, Schaub N, et al. Risk stratification in patients with unstable angina using absolute serial changes of 3 high-sensitivity troponin assays. *Am Heart J* 2013;**165**:371–378.e3. <https://doi.org/10.1016/j.ahj.2012.11.010>
 91. Reichlin T, Twerenbold R, Reiter M, Steuer S, Bassetti S, Balmelli C, et al. Introduction of high-sensitivity troponin assays: impact on myocardial infarction incidence and prognosis. *Am J Med* 2012;**125**:1205–1213.e1. <https://doi.org/10.1016/j.amjmed.2012.07.015>
 92. Shah ASV, Anand A, Strachan FE, Ferry AV, Lee KK, Chapman AR, et al. High-sensitivity troponin in the evaluation of patients with suspected acute coronary syndrome: a stepped-wedge, cluster-randomised controlled trial. *Lancet* 2018;**392**:919–928. [https://doi.org/10.1016/S0140-6736\(18\)31923-8](https://doi.org/10.1016/S0140-6736(18)31923-8)
 93. Mueller C, Giannitsis E, Mockel M, Huber K, Mair J, Plebani M, et al. Rapid rule out of acute myocardial infarction: novel biomarker-based strategies. *Eur Heart J Acute Cardiovasc Care* 2017;**6**:218–222. <https://doi.org/10.1177/2048872616653229>
 94. Mockel M, Giannitsis E, Mueller C, Huber K, Jaffe AS, Mair J, et al. Rule-in of acute myocardial infarction: focus on troponin. *Eur Heart J Acute Cardiovasc Care* 2017;**6**:212–217. <https://doi.org/10.1177/2048872616653228>
 95. Katus H, Ziegler A, Ekinci O, Giannitsis E, Stough WG, Achenbach S, et al. Early diagnosis of acute coronary syndrome. *Eur Heart J* 2017;**38**:3049–3055. <https://doi.org/10.1093/eurheartj/ehx492>
 96. Neumann JT, Twerenbold R, Ojeda F, Sörensen NA, Chapman AR, Shah ASV, et al. Application of high-sensitivity troponin in suspected myocardial infarction. *N Engl J Med* 2019;**380**:2529–2540. <https://doi.org/10.1056/NEJMoa1803377>
 97. Lopez-Ayala P, Nestelberger T, Boeddinghaus J, Koechlin L, Ratmann PD, Strebel I, et al. Novel criteria for the observe-zone of the ESC 0/1h-hs-cTnT algorithm. *Circulation* 2021;**144**:773–787. <https://doi.org/10.1161/CIRCULATIONAHA.120.052982>
 98. Ambavane A, Lindahl B, Giannitsis E, Roiz J, Mendivil J, Frankenstein L, et al. Economic evaluation of the one-hour rule-out and rule-in algorithm for acute myocardial infarction using the high-sensitivity cardiac troponin T assay in the emergency department. *PLoS One* 2017;**12**:e0187662. <https://doi.org/10.1371/journal.pone.0187662>
 99. Kaier TE, Twerenbold R, Puelacher C, Marjot J, Imambaccus N, Boeddinghaus J, et al. Direct comparison of cardiac myosin-binding protein C with cardiac troponins for the early diagnosis of acute myocardial infarction. *Circulation* 2017;**136**:1495–1508. <https://doi.org/10.1161/CIRCULATIONAHA.117.028084>
 100. Keller T, Tzikas S, Zeller T, Czyz E, Lillpopp L, Ojeda FM, et al. Copeptin improves early diagnosis of acute myocardial infarction. *J Am Coll Cardiol* 2010;**55**:2096–2106. <https://doi.org/10.1016/j.jacc.2010.01.029>
 101. Mockel M, Searle J, Hamm C, Slagman A, Blankenberg S, Huber K, et al. Early discharge using single cardiac troponin and copeptin testing in patients with suspected acute coronary syndrome (ACS): a randomized, controlled clinical process study. *Eur Heart J* 2015;**36**:369–376. <https://doi.org/10.1093/eurheartj/ehv178>
 102. Reichlin T, Hochholzer W, Stelzig C, Laule K, Freidank H, Morgenthaler NG, et al. Incremental value of copeptin for rapid rule out of acute myocardial infarction. *J Am Coll Cardiol* 2009;**54**:60–68. <https://doi.org/10.1016/j.jacc.2009.01.076>
 103. Boeddinghaus J, Reichlin T, Nestelberger T, Twerenbold R, Meili Y, Wildi K, et al. Early diagnosis of acute myocardial infarction in patients with mild elevations of cardiac troponin. *Clin Res Cardiol* 2017;**106**:457–467. <https://doi.org/10.1007/s00392-016-1075-9>
 104. Hillinger P, Twerenbold R, Jaeger C, Wildi K, Reichlin T, Gimenez MR, et al. Optimizing early rule-out strategies for acute myocardial infarction: utility of 1-hour copeptin. *Clin Chem* 2015;**61**:1466–1474. <https://doi.org/10.1373/clinchem.2015.242743>
 105. Mueller C, Mockel M, Giannitsis E, Huber K, Mair J, Plebani M, et al. Use of copeptin for rapid rule-out of acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2018;**7**:570–576. <https://doi.org/10.1177/2048872617710791>
 106. Mueller-Hennessen M, Lindahl B, Giannitsis E, Vafaei M, Biener M, Haushofer AC, et al. Combined testing of copeptin and high-sensitivity cardiac troponin T at presentation in comparison to other algorithms for rapid rule-out of acute myocardial infarction. *Int J Cardiol* 2019;**276**:261–267. <https://doi.org/10.1016/j.ijcard.2018.10.084>
 107. Stallone F, Schoenenberger AW, Puelacher C, Rubini Gimenez M, Walz B, Naduvilekoot Devasia A, et al. Incremental value of copeptin in suspected acute myocardial infarction very early after symptom onset. *Eur Heart J Acute Cardiovasc Care* 2016;**5**:407–415. <https://doi.org/10.1177/2048872616641289>
 108. Vargas KG, Kassem M, Mueller C, Wojta J, Huber K. Copeptin for the early rule-out of non-ST-elevation myocardial infarction. *Int J Cardiol* 2016;**223**:797–804. <https://doi.org/10.1016/j.ijcard.2016.08.304>
 109. Wildi K, Boeddinghaus J, Nestelberger T, Twerenbold R, Badertscher P, Wussler D, et al. Comparison of fourteen rule-out strategies for acute myocardial infarction. *Int J Cardiol* 2019;**283**:41–47. <https://doi.org/10.1016/j.ijcard.2018.11.140>
 110. Wildi K, Zellweger C, Twerenbold R, Jaeger C, Reichlin T, Haaf P, et al. Incremental value of copeptin to highly sensitive cardiac Troponin I for rapid rule-out of myocardial infarction. *Int J Cardiol* 2015;**190**:170–176. <https://doi.org/10.1016/j.ijcard.2015.04.133>
 111. Zellweger C, Wildi K, Twerenbold R, Reichlin T, Naduvilekoot A, Neuhaus JD, et al. Use of copeptin and high-sensitive cardiac troponin T for diagnosis and prognosis in patients with diabetes mellitus and suspected acute myocardial infarction. *Int J Cardiol* 2015;**190**:190–197. <https://doi.org/10.1016/j.ijcard.2015.04.134>
 112. Smulders MW, Kietselaer B, Wildberger JE, Dagnelie PC, Brunner-La Rocca H-P, Mingels AMA, et al. Initial imaging-guided strategy versus routine care in patients with non-ST-segment elevation myocardial infarction. *J Am Coll Cardiol* 2019;**74**:2466–2477. <https://doi.org/10.1016/j.jacc.2019.09.027>
 113. Fox K, Achenbach S, Bax J, Cosyns B, Delgado V, Dweck MR, et al. Multimodality imaging in cardiology: a statement on behalf of the Task Force on multimodality imaging of the European Association of Cardiovascular Imaging. *Eur Heart J* 2019;**40**:553–558. <https://doi.org/10.1093/eurheartj/ehy669>
 114. Gaibazzi N, Reverberi C, Badano L. Usefulness of contrast stress-echocardiography or exercise-electrocardiography to predict long-term acute coronary syndromes in patients presenting with chest pain without electrocardiographic abnormalities or 12-hour troponin elevation. *Am J Cardiol* 2011;**107**:161–167. <https://doi.org/10.1016/j.amjcard.2010.08.066>
 115. Shah BN, Balaji G, Alhajiri A, Ramzy IS, Ahmadvazir S, Senior R, et al. Incremental diagnostic and prognostic value of contemporary stress echocardiography in a chest pain unit: mortality and morbidity outcomes from a real-world setting. *Circ Cardiovasc Imaging* 2013;**6**:202–209. <https://doi.org/10.1161/CIRCIMAGING.112.980797>
 116. Sicari R, Nihoyannopoulos P, Evangelista A, Kasprzak J, Lancellotti P, Poldermans D, et al. Stress echocardiography expert consensus statement—executive summary: European Association of Echocardiography (EAE) (a registered branch of the ESC). *Eur Heart J* 2008;**30**:278–289. <https://doi.org/10.1093/eurheartj/ehn492>
 117. Ghadri JR, Wittstein IS, Prasad A, Sharkey S, Dote K, Akashi YJ, et al. International expert consensus document on Takotsubo syndrome (part II): diagnostic workup, outcome, and management. *Eur Heart J* 2018;**39**:2047–2062. <https://doi.org/10.1093/eurheartj/ehy077>
 118. Azmy C, Guerard S, Bonnet X, Gabrielli F, Skalli W. EOS orthopaedic imaging system to study patellofemoral kinematics: assessment of uncertainty. *Orthop Traumatol Surg Res* 2010;**96**:28–36. <https://doi.org/10.1016/j.rcot.2009.12.003>
 119. Hillinger P, Twerenbold R, Wildi K, Rubini Gimenez M, Jaeger C, Boeddinghaus J, et al. Gender-specific uncertainties in the diagnosis of acute coronary syndrome. *Clin Res Cardiol* 2017;**106**:28–37. <https://doi.org/10.1007/s00392-016-1020-y>
 120. Miller-Hodges E, Anand A, Shah ASV, Chapman AR, Gallacher P, Lee KK, et al. High-sensitivity cardiac troponin and the risk stratification of patients with renal impairment presenting with suspected acute coronary syndrome. *Circulation* 2018;**137**:425–435. <https://doi.org/10.1161/CIRCULATIONAHA.117.030320>

121. Gimenez MR, Twerenbold R, Boeddinghaus J, Nestelberger T, Puelacher C, Hillinger P, et al. Clinical effect of sex-specific cutoff values of high-sensitivity cardiac troponin T in suspected myocardial infarction. *JAMA Cardiol* 2016;**1**:912–920. <https://doi.org/10.1001/jamacardio.2016.2882>
122. Mueller-Hennessen M, Lindahl B, Giannitsis E, Biener M, Vafaei M, deFilippi CR, et al. Diagnostic and prognostic implications using age- and gender-specific cut-offs for high-sensitivity cardiac troponin T—sub-analysis from the TRAPID-AMI study. *Int J Cardiol* 2016;**209**:26–33. <https://doi.org/10.1016/j.ijcard.2016.01.213>
123. Sorensen NA, Neumann JT, Ojeda F, Schäfer S, Magnussen C, Keller T, et al. Relations of sex to diagnosis and outcomes in acute coronary syndrome. *J Am Heart Assoc* 2018;**7**:e007297. <https://doi.org/10.1161/JAHA.117.007297>
124. Mehta SR, Granger CB, Boden WE, Steg PG, Bassand J-P, Faxon DP, et al. Early versus delayed invasive intervention in acute coronary syndromes. *N Engl J Med* 2009;**360**:2165–2175. <https://doi.org/10.1056/NEJMoa0807986>
125. Kofoed KF, Kelbæk H, Hansen PR, Torp-Pedersen C, Høfsten D, Kløvgaard L, et al. Early versus standard care invasive examination and treatment of patients with non-ST-segment elevation acute coronary syndrome. *Circulation* 2018;**138**:2741–2750. <https://doi.org/10.1161/circulationaha.118.037152>
126. Butt JH, Kofoed KF, Kelbæk H, Hansen PR, Torp-Pedersen C, Høfsten D, et al. Importance of risk assessment in timing of invasive coronary evaluation and treatment of patients with non-ST-segment-elevation acute coronary syndrome: insights from the VERDICT trial. *J Am Heart Assoc* 2021;**10**:e022333. <https://doi.org/10.1161/jaha.121.022333>
127. Granger CB, Goldberg RJ, Dabbous O, Pieper KS, Eagle KA, Cannon CP, et al. Predictors of hospital mortality in the global registry of acute coronary events. *Arch Intern Med* 2003;**163**:2345–2353. <https://doi.org/10.1001/archinte.163.19.2345>
128. Moledina SM, Kontopantelis E, Wijeyesundera HC, Banerjee S, Van Spall HGC, Gale CP, et al. Ethnicity-dependent performance of the global registry of acute coronary events risk score for prediction of non-ST-segment elevation myocardial infarction in-hospital mortality: nationwide cohort study. *Eur Heart J* 2022;**43**:2289–2299. <https://doi.org/10.1093/eurheartj/ehac052>
129. Wallentin L, Becker RC, Budaj A, Cannon CP, Emanuelsson H, Held C, et al. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2009;**361**:1045–1057. <https://doi.org/10.1056/NEJMoa0904327>
130. Yusuf S, Zhao F, Mehta SR, Chrolavicius S, Tognoni G, Fox KK, et al. Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without ST-segment elevation. *N Engl J Med* 2001;**345**:494–502. <https://doi.org/10.1056/NEJMoa010746>
131. Puymirat E, Taldir G, Aissaoui N, Lemesle G, Lorgis L, Cuisset T, et al. Use of invasive strategy in non-ST-segment elevation myocardial infarction is a major determinant of improved long-term survival: FAST-MI (French Registry of Acute Coronary Syndrome). *JACC Cardiovasc Interv* 2012;**5**:893–902. <https://doi.org/10.1016/j.jcin.2012.05.008>
132. Bueno H, Rossello X, Pocock SJ, Van de Werf F, Chin CT, Danchin N, et al. In-hospital coronary revascularization rates and post-discharge mortality risk in non-ST-segment elevation acute coronary syndrome. *J Am Coll Cardiol* 2019;**74**:1454–1461. <https://doi.org/10.1016/j.jacc.2019.06.068>
133. Van de Werf F, Gore JM, Avezum A, Avezum Á, Gulba DC, Goodman SG, et al. Access to catheterisation facilities in patients admitted with acute coronary syndrome: multinational registry study. *BMJ* 2005;**330**:441. <https://doi.org/10.1136/bmj.383335.390718.82>
134. Bueno H. Medical management: the dark side of acute coronary syndromes. *Eur Heart J Cardiovasc Pharmacother* 2015;**1**:179–181. <https://doi.org/10.1093/ehjcvp/pvv017>
135. Giugliano RP, White JA, Bode C, Armstrong PW, Montalescot G, Lewis BS, et al. Early versus delayed, provisional eptifibatide in acute coronary syndromes. *N Engl J Med* 2009;**360**:2176–2190. <https://doi.org/10.1056/NEJMoa0901316>
136. Lindholm D, Varenhorst C, Cannon CP, Harrington RA, Himmelmann A, Maya J, et al. Ticagrelor vs. clopidogrel in patients with non-ST-elevation acute coronary syndrome with or without revascularization: results from the PLATO trial. *Eur Heart J* 2014;**35**:2083–2093. <https://doi.org/10.1093/eurheartj/ehu160>
137. Mega JL, Braunwald E, Wiwiot SD, Bassand J-P, Bhatt DL, Bode C, et al. Rivaroxaban in patients with a recent acute coronary syndrome. *N Engl J Med* 2012;**366**:9–19. <https://doi.org/10.1056/NEJMoa1112277>
138. Alexander JH, Lopes RD, James S, Kilaru R, He Y, Mohan P, et al. Apixaban with antiplatelet therapy after acute coronary syndrome. *N Engl J Med* 2011;**365**:699–708. <https://doi.org/10.1056/NEJMoa1105819>
139. Mauri L, Kereiakes DJ, Yeh RW, Driscoll-Shempp P, Cutlip DE, Steg PG, et al. Twelve or 30 months of dual antiplatelet therapy after drug-eluting stents. *N Engl J Med* 2014;**371**:2155–2166. <https://doi.org/10.1056/NEJMoa1409312>
140. Bonaca MP, Bhatt DL, Steg PG, Storey RF, Cohen M, Im K, et al. Ischaemic risk and efficacy of ticagrelor in relation to time from P2Y₁₂ inhibitor withdrawal in patients with prior myocardial infarction: insights from PEGASUS-TIMI 54. *Eur Heart J* 2016;**37**:1133–1142. <https://doi.org/10.1093/eurheartj/ehv531>
141. Bonaca MP, Bhatt DL, Cohen M, Steg PG, Storey RF, Jensen EC, et al. Long-term use of ticagrelor in patients with prior myocardial infarction. *N Engl J Med* 2015;**372**:1791–1800. <https://doi.org/10.1056/NEJMoa1500857>
142. Steg PG, Bhatt DL, Simon T, Fox K, Mehta SR, Harrington RA, et al. Ticagrelor in patients with stable coronary disease and diabetes. *N Engl J Med* 2019;**381**:1309–1320. <https://doi.org/10.1056/NEJMoa1908077>
143. Eikelboom JW, Connolly SJ, Bosch J, Dagenais GR, Hart RG, Shestakovsky O, et al. Rivaroxaban with or without aspirin in stable cardiovascular disease. *N Engl J Med* 2017;**377**:1319–1330. <https://doi.org/10.1056/NEJMoa1709118>
144. Connolly SJ, Eikelboom JW, Bosch J, Dagenais G, Dyal L, Lanais F, et al. Rivaroxaban with or without aspirin in patients with stable coronary artery disease: an international, randomised, double-blind, placebo-controlled trial. *Lancet* 2018;**391**:205–218. [https://doi.org/10.1016/S0140-6736\(17\)32458-3](https://doi.org/10.1016/S0140-6736(17)32458-3)
145. Fox KA, Mehta SR, Peters R, Zhao F, Lakkis N, Gersh BJ, et al. Benefits and risks of the combination of clopidogrel and aspirin in patients undergoing surgical revascularization for non-ST-elevation acute coronary syndrome: the Clopidogrel in Unstable angina to prevent Recurrent ischemic Events (CURE) trial. *Circulation* 2004;**110**:1202–1208. <https://doi.org/10.1161/01.Cir.0000140675.85342.1b>
146. Held C, Asenblad N, Bassand JP, Becker RC, Cannon CP, Claeys MJ, et al. Ticagrelor versus clopidogrel in patients with acute coronary syndromes undergoing coronary artery bypass surgery: results from the PLATO (Platelet Inhibition and Patient Outcomes) trial. *J Am Coll Cardiol* 2011;**57**:672–684. <https://doi.org/10.1016/j.jacc.2010.10.029>
147. Dewilde WJ, Oirbans T, Verheugt FW, Kelder JC, De Smet BJGL, Herrman J-P, et al. Use of clopidogrel with or without aspirin in patients taking oral anticoagulant therapy and undergoing percutaneous coronary intervention: an open-label, randomised, controlled trial. *Lancet* 2013;**381**:1107–1115. [https://doi.org/10.1016/S0140-6736\(12\)62177-1](https://doi.org/10.1016/S0140-6736(12)62177-1)
148. Fiedler KA, Maeng M, Mehilli J, Schulz-Schüpke S, Byrne RA, Sibbing D, et al. Duration of triple therapy in patients requiring oral anticoagulation after drug-eluting stent implantation: the ISAR-TRIPLE trial. *J Am Coll Cardiol* 2015;**65**:1619–1629. <https://doi.org/10.1016/j.jacc.2015.02.050>
149. Gibson CM, Mehran R, Bode C, Halperin J, Verheugt FW, Willgoose P, et al. Prevention of bleeding in patients with atrial fibrillation undergoing PCI. *N Engl J Med* 2016;**375**:2423–2434. <https://doi.org/10.1056/NEJMoa1611594>
150. Cannon CP, Bhatt DL, Oldgren J, Lip GYH, Ellis SG, Kimura T, et al. Dual antithrombotic therapy with dabigatran after PCI in atrial fibrillation. *N Engl J Med* 2017;**377**:1513–1524. <https://doi.org/10.1056/NEJMoa1708454>
151. Lopes RD, Heizer G, Aronson R, Vora AN, Massaro T, Mehran R, et al. Antithrombotic therapy after acute coronary syndrome or PCI in atrial fibrillation. *N Engl J Med* 2019;**380**:1509–1524. <https://doi.org/10.1056/NEJMoa1817083>
152. Vranckx P, Valgimigli M, Eckardt L, Tijssen J, Lewalter T, Gargiulo G, et al. Edoxaban-based versus vitamin K antagonist-based antithrombotic regimen after successful coronary stenting in patients with atrial fibrillation (ENTRUST-AF PCI): a randomised, open-label, phase 3b trial. *Lancet* 2019;**394**:1335–1343. [https://doi.org/10.1016/S0140-6736\(19\)31872-0](https://doi.org/10.1016/S0140-6736(19)31872-0)
153. Armstrong PW, Gershlick AH, Goldstein P, Wilcox R, Danays T, Lambert Y, et al. Fibrinolysis or primary PCI in ST-segment elevation myocardial infarction. *N Engl J Med* 2013;**368**:1379–1387. <https://doi.org/10.1056/NEJMoa1301092>
154. GUSTO Investigators. An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. *N Engl J Med* 1993;**329**:673–682. <https://doi.org/10.1056/nejm199309023291001>
155. Van De Werf F, Adegbe J, Ardissino D, Armstrong PW, Aylward P, Barbash G, et al. Single-bolus tenecteplase compared with front-loaded alteplase in acute myocardial infarction: the ASSENT-2 double-blind randomised trial. *Lancet* 1999;**354**:716–722. [https://doi.org/10.1016/S0140-6736\(99\)07403-6](https://doi.org/10.1016/S0140-6736(99)07403-6)
156. Group FTFC. Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients. *Lancet* 1994;**343**:311–322. [https://doi.org/10.1016/S0140-6736\(94\)91161-4](https://doi.org/10.1016/S0140-6736(94)91161-4)
157. Van de Werf F, Barron HV, Armstrong PW, Granger CB, Berlioli S, Barbash G, et al. Incidence and predictors of bleeding events after fibrinolytic therapy with fibrin-specific agents: a comparison of TNK-tPA and rt-PA. *Eur Heart J* 2001;**22**:2253–2261. <https://doi.org/10.1053/ehuj.2001.2686>
158. White HD. Thrombolytic therapy in the elderly. *Lancet* 2000;**356**:2028–2030. [https://doi.org/10.1016/S0140-6736\(00\)03397-3](https://doi.org/10.1016/S0140-6736(00)03397-3)
159. Investigators TGUStOOAGI. A comparison of reteplase with alteplase for acute myocardial infarction. *N Engl J Med* 1997;**337**:1118–1123. <https://doi.org/10.1056/nejm199710163371603>
160. Böttiger BW, Arntz HR, Chamberlain DA, Bluhmki E, Belmans A, Danays T, et al. Thrombolysis during resuscitation for out-of-hospital cardiac arrest. *N Engl J Med* 2008;**359**:2651–2662. <https://doi.org/10.1056/NEJMoa070570>
161. James SK, Roe MT, Cannon CP, Cornel JH, Horrow J, Husted S, et al. Ticagrelor versus clopidogrel in patients with acute coronary syndromes intended for non-invasive management: substudy from prospective randomised PLATElet inhibition and patient Outcomes (PLATO) trial. *BMJ* 2011;**342**:d3527. <https://doi.org/10.1136/bmj.d3527>
162. Husted S, James S, Becker RC, Horrow J, Katus H, Storey RF, et al. Ticagrelor versus clopidogrel in elderly patients with acute coronary syndromes: a substudy from the prospective randomized PLATElet inhibition and patient Outcomes (PLATO) trial.

- Circ Cardiovasc Qual Outcomes* 2012;**5**:680–688. <https://doi.org/10.1161/circoutcomes.111.964395>
163. Szummer K, Montez-Rath ME, Alfredsson J, Erlinge D, Lindahl B, Hofmann R, et al. Comparison between ticagrelor and clopidogrel in elderly patients with an acute coronary syndrome: insights from the SWEDEHEART registry. *Circulation* 2020;**142**: 1700–1708. <https://doi.org/10.1161/circulationaha.120.050645>
 164. Gimbel M, Qaderdan K, Willemsen L, Hermanides R, Bergmeijer T, de Vrey E, et al. Clopidogrel versus ticagrelor or prasugrel in patients aged 70 years or older with non-ST-elevation acute coronary syndrome (POPular AGE): the randomised, open-label, non-inferiority trial. *Lancet* 2020;**395**:1374–1381. [https://doi.org/10.1016/s0140-6736\(20\)30325-1](https://doi.org/10.1016/s0140-6736(20)30325-1)
 165. Sinning C, Ahrens I, Cariou A, Beygui F, Lamhaut L, Halvorsen S, et al. The cardiac arrest centre for the treatment of sudden cardiac arrest due to presumed cardiac cause—aims, function and structure: position paper of the association for Acute CardioVascular Care of the European Society of Cardiology (AVCV), European Association of Percutaneous Coronary Interventions (EAPCI), European Heart Rhythm Association (EHRA), European Resuscitation Council (ERC), European Society for Emergency Medicine (EUSEM) and European Society of Intensive Care Medicine (ESICM). *Eur Heart J Acute Cardiovasc Care* 2020;**9**:S193–S202. <https://doi.org/10.1177/2048872620963492>
 166. Yeung J, Matsuyama T, Bray J, Reynolds J, Skrifvars MB. Does care at a cardiac arrest centre improve outcome after out-of-hospital cardiac arrest?—A systematic review. *Resuscitation* 2019;**137**:102–115. <https://doi.org/10.1016/j.resuscitation.2019.02.006>
 167. Wnent J, Seewald S, Heringlake M, Lemke H, Brauer K, Lefering R, et al. Choice of hospital after out-of-hospital cardiac arrest—a decision with far-reaching consequences: a study in a large German city. *Crit Care* 2012;**16**:R164. <https://doi.org/10.1186/cc11516>
 168. Kragholm K, Malta Hansen C, Dupre ME, Xian Y, Strauss B, Tyson C, et al. Direct transport to a percutaneous cardiac intervention center and outcomes in patients with out-of-hospital cardiac arrest. *Circ Cardiovasc Qual Outcomes* 2017;**10**:e003414. <https://doi.org/10.1161/circoutcomes.116.003414>
 169. Dafaalla M, Rashid M, Sun L, Quinn T, Timmis A, Wijeyesundera H, et al. Impact of availability of catheter laboratory facilities on management and outcomes of acute myocardial infarction presenting with out of hospital cardiac arrest. *Resuscitation* 2022;**170**: 327–334. <https://doi.org/10.1016/j.resuscitation.2021.10.031>
 170. Kelham M, Jones TN, Rathod KS, Guttman O, Proudfoot A, Rees P, et al. An observational study assessing the impact of a cardiac arrest centre on patient outcomes after out-of-hospital cardiac arrest (OHCA). *Eur Heart J Acute Cardiovasc Care* 2020;**9**: S67–S73. <https://doi.org/10.1177/2048872620974606>
 171. Bonnefoy-Cudraz E, Bueno H, Casella G, De Maria E, Fitzsimons D, Halvorsen S, et al. Editor's choice—acute cardiovascular care association position paper on intensive cardiovascular care units: an update on their definition, structure, organisation and function. *Eur Heart J Acute Cardiovasc Care* 2018;**7**:80–95. <https://doi.org/10.1177/2048872617724269>
 172. van Diepen S, Katz JN, Morrow DA. Will cardiac intensive care unit admissions warrant appropriate use criteria in the future? *Circulation* 2019;**140**:267–269. <https://doi.org/10.1161/circulationaha.118.039125>
 173. Claeys MJ, Roubille F, Casella G, Zukermann R, Nikolaou N, De Luca L, et al. Organization of intensive cardiac care units in Europe: results of a multinational survey. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:993–1001. <https://doi.org/10.1177/2048872619883997>
 174. Melberg T, Jørgensen M, Ørn S, Solli T, Edland U, Dickstein K. Safety and health status following early discharge in patients with acute myocardial infarction treated with primary PCI: a randomized trial. *Eur J Prev Cardiol* 2015;**22**:1427–1434. <https://doi.org/10.1177/2047487314559276>
 175. Tickoo S, Bhardwaj A, Fonarow GC, Liang L, Bhatt DL, Cannon CP. Relation between hospital length of stay and quality of care in patients with acute coronary syndromes (from the American Heart Association's get with the guidelines—coronary artery disease data set). *Am J Cardiol* 2016;**117**:201–205. <https://doi.org/10.1016/j.amjcard.2015.10.027>
 176. Eagle KA, Lim MJ, Dabbous OH, Pieper KS, Goldberg RJ, Van de Werf F, et al. A validated prediction model for all forms of acute coronary syndrome: estimating the risk of 6-month postdischarge death in an international registry. *JAMA* 2004;**291**: 2727–2733. <https://doi.org/10.1001/jama.291.22.2727>
 177. Fox KA, Dabbous OH, Goldberg RJ, Pieper KS, Eagle KA, Van de Werf F, et al. Prediction of risk of death and myocardial infarction in the six months after presentation with acute coronary syndrome: prospective multinational observational study (GRACE). *BMJ* 2006;**333**:1091. <https://doi.org/10.1136/bmj.38985.646481.55>
 178. Fox KA, Fitzgerald G, Puymirat E, Huang W, Carruthers K, Simon T, et al. Should patients with acute coronary disease be stratified for management according to their risk? Derivation, external validation and outcomes using the updated GRACE risk score. *BMJ Open* 2014;**4**:e004425. <https://doi.org/10.1136/bmjopen-2013-004425>
 179. Fox KA, Eagle KA, Gore JM, Steg PG, Anderson FA. The global registry of acute coronary events, 1999 to 2009—GRACE. *Heart* 2010;**96**:1095–1101. <https://doi.org/10.1136/hrt.2009.190827>
 180. Bing R, Goodman SG, Yan AT, Fox K, Gale CP, Hyun K, et al. Use of clinical risk stratification in non-ST elevation acute coronary syndromes: an analysis from the CONCORDANCE registry. *Eur Heart J Qual Care Clin Outcomes* 2018;**4**:309–317. <https://doi.org/10.1093/ehjqcco/qcy002>
 181. Chew DP, Junbo G, Parsonage W, Kerker P, Sulimov VA, Horsfall M, et al. Perceived risk of ischemic and bleeding events in acute coronary syndromes. *Circ Cardiovasc Qual Outcomes* 2013;**6**:299–308. <https://doi.org/10.1161/circoutcomes.111.000072>
 182. Hall M, Bebb OJ, Dondo TB, Yan AT, Goodman SG, Bueno H, et al. Guideline-indicated treatments and diagnostics, GRACE risk score, and survival for non-ST elevation myocardial infarction. *Eur Heart J* 2018;**39**:3798–3806. <https://doi.org/10.1093/eurheartj/ehy517>
 183. Fox KA, Anderson FA Jr, Dabbous OH, Steg PG, Lopez-Sendon J, Van de Werf F, et al. Intervention in acute coronary syndromes: do patients undergo intervention on the basis of their risk characteristics? The Global Registry of Acute Coronary Events (GRACE). *Heart* 2007;**93**:177–182. <https://doi.org/10.1136/hrt.2005.084830>
 184. Saar A, Marandi T, Ainla T, Fischer K, Blöndal M, Eha J. The risk-treatment paradox in non-ST-elevation myocardial infarction patients according to their estimated GRACE risk. *Int J Cardiol* 2018;**272**:26–32. <https://doi.org/10.1016/j.ijcard.2018.08.015>
 185. Chew DP, Astley CM, Luker H, Alprandi-Costa B, Hillis G, Chow CK, et al. A cluster randomized trial of objective risk assessment versus standard care for acute coronary syndromes: rationale and design of the Australian GRACE Risk score Intervention Study (AGRIS). *Am Heart J* 2015;**170**:995–1004.e1. <https://doi.org/10.1016/j.ahj.2015.07.032>
 186. Everett CC, Fox KA, Reynolds C, Fernandez C, Sharples L, Stocken DD, et al. Evaluation of the impact of the GRACE risk score on the management and outcome of patients hospitalised with non-ST elevation acute coronary syndrome in the UK: protocol of the UKGRIS cluster-randomised registry-based trial. *BMJ Open* 2019;**9**: e032165. <https://doi.org/10.1136/bmjopen-2019-032165>
 187. Chew DP, Hyun K, Morton E, Horsfall M, Hillis GS, Chow CK, et al. Objective risk assessment vs standard care for acute coronary syndromes: a randomized clinical trial. *JAMA Cardiol* 2021;**6**:304–313. <https://doi.org/10.1001/jamacardio.2020.6314>
 188. Hall M, Dondo TB, Yan AT, Goodman SG, Bueno H, Chew DP, et al. Association of clinical factors and therapeutic strategies with improvements in survival following non-ST-elevation myocardial infarction, 2003–2013. *JAMA* 2016;**316**:1073–1082. <https://doi.org/10.1001/jama.2016.10766>
 189. Bricker RS, Valle JA, Plomondon ME, Armstrong EJ, Waldo SW. Causes of mortality after percutaneous coronary intervention. *Circ Cardiovasc Qual Outcomes* 2019;**12**: e005355. <https://doi.org/10.1161/circoutcomes.118.005355>
 190. Chang WC, Boersma E, Granger CB, Harrington RA, Califf RM, Simoons ML, et al. Dynamic prognostication in non-ST-elevation acute coronary syndromes: insights from GUSTO-III and PURSUIT. *Am Heart J* 2004;**148**:62–71. <https://doi.org/10.1016/j.ahj.2003.05.004>
 191. Fox KA, Anderson FA Jr, Goodman SG, Steg PG, Pieper K, Quill A, et al. Time course of events in acute coronary syndromes: implications for clinical practice from the GRACE registry. *Nat Clin Pract Cardiovasc Med* 2008;**5**:580–589. <https://doi.org/10.1038/npcardio1302>
 192. Spoon DB, Psaltis PJ, Singh M, Holmes DR, Gersh BJ, Rihal CS, et al. Trends in cause of death after percutaneous coronary intervention. *Circulation* 2014;**129**:1286–1294. <https://doi.org/10.1161/circulationaha.113.006518>
 193. Simms AD, Reynolds S, Pieper K, Baxter PD, Cattle BA, Batin PD, et al. Evaluation of the NICE mini-GRACE risk scores for acute myocardial infarction using the Myocardial Ischaemia National Audit Project (MINAP) 2003–2009: National Institute for Cardiovascular Outcomes Research (NICOR). *Heart* 2013;**99**:35–40. <https://doi.org/10.1136/heartjnl-2012-302632>
 194. Meune C, Drexler B, Haaf P, Reichlin T, Reiter M, Meissner J, et al. The GRACE score's performance in predicting in-hospital and 1-year outcome in the era of high-sensitivity cardiac troponin assays and B-type natriuretic peptide. *Heart* 2011;**97**:1479–1483. <https://doi.org/10.1136/hrt.2010.220988>
 195. Elbarouni B, Goodman SG, Yan RT, Welsh RC, Kornder JM, DeYoung JP, et al. Validation of the Global Registry of Acute Coronary Event (GRACE) risk score for in-hospital mortality in patients with acute coronary syndrome in Canada. *Am Heart J* 2009;**158**:392–399. <https://doi.org/10.1016/j.ahj.2009.06.010>
 196. Lin A, Devlin G, Lee M, Kerr AJ. Performance of the GRACE scores in a New Zealand acute coronary syndrome cohort. *Heart* 2014;**100**:1960–1966. <https://doi.org/10.1136/heartjnl-2014-306062>
 197. Wenzl FA, Kraler S, Ambler W, Weston C, Herzog SA, Räber L, et al. Sex-specific evaluation and redevelopment of the GRACE score in non-ST-segment elevation acute coronary syndromes in populations from the UK and Switzerland: a multinational analysis with external cohort validation. *Lancet* 2022;**400**:744–756. [https://doi.org/10.1016/s0140-6736\(22\)01483-0](https://doi.org/10.1016/s0140-6736(22)01483-0)
 198. Redfors B, Chen S, Crowley A, Ben-Yehuda O, Gersh BJ, Lembo NJ, et al. B-Type natriuretic peptide assessment in patients undergoing revascularization for left main coronary artery disease. *Circulation* 2018;**138**:469–478. <https://doi.org/10.1161/CIRCULATIONAHA.118.033631>
 199. Michou E, Fahri G, Mueller C. Quantifying heart failure using natriuretic peptides may help the HEART team in decision-making. *Eur Heart J* 2019;**40**:3406–3408. <https://doi.org/10.1093/eurheartj/ehz572>

200. Zhang C, Jiang L, Xu L, Tian J, Liu J, Zhao X, et al. Implications of N-terminal pro-B-type natriuretic peptide in patients with three-vessel disease. *Eur Heart J* 2019;**40**: 3397–3405. <https://doi.org/10.1093/eurheartj/ehz394>
201. Shen S, Ye J, Wu X, Li X. Association of N-terminal pro-brain natriuretic peptide level with adverse outcomes in patients with acute myocardial infarction: a meta-analysis. *Heart Lung* 2021;**50**:863–869. <https://doi.org/10.1016/j.hrtlung.2021.07.007>
202. Thygesen K, Mair J, Mueller C, Huber K, Weber M, Plebani M, et al. Recommendations for the use of natriuretic peptides in acute cardiac care: a position statement from the Study Group on Biomarkers in Cardiology of the ESC Working Group on Acute Cardiac Care. *Eur Heart J* 2012;**33**:2001–2006. <https://doi.org/10.1093/eurheartj/ehq509>
203. de Lemos JA, Morrow DA, Bentley JH, Omland T, Sabatine MS, McCabe CH, et al. The prognostic value of B-type natriuretic peptide in patients with acute coronary syndromes. *N Engl J Med* 2001;**345**:1014–1021. <https://doi.org/10.1056/NEJMoa011053>
204. Morrow DA, de Lemos JA, Sabatine MS, Murphy SA, Demopoulos LA, DiBattiste PM, et al. Evaluation of B-type natriuretic peptide for risk assessment in unstable angina/non-ST-elevation myocardial infarction: B-type natriuretic peptide and prognosis in TACTICS-TIMI 18. *J Am Coll Cardiol* 2003;**41**:1264–1272. [https://doi.org/10.1016/s0735-1097\(03\)00168-2](https://doi.org/10.1016/s0735-1097(03)00168-2)
205. O'Donoghue M, de Lemos JA, Morrow DA, Murphy SA, Buros JL, Cannon CP, et al. Prognostic utility of heart-type fatty acid binding protein in patients with acute coronary syndromes. *Circulation* 2006;**114**:550–557. <https://doi.org/10.1161/circulationaha.106.641936>
206. Balmelli C, Meune C, Twerenbold R, Reichlin T, Rieder S, Drexler B, et al. Comparison of the performances of cardiac troponins, including sensitive assays, and copeptin in the diagnostic of acute myocardial infarction and long-term prognosis between women and men. *Am Heart J* 2013;**166**:30–37. <https://doi.org/10.1016/j.ahj.2013.03.014>
207. Dhillon OS, Khan SQ, Narayan HK, Ng KH, Struck J, Quinn PA, et al. Prognostic value of mid-regional pro-adrenomedullin levels taken on admission and discharge in non-ST-elevation myocardial infarction: the LAMP (Leicester Acute Myocardial Infarction Peptide) II study. *J Am Coll Cardiol* 2010;**56**:125–133. <https://doi.org/10.1016/j.jacc.2010.01.060>
208. Maisel A, Mueller C, Neath SX, Christenson RH, Morgenthaler NG, McCord J, et al. Copeptin helps in the early detection of patients with acute myocardial infarction: primary results of the CHOPIN trial (Copeptin Helps in the early detection Of Patients with acute myocardial infarction). *J Am Coll Cardiol* 2013;**62**:150–160. <https://doi.org/10.1016/j.jacc.2013.04.011>
209. Sabatine MS, Morrow DA, de Lemos JA, Gibson CM, Murphy SA, Rifai N, et al. Multimarker approach to risk stratification in non-ST elevation acute coronary syndromes: simultaneous assessment of troponin I, C-reactive protein, and B-type natriuretic peptide. *Circulation* 2002;**105**:1760–1763. <https://doi.org/10.1161/01.cir.0000015464.18023.0a>
210. Wollert KC, Kempf T, Peter T, Olofsson S, James S, Johnston N, et al. Prognostic value of growth-differentiation factor-15 in patients with non-ST-elevation acute coronary syndrome. *Circulation* 2007;**115**:962–971. <https://doi.org/10.1161/circulationaha.106.650846>
211. Subherwal S, Bach RG, Chen AY, Gage BF, Rao SV, Newby LK, et al. Baseline risk of major bleeding in non-ST-segment-elevation myocardial infarction: the CRUSADE (Can Rapid risk stratification of Unstable angina patients Suppress ADverse outcomes with Early implementation of the ACC/AHA Guidelines) bleeding score. *Circulation* 2009;**119**:1873–1882. <https://doi.org/10.1161/circulationaha.108.828541>
212. Abu-Assi E, Raposeiras-Roubin S, Lear P, Cabanas-Grandio P, Gironde M, Rodríguez-Cordero M, et al. Comparing the predictive validity of three contemporary bleeding risk scores in acute coronary syndrome. *Eur Heart J Acute Cardiovasc Care* 2012;**1**:222–231. <https://doi.org/10.1177/2048872612453924>
213. Mehran R, Pocock SJ, Nikolsky E, Clayton T, Dangas GD, Kirtane AJ, et al. A risk score to predict bleeding in patients with acute coronary syndromes. *J Am Coll Cardiol* 2010;**55**:2556–2566. <https://doi.org/10.1016/j.jacc.2009.09.076>
214. Urban P, Mehran R, Collieran R, Angiolillo DJ, Byrne RA, Capodanno D, et al. Defining high bleeding risk in patients undergoing percutaneous coronary intervention: a consensus document from the academic research consortium for high bleeding risk. *Eur Heart J* 2019;**40**:2632–2653. <https://doi.org/10.1093/eurheartj/ehz372>
215. Natsuaki M, Morimoto T, Shiomi H, Yamaji K, Watanabe H, Shizuta S, et al. Application of the academic research consortium high bleeding risk criteria in an all-comers registry of percutaneous coronary intervention. *Circ Cardiovasc Interv* 2019;**12**:e008307. <https://doi.org/10.1161/circinterventions.119.008307>
216. Ueki Y, Bär S, Losdat S, Otsuka T, Zanchin C, Zanchin T, et al. Validation of the Academic Research Consortium for High Bleeding Risk (ARC-HBR) criteria in patients undergoing percutaneous coronary intervention and comparison with contemporary bleeding risk scores. *Eurointervention* 2020;**16**:371–379. <https://doi.org/10.4244/eij-d-20-00052>
217. Capodanno D, Morice MC, Angiolillo DJ, Bhatt DL, Byrne RA, Collieran R, et al. Trial design principles for patients at high bleeding risk undergoing PCI: JACC scientific expert panel. *J Am Coll Cardiol* 2020;**76**:1468–1483. <https://doi.org/10.1016/j.jacc.2020.06.085>
218. Urban P, Meredith IT, Abizaid A, Pocock SJ, Carrié D, Naber C, et al. Polymer-free drug-coated coronary stents in patients at high bleeding risk. *N Engl J Med* 2015;**373**:2038–2047. <https://doi.org/10.1056/NEJMoa1503943>
219. Valgimigli M, Patilalakas A, Thury A, McFadden E, Colangelo S, Campo G, et al. Zotarolimus-eluting versus bare-metal stents in uncertain drug-eluting stent candidates. *J Am Coll Cardiol* 2015;**65**:805–815. <https://doi.org/10.1016/j.jacc.2014.11.053>
220. Varenne O, Cook S, Sideris G, Kedev S, Cuisset T, Carrié D, et al. Drug-eluting stents in elderly patients with coronary artery disease (SENIOR): a randomised single-blind trial. *Lancet* 2018;**391**:41–50. [https://doi.org/10.1016/s0140-6736\(17\)32713-7](https://doi.org/10.1016/s0140-6736(17)32713-7)
221. Yeh RW, Secemsky EA, Kereiakes DJ, Normand S-LT, Gershlick AH, Cohen DJ, et al. Development and validation of a prediction rule for benefit and harm of dual antiplatelet therapy beyond 1 year after percutaneous coronary intervention. *JAMA* 2016;**315**:1735–1749. <https://doi.org/10.1001/jama.2016.3775>
222. Costa F, van Klaveren D, James S, Heg D, Räber L, Feres F, et al. Derivation and validation of the predicting bleeding complications in patients undergoing stent implantation and subsequent dual antiplatelet therapy (PRECISE-DAPT) score: a pooled analysis of individual-patient datasets from clinical trials. *Lancet* 2017;**389**:1025–1034. [https://doi.org/10.1016/s0140-6736\(17\)30397-5](https://doi.org/10.1016/s0140-6736(17)30397-5)
223. Bianco M, D'Ascenzo F, Raposeiras Roubin S, Kinnaird T, Peyracchia M, Ariza-Solé A, et al. Comparative external validation of the PRECISE-DAPT and PARIS risk scores in 4424 acute coronary syndrome patients treated with prasugrel or ticagrelor. *Int J Cardiol* 2020;**301**:200–206. <https://doi.org/10.1016/j.ijcard.2019.11.132>
224. Piccolo R, Gargiulo G, Franzone A, Santucci A, Ariotti S, Baldo A, et al. Use of the dual-antiplatelet therapy score to guide treatment duration after percutaneous coronary intervention. *Ann Intern Med* 2017;**167**:17–25. <https://doi.org/10.7326/m16-2389>
225. Lambert LJ, Brophy JM, Racine N, Rinfret S, L'Allier PL, Brown KA, et al. Outcomes of patients with ST-elevation myocardial infarction receiving and not receiving reperfusion therapy: the importance of examining all patients. *Can J Cardiol* 2016;**32**:1325.e11–1325.e18. <https://doi.org/10.1016/j.cjca.2016.02.073>
226. Hong SJ, Kim BK, Shin DH, Nam C-M, Kim J-S, Ko Y-G, et al. Effect of intravascular ultrasound-guided vs angiography-guided everolimus-eluting stent implantation: the IVUS-XPL randomized clinical trial. *JAMA* 2015;**314**:2155–2163. <https://doi.org/10.1001/jama.2015.15454>
227. Zhang J, Gao X, Kan J, Ge Z, Han L, Lu S, et al. Intravascular ultrasound versus angiography-guided drug-eluting stent implantation: the ULTIMATE trial. *J Am Coll Cardiol* 2018;**72**:3126–3137. <https://doi.org/10.1016/j.jacc.2018.09.013>
228. Gao XF, Ge Z, Kong XQ, Kan J, Han L, Lu S, et al. 3-Year outcomes of the ULTIMATE trial comparing intravascular ultrasound versus angiography-guided drug-eluting stent implantation. *JACC Cardiovasc Interv* 2021;**14**:247–257. <https://doi.org/10.1016/j.jcin.2020.10.001>
229. Meneveau N, Souteyrand G, Motreff P, Caussin C, Amabile N, Ohlmann P, et al. Optical coherence tomography to optimize results of percutaneous coronary intervention in patients with non-ST-elevation acute coronary syndrome: results of the multicenter, randomized DOCTORS study (Does Optical Coherence Tomography Optimize Results of Stenting). *Circulation* 2016;**134**:906–917. <https://doi.org/10.1161/CIRCULATIONAHA.116.024393>
230. Kala P, Cervinka P, Jakl M, Kanovsky J, Kupec A, Spacek R, et al. OCT Guidance during stent implantation in primary PCI: a randomized multicenter study with nine months of optical coherence tomography follow-up. *Int J Cardiol* 2018;**250**:98–103. <https://doi.org/10.1016/j.ijcard.2017.10.059>
231. Ali ZA, Maehara A, Genereux P, Shlofmitz RA, Fabbiochi F, Nazif TM, et al. Optical coherence tomography compared with intravascular ultrasound and with angiography to guide coronary stent implantation (ILUMIEN III: OPTIMIZE PCI): a randomised controlled trial. *Lancet* 2016;**388**:2618–2628. [https://doi.org/10.1016/S0140-6736\(16\)31922-5](https://doi.org/10.1016/S0140-6736(16)31922-5)
232. Ali ZA, Karimi Galougahi K, Maehara A, Shlofmitz RA, Fabbiochi F, Guagliumi G, et al. Outcomes of optical coherence tomography compared with intravascular ultrasound and with angiography to guide coronary stent implantation: one-year results from the ILUMIEN III: OPTIMIZE PCI trial. *Eurointervention* 2021;**16**:1085–1091. <https://doi.org/10.4244/eij-d-20-00498>
233. Kubo T, Shinko T, Okamura T, Hibi K, Nakazawa G, Morino Y, et al. Optical frequency domain imaging vs. intravascular ultrasound in percutaneous coronary intervention (OPINION trial): one-year angiographic and clinical results. *Eur Heart J* 2017;**38**:3139–3147. <https://doi.org/10.1093/eurheartj/ehx351>
234. Carrick D, Haig C, Carberry J, May VTY, McCartney P, Welsh P, et al. Microvascular resistance of the culprit coronary artery in acute ST-elevation myocardial infarction. *JCI Insight* 2016;**1**:e85768. <https://doi.org/10.1172/jci.insight.85768>
235. Cuculi F, Dall'Armellina E, Manhiot C, De Caterina AR, Colyer S, Ferreira V, et al. Early change in invasive measures of microvascular function can predict myocardial recovery following PCI for ST-elevation myocardial infarction. *Eur Heart J* 2014;**35**:1971–1980. <https://doi.org/10.1093/eurheartj/ehz434>
236. Carrick D, Haig C, Ahmed N, Carberry J, Yue May VT, McEntegart M, et al. Comparative prognostic utility of indexes of microvascular function alone or in combination in patients with an acute ST-segment-elevation myocardial infarction. *Circulation* 2016;**134**:1833–1847. <https://doi.org/10.1161/circulationaha.116.022603>

237. Murai T, Yonetsu T, Kanaji Y, Usui E, Hoshino M, Hada M, et al. Prognostic value of the index of microcirculatory resistance after percutaneous coronary intervention in patients with non-ST-segment elevation acute coronary syndrome. *Catheter Cardiovasc Interv* 2018;**92**:1063–1074. <https://doi.org/10.1002/ccd.27529>
238. Maznycka AM, Oldroyd KG, McCartney P, McEntegart M, Berry C. The potential use of the index of microcirculatory resistance to guide stratification of patients for adjunctive therapy in acute myocardial infarction. *JACC Cardiovasc Interv* 2019;**12**:951–966. <https://doi.org/10.1016/j.jcin.2019.01.246>
239. van Leeuwen MAH, van der Hoeven NW, Janssens GN, Everaars H, Nap A, Lemkes JS, et al. Evaluation of microvascular injury in revascularized patients with ST-segment-elevation myocardial infarction treated with ticagrelor versus prasugrel. *Circulation* 2019;**139**:636–646. <https://doi.org/10.1161/circulationaha.118.035931>
240. Maznycka AM, McCartney PJ, Oldroyd KG, Lindsay M, McEntegart M, Eteiba H, et al. Effects of intracoronary alteplase on microvascular function in acute myocardial infarction. *J Am Heart Assoc* 2020;**9**:e014066. <https://doi.org/10.1161/jaha.119.014066>
241. Kapur NK, Paruchuri V, Urbano-Morales JA, Mackey EE, Daly GH, Qiao X, et al. Mechanically unloading the left ventricle before coronary reperfusion reduces left ventricular wall stress and myocardial infarct size. *Circulation* 2013;**128**:328–336. <https://doi.org/10.1161/circulationaha.112.000029>
242. Watanabe S, Fish K, Kovacic JC, Bikou O, Leonardson L, Nomoto K, et al. Left ventricular unloading using an impella CP improves coronary flow and infarct zone perfusion in ischemic heart failure. *J Am Heart Assoc* 2018;**7**:e006462. <https://doi.org/10.1161/jaha.117.006462>
243. Esposito ML, Zhang Y, Qiao X, Reyelt L, Paruchuri V, Schnitzler GR, et al. Left ventricular unloading before reperfusion promotes functional recovery after acute myocardial infarction. *J Am Coll Cardiol* 2018;**72**:501–514. <https://doi.org/10.1016/j.jacc.2018.05.034>
244. Patel MR, Smalling RW, Thiele H, Barnhart HX, Zhou Y, Chandra P, et al. Intra-aortic balloon counterpulsation and infarct size in patients with acute anterior myocardial infarction without shock: the CRISP AMI randomized trial. *JAMA* 2011;**306**:1329–1337. <https://doi.org/10.1001/jama.2011.1280>
245. Kapur NK, Alkhouli MA, DeMartini TJ, Faraz H, George ZH, Goodwin MJ, et al. Unloading the left ventricle before reperfusion in patients with anterior ST-segment-elevation myocardial infarction. *Circulation* 2019;**139**:337–346. <https://doi.org/10.1161/circulationaha.118.038269>
246. Erlinge D. A review of mild hypothermia as an adjunctive treatment for ST-elevation myocardial infarction. *Ther Hypothermia Temp Manag* 2011;**1**:129–141. <https://doi.org/10.1089/ther.2011.0008>
247. Dixon SR, Whitbourn RJ, Dae MW, Nguyen TT, O'Neill VV, Grube E, et al. Induction of mild systemic hypothermia with endovascular cooling during primary percutaneous coronary intervention for acute myocardial infarction. *J Am Coll Cardiol* 2002;**40**:1928–1934. [https://doi.org/10.1016/s0735-1097\(02\)02567-6](https://doi.org/10.1016/s0735-1097(02)02567-6)
248. Götzberg M, Olivecrona GK, Koul S, Carlsson M, Engblom H, Ugander M, et al. A pilot study of rapid cooling by cold saline and endovascular cooling before reperfusion in patients with ST-elevation myocardial infarction. *Circ Cardiovasc Interv* 2010;**3**:400–407. <https://doi.org/10.1161/circinterventions.110.957902>
249. Erlinge D, Götzberg M, Lang I, Holzer M, Noc M, Clemmensen P, et al. Rapid endovascular catheter core cooling combined with cold saline as an adjunct to percutaneous coronary intervention for the treatment of acute myocardial infarction. The CHILL-MI trial: a randomized controlled study of the use of central venous catheter core cooling combined with cold saline as an adjunct to percutaneous coronary intervention for the treatment of acute myocardial infarction. *J Am Coll Cardiol* 2014;**63**:1857–1865. <https://doi.org/10.1016/j.jacc.2013.12.027>
250. Nichol G, Strickland W, Shavell D, Maehara A, Ben-Yehuda O, Genereux P, et al. Prospective, multicenter, randomized, controlled pilot trial of peritoneal hypothermia in patients with ST-segment-elevation myocardial infarction. *Circ Cardiovasc Interv* 2015;**8**:e001965. <https://doi.org/10.1161/circinterventions.114.001965>
251. Noc M, Erlinge D, Neskovic AN, Kafedzic S, Merkely B, Zima E, et al. COOL AMI EU pilot trial: a multicentre, prospective, randomised controlled trial to assess cooling as an adjunctive therapy to percutaneous intervention in patients with acute myocardial infarction. *Eurointervention* 2017;**13**:e531–e539. <https://doi.org/10.4244/eij-d-17-00279>
252. Ibáñez B, Heusch G, Ovize M, Van de Werf F. Evolving therapies for myocardial ischemia/reperfusion injury. *J Am Coll Cardiol* 2015;**65**:1454–1471. <https://doi.org/10.1016/j.jacc.2015.02.032>
253. Hausenloy DJ, Botker HE, Engstrom T, Erlinge D, Heusch G, Ibanez B, et al. Targeting reperfusion injury in patients with ST-segment elevation myocardial infarction: trials and tribulations. *Eur Heart J* 2017;**38**:935–941. <https://doi.org/10.1093/eurheartj/ehw145>
254. García-Prieto J, Villena-Gutiérrez R, Gómez M, Bernardo E, Pun-García A, García-Lunar I, et al. Neutrophil stunning by metoprolol reduces infarct size. *Nat Commun* 2017;**8**:14780. <https://doi.org/10.1038/ncomms14780>
255. Ibanez B, Macaya C, Sánchez-Brunete V, Pizarro G, Fernández-Friera L, Mateos A, et al. Effect of early metoprolol on infarct size in ST-segment-elevation myocardial infarction patients undergoing primary percutaneous coronary intervention: the Effect of Metoprolol in Cardioprotection During an Acute Myocardial Infarction (METOCARD-CNIC) trial. *Circulation* 2013;**128**:1495–1503. <https://doi.org/10.1161/circulationaha.113.003653>
256. Pizarro G, Fernández-Friera L, Fuster V, Fernández-Jiménez R, García-Ruiz JM, García-Álvarez A, et al. Long-term benefit of early pre-reperfusion metoprolol administration in patients with acute myocardial infarction: results from the METOCARD-CNIC trial (Effect of Metoprolol in Cardioprotection During an Acute Myocardial Infarction). *J Am Coll Cardiol* 2014;**63**:2356–2362. <https://doi.org/10.1016/j.jacc.2014.03.014>
257. Roolink V, Ibáñez B, Ottavanger JP, Pizarro G, van Royen N, Mateos A, et al. Early intravenous beta-blockers in patients with ST-segment elevation myocardial infarction before primary percutaneous coronary intervention. *J Am Coll Cardiol* 2016;**67**:2705–2715. <https://doi.org/10.1016/j.jacc.2016.03.522>
258. García-Ruiz JM, Fernández-Jiménez R, García-Álvarez A, Pizarro G, Galán-Arriola C, Fernández-Friera L, et al. Impact of the timing of metoprolol administration during STEMI on infarct size and ventricular function. *J Am Coll Cardiol* 2016;**67**:2093–2104. <https://doi.org/10.1016/j.jacc.2016.02.050>
259. Lobo-Gonzalez M, Galán-Arriola C, Rossello X, González-Del-Hoyo M, Vilchez JP, Higuero-Verdejo MI, et al. Metoprolol blunts the time-dependent progression of infarct size. *Basic Res Cardiol* 2020;**115**:55. <https://doi.org/10.1007/s00395-020-0812-4>
260. Clemente-Moragón A, Gómez M, Villena-Gutiérrez R, Lalama DV, García-Prieto J, Martínez F, et al. Metoprolol exerts a non-class effect against ischaemia-reperfusion injury by abrogating exacerbated inflammation. *Eur Heart J* 2020;**41**:4425–4440. <https://doi.org/10.1093/eurheartj/ehaa733>
261. Belle L, Motreff P, Mangin L, Rangé G, Marcaggi X, Marie A, et al. Comparison of immediate with delayed stenting using the minimalist immediate mechanical intervention approach in acute ST-segment-elevation myocardial infarction: the MIMI study. *Circ Cardiovasc Interv* 2016;**9**:e003388. <https://doi.org/10.1161/circinterventions.115.003388>
262. Carrick D, Oldroyd KG, McEntegart M, Haig C, Petrie MC, Eteiba H, et al. A randomized trial of deferred stenting versus immediate stenting to prevent no- or slow-reflow in acute ST-segment elevation myocardial infarction (DEFER-STEMI). *J Am Coll Cardiol* 2014;**63**:2088–2098. <https://doi.org/10.1016/j.jacc.2014.02.530>
263. Kelbaek H, Høfsten DE, Køber L, Helqvist S, Kløvgård L, Holmvang L, et al. Deferred versus conventional stent implantation in patients with ST-segment elevation myocardial infarction (DANAMI 3-DEFER): an open-label, randomised controlled trial. *Lancet* 2016;**387**:2199–2206. [https://doi.org/10.1016/s0140-6736\(16\)30072-1](https://doi.org/10.1016/s0140-6736(16)30072-1)
264. Lønborg J, Engstrøm T, Ahtarovski KA, Nepper-Christensen L, Helqvist S, Vejstrup N, et al. Myocardial damage in patients with deferred stenting after STEMI: a DANAMI-3-DEFER substudy. *J Am Coll Cardiol* 2017;**69**:2794–2804. <https://doi.org/10.1016/j.jacc.2017.03.601>
265. Kim JS, Lee HJ, Woong Yu C, Kim YM, Hong SJ, Park JH, et al. INNOVATION Study (impact of immediate stent implantation versus deferred stent implantation on infarct size and microvascular perfusion in patients with ST-segment-elevation myocardial infarction). *Circ Cardiovasc Interv* 2016;**9**:e004101. <https://doi.org/10.1161/circinterventions.116.004101>
266. Qiao J, Pan L, Zhang B, Wang J, Zhao Y, Yang R, et al. Deferred versus immediate stenting in patients with ST-segment elevation myocardial infarction: a systematic review and meta-analysis. *J Am Heart Assoc* 2017;**6**:e004838. <https://doi.org/10.1161/jaha.116.004838>
267. Ali Z, Landmesser U, Karimi Galougahi K, Maehara A, Matsumura M, Shlofmitz RA, et al. Optical coherence tomography-guided coronary stent implantation compared to angiography: a multicentre randomised trial in PCI—design and rationale of ILUMIEN IV: OPTIMAL PCI. *Eurointervention* 2021;**16**:1092–1099. <https://doi.org/10.4244/eij-d-20-00501>
268. Pizzi C, Xhyheri B, Costa GM, Faustino M, Flacco ME, Gualano MR, et al. Nonobstructive versus obstructive coronary artery disease in acute coronary syndrome: a meta-analysis. *J Am Heart Assoc* 2016;**5**:e004185. <https://doi.org/10.1161/jaha.116.004185>
269. Safdar B, Spatz ES, Dreyer RP, Beltrame JF, Lichtman JH, Spertus JA, et al. Presentation, clinical profile, and prognosis of young patients with Myocardial Infarction with Nonobstructive Coronary Arteries (MINOCA): results from the VIRGO study. *J Am Heart Assoc* 2018;**7**:e009174. <https://doi.org/10.1161/jaha.118.009174>
270. Ciliberti G, Coiro S, Tritto I, Benedetti M, Guerra F, Del Pinto M, et al. Predictors of poor clinical outcomes in patients with acute myocardial infarction and non-obstructed coronary arteries (MINOCA). *Int J Cardiol* 2018;**267**:41–45. <https://doi.org/10.1016/j.ijcard.2018.03.092>
271. Kunadian V, Chieffo A, Camici PG, Berry C, Escaned J, Maas AHEM, et al. An EAPCI expert consensus document on ischaemia with non-obstructive coronary arteries in collaboration with European Society of Cardiology Working Group on coronary pathophysiology & microcirculation endorsed by coronary vasomotor disorders international study group. *Eur Heart J* 2020;**41**:3504–3520. <https://doi.org/10.1093/eurheartj/ehaa503>
272. Lindahl B, Baron T, Erlinge D, Hadziosmanovic N, Nordenskjöld A, Gard A, et al. Medical therapy for secondary prevention and long-term outcome in patients with myocardial infarction with nonobstructive coronary artery disease. *Circulation* 2017;**135**:1481–1489. <https://doi.org/10.1161/circulationaha.116.026336>

273. Grodzinsky A, Arnold SV, Gosch K, Spertus JA, Foody JM, Beltrame J, et al. Angina frequency after acute myocardial infarction in patients without obstructive coronary artery disease. *Eur Heart J Qual Care Clin Outcomes* 2015;**1**:92–99. <https://doi.org/10.1093/ehjqqc/qcv014>
274. Vijayalakshmi K, De Belder MA. Angiographic and physiologic assessment of coronary flow and myocardial perfusion in the cardiac catheterization laboratory. *Acute Card Care* 2008;**10**:69–78. <https://doi.org/10.1080/17482940701606905>
275. Kunadian V, Harrigan C, Zorkun C, Palmer AM, Ogando KJ, Biller LH, et al. Use of the TIMI frame count in the assessment of coronary artery blood flow and microvascular function over the past 15 years. *J Thromb Thrombolysis* 2009;**27**:316–328. <https://doi.org/10.1007/s11239-008-0220-3>
276. Adlam D, Alfonso F, Maas A, Vrints C. European Society of Cardiology, Acute Cardiovascular Care Association, SCAD study group: a position paper on spontaneous coronary artery dissection. *Eur Heart J* 2018;**39**:3353–3368. <https://doi.org/10.1093/eurheartj/ehy080>
277. Lee BK, Lim HS, Fearon WF, Yong AS, Yamada R, Tanaka S, et al. Invasive evaluation of patients with angina in the absence of obstructive coronary artery disease. *Circulation* 2015;**131**:1054–1060. <https://doi.org/10.1161/circulationaha.114.012636>
278. Ford TJ, Stanley B, Good R, Rocchiccioli P, McEntegart M, Watkins S, et al. Stratified medical therapy using invasive coronary function testing in angina: the CorMicA trial. *J Am Coll Cardiol* 2018;**72**:2841–2855. <https://doi.org/10.1016/j.jacc.2018.09.006>
279. Ford TJ, Ong P, Sechtem U, Beltrame J, Camici PG, Crea F, et al. Assessment of vascular dysfunction in patients without obstructive coronary artery disease: why, how, and when. *JACC Cardiovasc Interv* 2020;**13**:1847–1864. <https://doi.org/10.1016/j.jcin.2020.05.052>
280. Ciliberti G, Seshasai SRK, Ambrosio G, Kaski JC. Safety of intracoronary provocative testing for the diagnosis of coronary artery spasm. *Int J Cardiol* 2017;**244**:77–83. <https://doi.org/10.1016/j.ijcard.2017.05.109>
281. Montone RA, Niccoli G, Fracassi F, Russo M, Gurgoglione F, Cammà G, et al. Patients with acute myocardial infarction and non-obstructive coronary arteries: safety and prognostic relevance of invasive coronary provocative tests. *Eur Heart J* 2018;**39**:91–98. <https://doi.org/10.1093/eurheartj/ehx667>
282. Kubo T, Imanishi T, Takarada S, Kuroi A, Ueno S, Yamano T, et al. Assessment of culprit lesion morphology in acute myocardial infarction: ability of optical coherence tomography compared with intravascular ultrasound and coronary angiography. *J Am Coll Cardiol* 2007;**50**:933–939. <https://doi.org/10.1016/j.jacc.2007.04.082>
283. Sinclair H, Bourantas C, Bagnall A, Mintz GS, Kunadian V. OCT for the identification of vulnerable plaque in acute coronary syndrome. *JACC Cardiovasc Imaging* 2015;**8**:198–209. <https://doi.org/10.1016/j.jcmg.2014.12.005>
284. Johnson TW, Räber L, di Mario C, Bourantas C, Jia H, Mattesini A, et al. Clinical use of intracoronary imaging. Part 2: acute coronary syndromes, ambiguous coronary angiography findings, and guiding interventional decision-making: an expert consensus document of the European Association of Percutaneous Cardiovascular Interventions. *Eur Heart J* 2019;**40**:2566–2584. <https://doi.org/10.1093/eurheartj/ehz332>
285. Konstantinides SV, Torbicki A, Agnelli G, Danchin N, Fitzmaurice D, Galis N, et al. 2014 ESC guidelines on the diagnosis and management of acute pulmonary embolism. *Eur Heart J* 2014;**35**:3033–3080, 3069a–3069k. <https://doi.org/10.1093/eurheartj/ehu283>
286. Imazio M, Negro A, Belli R, Beqaraj F, Forno D, Giammaria M, et al. Frequency and prognostic significance of pericarditis following acute myocardial infarction treated by primary percutaneous coronary intervention. *Am J Cardiol* 2009;**103**:1525–1529. <https://doi.org/10.1016/j.amjcard.2009.01.366>
287. Imazio M, Hoi BD. Post-cardiac injury syndromes. An emerging cause of pericardial diseases. *Int J Cardiol* 2013;**168**:648–652. <https://doi.org/10.1016/j.ijcard.2012.09.052>
288. Rossello X, Wiegerinck RF, Alguersuari J, Bardaji A, Worner F, Sutil M, et al. New electrocardiographic criteria to differentiate acute pericarditis and myocardial infarction. *Am J Med* 2014;**127**:233–239. <https://doi.org/10.1016/j.amjmed.2013.11.006>
289. Adler Y, Charron P, Imazio M, Badano L, Barón-Esquivias G, Bogaert J, et al. 2015 ESC Guidelines for the diagnosis and management of pericardial diseases: the Task Force for the diagnosis and management of pericardial diseases of the European Society of Cardiology (ESC). Endorsed by: the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J* 2015;**36**:2921–2964. <https://doi.org/10.1093/eurheartj/ehv318>
290. Verma BR, Montane B, Chetrit M, Khayata M, Furqan MM, Ayoub C, et al. Pericarditis and post-cardiac injury syndrome as a sequelae of acute myocardial infarction. *Curr Cardiol Rep* 2020;**22**:127. <https://doi.org/10.1007/s11886-020-01371-5>
291. Imazio M, Lazaros G, Picardi E, Vasileiou P, Orlando F, Carraro M, et al. Incidence and prognostic significance of new onset atrial fibrillation/flutter in acute pericarditis. *Heart* 2015;**101**:1463–1467. <https://doi.org/10.1136/heartjnl-2014-307398>
292. Hansson EC, Shams Hakimi C, Åström-Olsson K, Hesse C, Wallén H, Dellborg M, et al. Effects of ex vivo platelet supplementation on platelet aggregability in blood samples from patients treated with acetylsalicylic acid, clopidogrel, or ticagrelor. *Br J Anaesth* 2014;**112**:570–575. <https://doi.org/10.1093/bja/aet339>
293. Zafar MU, Santos-Gallego C, Vorchheimer DA, Viles-Gonzalez JF, Elmariah S, Giannarelli C, et al. Platelet function normalization after a prasugrel loading-dose: time-dependent effect of platelet supplementation. *J Thromb Haemost* 2013;**11**:100–106. <https://doi.org/10.1111/jth.12058>
294. Buchanan A, Newton P, Pehrsson S, Inghardt T, Antonsson T, Svensson P, et al. Structural and functional characterization of a specific antidote for ticagrelor. *Blood* 2015;**125**:3484–3490. <https://doi.org/10.1182/blood-2015-01-622928>
295. Agno VV, Crowther M, Steidl L, Ultori C, Mera V, Dentali F, et al. Low dose oral vitamin K to reverse acenocoumarol-induced coagulopathy: a randomized controlled trial. *Thromb Haemost* 2002;**88**:48–51. <https://doi.org/10.1055/s-0037-1613152>
296. Crowther MA, Julian J, McCarty D, Douketis J, Kovacs M, Biagioni L, et al. Treatment of warfarin-associated coagulopathy with oral vitamin K: a randomised controlled trial. *Lancet* 2000;**356**:1551–1553. [https://doi.org/10.1016/s0140-6736\(00\)03125-1](https://doi.org/10.1016/s0140-6736(00)03125-1)
297. Ageno W, Gallus AS, Wittkowsky A, Crowther M, Hylek EM, Palareti G, et al. Oral anticoagulant therapy: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of chest physicians evidence-based clinical practice guidelines. *Chest* 2012;**141**:e44S–e88S. <https://doi.org/10.1378/chest.11-2292>
298. Pollack CVJ, Reilly PA, van Ryn J, Eikelboom JW, Glund S, Bernstein RA, et al. Idarucizumab for dabigatran reversal—full cohort analysis. *N Engl J Med* 2017;**377**:431–441. <https://doi.org/10.1056/NEJMoa1707278>
299. Tomaselli GF, Mahaffey KW, Cuker A, Dobesh PP, Doherty JU, Eikelboom JW, et al. 2017 ACC expert consensus decision pathway on management of bleeding in patients on oral anticoagulants: a report of the American College of Cardiology Task Force on expert consensus decision pathways. *J Am Coll Cardiol* 2017;**70**:3042–3067. <https://doi.org/10.1016/j.jacc.2017.09.1085>
300. Eerenberg ES, Kamphuisen PW, Sijpkens MK, Meijers JC, Buller HR, Levi M, et al. Reversal of rivaroxaban and dabigatran by prothrombin complex concentrate: a randomized, placebo-controlled, crossover study in healthy subjects. *Circulation* 2011;**124**:1573–1579. <https://doi.org/10.1161/circulationaha.111.029017>
301. Connolly SJ, Crowther M, Eikelboom JW, Gibson CM, Curnutte JT, Lawrence JH, et al. Full study report of andexanet alfa for bleeding associated with factor Xa inhibitors. *N Engl J Med* 2019;**380**:1326–1335. <https://doi.org/10.1056/NEJMoa1814051>
302. Connolly SJ, Milling TJ Jr, Eikelboom JW, Gibson CM, Curnutte JT, Gold A, et al. Andexanet alfa for acute major bleeding associated with factor Xa inhibitors. *N Engl J Med* 2016;**375**:1131–1141. <https://doi.org/10.1056/NEJMoa1607887>
303. Ndrepepa G, Guerra E, Schulz S, Fusaro M, Cassese S, Kastrati A, et al. Weight of the bleeding impact on early and late mortality after percutaneous coronary intervention. *J Thromb Thrombolysis* 2015;**39**:35–42. <https://doi.org/10.1007/s11239-014-1084-3>
304. Leibundgut G, Pache J, Schulz S, Berger PB, Ferenc M, Gick M, et al. Collagen plug vascular closure devices and reduced risk of bleeding with bivalirudin versus heparin plus abciximab in patients undergoing percutaneous coronary intervention for non ST-segment elevation myocardial infarction. *J Interv Cardiol* 2013;**26**:623–629. <https://doi.org/10.1111/joi.12067>
305. Ndrepepa G, Neumann FJ, Richardt G, Schulz S, Tölg R, Stoyanov KM, et al. Prognostic value of access and non-access sites bleeding after percutaneous coronary intervention. *Circ Cardiovasc Interv* 2013;**6**:354–361. <https://doi.org/10.1161/circinterventions.113.000433>
306. Marso SP, Amin AP, House JA, Kennedy KF, Spertus JA, Rao SV, et al. Association between use of bleeding avoidance strategies and risk of periprocedural bleeding among patients undergoing percutaneous coronary intervention. *JAMA* 2010;**303**:2156–2164. <https://doi.org/10.1001/jama.2010.708>
307. Valgimigli M, Gagnor A, Calabró P, Frigoli E, Leonardi S, Zaro T, et al. Radial versus femoral access in patients with acute coronary syndromes undergoing invasive management: a randomised multicentre trial. *Lancet* 2015;**385**:2465–2476. [https://doi.org/10.1016/s0140-6736\(15\)60292-6](https://doi.org/10.1016/s0140-6736(15)60292-6)
308. Schulz-Schüpke S, Helde S, Gewalt S, Ibrahim T, Linhardt M, Haas K, et al. Comparison of vascular closure devices vs manual compression after femoral artery puncture: the ISAR-CLOSURE randomized clinical trial. *JAMA* 2014;**312**:1981–1987. <https://doi.org/10.1001/jama.2014.15305>
309. Shi J, Ji H, Ren F, Wang G, Xu M, Xue Y, et al. Protective effects of tranexamic acid on clopidogrel before coronary artery bypass grafting: a multicenter randomized trial. *JAMA Surg* 2013;**148**:538–547. <https://doi.org/10.1001/jamasurg.2013.1560>
310. Paone G, Likosky DS, Brewer R, Theurer PF, Bell GF, Cogan CM, et al. Transfusion of 1 and 2 units of red blood cells is associated with increased morbidity and mortality. *Ann Thorac Surg* 2014;**97**:87–93; discussion 93–84. <https://doi.org/10.1016/j.athoracsurg.2013.07.020>
311. Mehta RH, Sheng S, O'Brien SM, Grover FL, Gammie JS, Ferguson TB, et al. Reoperation for bleeding in patients undergoing coronary artery bypass surgery: incidence, risk factors, time trends, and outcomes. *Circ Cardiovasc Qual Outcomes* 2009;**2**:583–590. <https://doi.org/10.1161/circoutcomes.109.858811>
312. Hardy JF, Bélice S, Van der Linden P. Efficacy and safety of activated recombinant factor VII in cardiac surgical patients. *Curr Opin Anaesthesiol* 2009;**22**:95–99. <https://doi.org/10.1097/ACO.0b013e32831a40a3>
313. Rao SV, Jollis JG, Harrington RA, Granger CB, Newby LK, Armstrong PW, et al. Relationship of blood transfusion and clinical outcomes in patients with acute coronary syndromes. *JAMA* 2004;**292**:1555–1562. <https://doi.org/10.1001/jama.292.13.1555>

314. Sherwood MW, Wang Y, Curtis JP, Peterson ED, Rao SV. Patterns and outcomes of red blood cell transfusion in patients undergoing percutaneous coronary intervention. *JAMA* 2014;**311**:836–843. <https://doi.org/10.1001/jama.2014.980>
315. Nikolsky E, Mehran R, Sadeghi HM, Grines CL, Cox DA, Garcia E, et al. Prognostic impact of blood transfusion after primary angioplasty for acute myocardial infarction: analysis from the CADILLAC (Controlled Abciximab and Device Investigation to Lower Late Angioplasty Complications) trial. *JACC Cardiovasc Interv* 2009;**2**:624–632. <https://doi.org/10.1016/j.jcin.2009.05.004>
316. Kansagara D, Dyer E, Englander H, Fu R, Freeman M, Kagen D, et al. Treatment of anemia in patients with heart disease: a systematic review. *Ann Intern Med* 2013;**159**:746–757. <https://doi.org/10.7326/0003-4819-159-11-201312030-00007>
317. Carson JL, Carless PA, Hebert PC. Transfusion thresholds and other strategies for guiding allogeneic red blood cell transfusion. *Cochrane Database Syst Rev* 2012;**4**:CD002042. <https://doi.org/10.1002/14651858.CD002042.pub3>
318. Chatterjee S, Wetterslev J, Sharma A, Lichstein E, Mukherjee D. Association of blood transfusion with increased mortality in myocardial infarction: a meta-analysis and diversity-adjusted study sequential analysis. *JAMA Intern Med* 2013;**173**:132–139. <https://doi.org/10.1001/2013.jamainternmed.1001>
319. Cooper HA, Rao SV, Greenberg MD, Rumsey MP, McKenzie M, Alcorn KW, et al. Conservative versus liberal red cell transfusion in acute myocardial infarction (the CRIT Randomized Pilot Study). *Am J Cardiol* 2011;**108**:1108–1111. <https://doi.org/10.1016/j.amjcard.2011.06.014>
320. Alexander KP, Chen AY, Wang TY, Rao SV, Newby LK, LaPointe NMA, et al. Transfusion practice and outcomes in non-ST-segment elevation acute coronary syndromes. *Am Heart J* 2008;**155**:1047–1053. <https://doi.org/10.1016/j.ahj.2008.01.009>
321. Holst LB, Petersen MW, Haase N, Perner A, Wetterslev J. Restrictive versus liberal transfusion strategy for red blood cell transfusion: systematic review of randomised trials with meta-analysis and trial sequential analysis. *BMJ* 2015;**350**:h1354. <https://doi.org/10.1136/bmj.h1354>
322. Murphy GJ, Pike K, Rogers CA, Wordsworth S, Stokes EA, Angelini GD, et al. Liberal or restrictive transfusion after cardiac surgery. *N Engl J Med* 2015;**372**:997–1008. <https://doi.org/10.1056/NEJMoa1403612>
323. Wessler JD, Giugliano RP. Risk of thrombocytopenia with glycoprotein IIb/IIIa inhibitors across drugs and patient populations: a meta-analysis of 29 large placebo-controlled randomized trials. *Eur Heart J Cardiovasc Pharmacother* 2015;**1**:97–106. <https://doi.org/10.1093/ehjcvp/pvu008>
324. Dasgupta H, Blankenship JC, Wood GC, Frey CM, Demko SL, Menapace FJ, et al. Thrombocytopenia complicating treatment with intravenous glycoprotein IIb/IIIa receptor inhibitors: a pooled analysis. *Am Heart J* 2000;**140**:206–211. <https://doi.org/10.1016/j.mhj.2000.107554>
325. Bakchoul T, Marini L. Drug-associated thrombocytopenia. *Hematology Am Soc Hematol Educ Program* 2018;**2018**:576–583. <https://doi.org/10.1182/asheducation-2018.1.576>
326. Arnold DM, Nazi I, Warkentin TE, Smith JW, Tóth LJ, George JN, et al. Approach to the diagnosis and management of drug-induced immune thrombocytopenia. *Transfus Med Rev* 2013;**27**:137–145. <https://doi.org/10.1016/j.tmr.2013.05.005>
327. Greinacher A, Selleng S. How I evaluate and treat thrombocytopenia in the intensive care unit patient. *Blood* 2016;**128**:3032–3042. <https://doi.org/10.1182/blood-2016-09-693655>
328. Warkentin TE, Levine MN, Hirsh J, Horsewood P, Roberts RS, Gent M, et al. Heparin-induced thrombocytopenia in patients treated with low-molecular-weight heparin or unfractionated heparin. *N Engl J Med* 1995;**332**:1330–1336. <https://doi.org/10.1056/nejm199505183322003>
329. Kelton JG, Arnold DM, Bates SM. Nonheparin anticoagulants for heparin-induced thrombocytopenia. *N Engl J Med* 2013;**368**:737–744. <https://doi.org/10.1056/NEJMct1206642>
330. Warkentin TE, Kelton JG. Temporal aspects of heparin-induced thrombocytopenia. *N Engl J Med* 2001;**344**:1286–1292. <https://doi.org/10.1056/nejm200104263441704>
331. Schindewolf M, Steindl J, Beyer-Westendorf J, Schellong S, Dohmen PM, Brachmann J, et al. Use of fondaparinux off-label or approved anticoagulants for management of heparin-induced thrombocytopenia. *J Am Coll Cardiol* 2017;**70**:2636–2648. <https://doi.org/10.1016/j.jacc.2017.09.1099>
332. Sinclair H, Kunadian V. Coronary revascularisation in older patients with non-ST elevation acute coronary syndromes. *Heart* 2016;**102**:416–424. <https://doi.org/10.1136/heartjnl-2015-307859>
333. Richter D, Guasti L, Walker D, Lambrinou E, Lionis C, Abreu A, et al. Frailty in cardiology: definition, assessment and clinical implications for general cardiology. A consensus document of the Council for Cardiology Practice (CCP), Association for Acute Cardio Vascular Care (ACVC), Association of Cardiovascular Nursing and Allied Professions (ACNAP), European Association of Preventive Cardiology (EAPC), European Heart Rhythm Association (EHRA), Council on Valvular Heart Diseases (VHD), Council on Hypertension (CHT), Council of Cardio-Oncology (CCO), Working Group (WG) Aorta and Peripheral Vascular Diseases, WG e-Cardiology, WG Thrombosis, of the European Society of Cardiology, European Primary Care Cardiology Society (EPCCS). *Eur J Prev Cardiol* 2022;**29**:216–227. <https://doi.org/10.1093/eurjpc/zwaa167>
334. Lopes RD, White JA, Tricoci P, White HD, Armstrong PW, Braunwald E, et al. Age, treatment, and outcomes in high-risk non-ST-segment elevation acute coronary syndrome patients: insights from the EARLY ACS trial. *Int J Cardiol* 2013;**167**:2580–2587. <https://doi.org/10.1016/j.ijcard.2012.06.053>
335. Brieger D, Eagle KA, Goodman SG, Steg PG, Budaj A, White K, et al. Acute coronary syndromes without chest pain, an underdiagnosed and undertreated high-risk group: insights from the Global Registry of Acute Coronary Events. *Chest* 2004;**126**:461–469. <https://doi.org/10.1378/chest.126.2.461>
336. Goldberg RJ, Steg PG, Sadiq I, Granger CB, Jackson EA, Budaj A, et al. Extent of, and factors associated with, delay to hospital presentation in patients with acute coronary disease (the GRACE registry). *Am J Cardiol* 2002;**89**:791–796. [https://doi.org/10.1016/s0002-9149\(02\)02186-0](https://doi.org/10.1016/s0002-9149(02)02186-0)
337. Rosengren A, Wallentin L, Simoons-Smit AM, Gitt AK, Behar S, Battler A, et al. Age, clinical presentation, and outcome of acute coronary syndromes in the Euroheart acute coronary syndrome survey. *Eur Heart J* 2006;**27**:789–795. <https://doi.org/10.1093/eurheartj/ehi774>
338. Reiter M, Twerenbold R, Reichlin T, Haaf P, Peter F, Meissner J, et al. Early diagnosis of acute myocardial infarction in the elderly using more sensitive cardiac troponin assays. *Eur Heart J* 2011;**32**:1379–1389. <https://doi.org/10.1093/eurheartj/ehr033>
339. Zhang F, Mohamed MO, Ensor J, Peat G, Mamas MA. Temporal trends in comorbidity burden and impact on prognosis in patients with acute coronary syndrome using the elixhauser comorbidity index score. *Am J Cardiol* 2020;**125**:1603–1611. <https://doi.org/10.1016/j.amjcard.2020.02.044>
340. Borovac JA, Mohamed MO, Kontopantelis E, Alkhoul M, Alraies MC, Cheng RK, et al. Frailty among patients with acute ST-elevation myocardial infarction in the United States: the impact of the primary percutaneous coronary intervention on in-hospital outcomes. *J Invasive Cardiol* 2022;**34**:E55–E64.
341. Cholesterol Treatment Trialists' Collaboration. Efficacy and safety of statin therapy in older people: a meta-analysis of individual participant data from 28 randomised controlled trials. *Lancet* 2019;**393**:407–415. [https://doi.org/10.1016/s0140-6736\(18\)31942-1](https://doi.org/10.1016/s0140-6736(18)31942-1)
342. Bach RG, Cannon CP, Giugliano RP, White JA, Lokhnygina Y, Bohula EA, et al. Effect of simvastatin-ezetimibe compared with simvastatin monotherapy after acute coronary syndrome among patients 75 years or older: a secondary analysis of a randomized clinical trial. *JAMA Cardiol* 2019;**4**:846–854. <https://doi.org/10.1001/jamacardio.2019.2306>
343. Alexander KP, Newby LK, Cannon CP, Armstrong PW, Gibler WB, Rich MW, et al. Acute coronary care in the elderly, part I: non-ST-segment-elevation acute coronary syndromes: a scientific statement for healthcare professionals from the American Heart Association Council on Clinical Cardiology: in collaboration with the Society of Geriatric Cardiology. *Circulation* 2007;**115**:2549–2569. <https://doi.org/10.1161/CIRCULATIONAHA.107.182615>
344. Alexander KP, Chen AY, Roe MT, Newby LK, Gibson CM, Allen-LaPointe NM, et al. Excess dosing of antiplatelet and antithrombin agents in the treatment of non-ST-segment elevation acute coronary syndromes. *JAMA* 2005;**294**:3108–3116. <https://doi.org/10.1001/jama.294.24.3108>
345. Valgimigli M, Frigoli E, Heg D, Tijssen J, Juni P, Vranckx P, et al. Dual antiplatelet therapy after PCI in patients at high bleeding risk. *N Engl J Med* 2021;**385**:1643–1655. <https://doi.org/10.1056/NEJMoa2108749>
346. Roe MT, Armstrong PW, Fox KA, White HD, Prabhakaran D, Goodman SG, et al. Prasugrel versus clopidogrel for acute coronary syndromes without revascularization. *N Engl J Med* 2012;**367**:1297–1309. <https://doi.org/10.1056/NEJMoa1205512>
347. Savonitto S, Ferri LA, Piatti L, Grosseto D, Piovaccari G, Morici N, et al. Comparison of reduced-dose prasugrel and standard-dose clopidogrel in elderly patients with acute coronary syndromes undergoing early percutaneous revascularization. *Circulation* 2018;**137**:2435–2445. <https://doi.org/10.1161/circulationaha.117.032180>
348. Mehran R, Baber U, Sharma SK, Cohen DJ, Angiolillo DJ, Briguori C, et al. Ticagrelor with or without aspirin in high-risk patients after PCI. *N Engl J Med* 2019;**381**:2032–2042. <https://doi.org/10.1056/NEJMoa1908419>
349. Angiolillo DJ, Cao D, Baber U, Sartori S, Zhang Z, Dangas G, et al. Impact of age on the safety and efficacy of ticagrelor monotherapy in patients undergoing PCI. *JACC Cardiovasc Interv* 2021;**14**:1434–1446. <https://doi.org/10.1016/j.jcin.2021.04.043>
350. Escaned J, Cao D, Baber U, Nicolas J, Sartori S, Zhang Z, et al. Ticagrelor monotherapy in patients at high bleeding risk undergoing percutaneous coronary intervention: TWILIGHT-HBR. *Eur Heart J* 2021;**42**:4624–4634. <https://doi.org/10.1093/eurheartj/ehab702>
351. Gargiulo G, Goette A, Tijssen J, Eckardt L, Lewalter T, Vranckx P, et al. Safety and efficacy outcomes of double vs. triple antithrombotic therapy in patients with atrial fibrillation following percutaneous coronary intervention: a systematic review and meta-analysis of non-vitamin K antagonist oral anticoagulant-based randomized clinical trials. *Eur Heart J* 2019;**40**:3757–3767. <https://doi.org/10.1093/eurheartj/ehz732>
352. Nanda S, Nelson-Piercy C, Mackillop L. Cardiac disease in pregnancy. *Clin Med (Lond)* 2012;**12**:553–560. <https://doi.org/10.7861/clinmedicine.12-6-553>
353. Bush N, Nelson-Piercy C, Spark P, Kurinczuk JJ, Brocklehurst P, Knight M, et al. Myocardial infarction in pregnancy and postpartum in the UK. *Eur J Prev Cardiol* 2013;**20**:12–20. <https://doi.org/10.1177/1741826711432117>

354. Berg CJ, Callaghan WM, Syverson C, Henderson Z. Pregnancy-related mortality in the United States, 1998 to 2005. *Obstet Gynecol* 2010;**116**:1302–1309. <https://doi.org/10.1097/AOG.0b013e3181f8fb11>
355. James AH, Jamison MG, Biswas MS, Brancazio LR, Swamy GK, Myers ER, et al. Acute myocardial infarction in pregnancy: a United States population-based study. *Circulation* 2006;**113**:1564–1571. <https://doi.org/10.1161/circulationaha.105.576751>
356. Elkayam U, Jalnapurkar S, Barakkat MN, Khatri N, Kealey AJ, Mehra A, et al. Pregnancy-associated acute myocardial infarction: a review of contemporary experience in 150 cases between 2006 and 2011. *Circulation* 2014;**129**:1695–1702. <https://doi.org/10.1161/circulationaha.113.002054>
357. Ahmed R, Dunford J, Mehran R, Robson S, Kunadian V. Pre-eclampsia and future cardiovascular risk among women: a review. *J Am Coll Cardiol* 2014;**63**:1815–1822. <https://doi.org/10.1016/j.jacc.2014.02.529>
358. Tweet MS, Hayes SN, Pitta SR, Simari RD, Lerman A, Lennon RJ, et al. Clinical features, management, and prognosis of spontaneous coronary artery dissection. *Circulation* 2012;**126**:579–588. <https://doi.org/10.1161/circulationaha.112.105718>
359. Saw J, Humphries K, Aymong E, Sedlak T, Prakash R, Starovoytov A, et al. Spontaneous coronary artery dissection: clinical outcomes and risk of recurrence. *J Am Coll Cardiol* 2017;**70**:1148–1158. <https://doi.org/10.1016/j.jacc.2017.06.053>
360. Alfonso F, Paulo M, Lennie V, Dutary J, Bernardo E, Jiménez-Quevedo P, et al. Spontaneous coronary artery dissection: long-term follow-up of a large series of patients prospectively managed with a 'conservative' therapeutic strategy. *JACC Cardiovasc Interv* 2012;**5**:1062–1070. <https://doi.org/10.1016/j.jcin.2012.06.014>
361. Beltrame JF, Crea F, Kaski JC, Ogawa H, Ong P, Sehtem U, et al. International standardization of diagnostic criteria for vasospastic angina. *Eur Heart J* 2017;**38**:2565–2568. <https://doi.org/10.1093/eurheartj/ehv351>
362. Regitz-Zagrosek V, Roos-Hesselink JW, Bauersachs J, Blomström-Lundqvist C, Cifková R, De Bonis M, et al. 2018 ESC Guidelines for the management of cardiovascular diseases during pregnancy. *Eur Heart J* 2018;**39**:3165–3241. <https://doi.org/10.1093/eurheartj/ehy340>
363. Bortnick AE, Shahid M, Shitole SG, Park M, Broder A, Rodriguez CJ, et al. Outcomes of ST-elevation myocardial infarction by age and sex in a low-income urban community: the Montefiore STEMI registry. *Clin Cardiol* 2020;**43**:1100–1109. <https://doi.org/10.1002/clc.23412>
364. Kadri AN, Khodor S, Ali A, Nusairat L, Mahmood A, Nahhas G, et al. National trends of tobacco, alcohol, and drug use in patients admitted with acute myocardial infarction. *Cardiovasc Revasc Med* 2021;**26**:26–31. <https://doi.org/10.1016/j.carrev.2020.10.021>
365. Desai R, Fong HK, Shah K, Kaur VP, Savani S, Gangani K, et al. Rising trends in hospitalizations for cardiovascular events among young Cannabis users (18–39 years) without other substance abuse. *Medicina (Kaunas)* 2019;**55**:438. <https://doi.org/10.3390/medicina55080438>
366. Bonsignore A, Barranco R, Morando A, Fraternali Orcioni G, Ventura F. MDMA induced cardio-toxicity and pathological myocardial effects: a systematic review of experimental data and autopsy findings. *Cardiovasc Toxicol* 2019;**19**:493–499. <https://doi.org/10.1007/s12012-019-09526-9>
367. Carrillo X, Vilalta V, Cediell G, Fernandez-Nofrerias E, Rodriguez-Leor O, Mauri J, et al. Trends in prevalence and outcomes of acute coronary syndrome associated with cocaine consumption: the RUTI-cocaine study. *Int J Cardiol* 2019;**283**:23–27. <https://doi.org/10.1016/j.ijcard.2018.12.026>
368. Bharadwaj A, Potts J, Mohamed MO, Parwani P, Swamy P, Lopez-Mattei JC, et al. Acute myocardial infarction treatments and outcomes in 6.5 million patients with a current or historical diagnosis of cancer in the USA. *Eur Heart J* 2020;**41**:2183–2193. <https://doi.org/10.1093/eurheartj/ehz851>
369. Gevaert SA, Halvorsen S, Sinnaeve PR, Sambola A, Gulati G, Lancellotti P, et al. Evaluation and management of cancer patients presenting with acute cardiovascular disease: a consensus document of the Acute CardioVascular Care (ACVC) association and the ESC Council of Cardio-Oncology-Part 1: acute coronary syndromes and acute pericardial diseases. *Eur Heart J Acute Cardiovasc Care* 2021;**10**:947–959. <https://doi.org/10.1093/ehjacc/zuab056>
370. Iliescu CA, Grines CL, Herrmann J, Yang EH, Cilingiroglu M, Charitakis K, et al. SCAI Expert consensus statement: evaluation, management, and special considerations of cardio-oncology patients in the cardiac catheterization laboratory (endorsed by the Cardiological Society of India, and Sociedad Latino Americana de Cardiología Intervencionista). *Catheter Cardiovasc Interv* 2016;**87**:E202–E223. <https://doi.org/10.1002/ccd.26379>
371. Brosius FC III, Waller BF, Roberts WC. Radiation heart disease: analysis of 16 young (aged 15 to 33 years) necropsy patients who received over 3,500 rads to the heart. *Am J Med* 1981;**70**:519–530. [https://doi.org/10.1016/0002-9343\(81\)90574-x](https://doi.org/10.1016/0002-9343(81)90574-x)
372. Veinot JP, Edwards WD. Pathology of radiation-induced heart disease: a surgical and autopsy study of 27 cases. *Hum Pathol* 1996;**27**:766–773. [https://doi.org/10.1016/s0046-8177\(96\)90447-5](https://doi.org/10.1016/s0046-8177(96)90447-5)
373. Mohamed MO, Van Spall HGC, Kontopantelis E, Alkhouli M, Barac A, Elgendy IY, et al. Effect of primary percutaneous coronary intervention on in-hospital outcomes among active cancer patients presenting with ST-elevation myocardial infarction: a propensity score matching analysis. *Eur Heart J Acute Cardiovasc Care* 2021;**10**:829–839. <https://doi.org/10.1093/ehjacc/zuab032>
374. Guddati AK, Joy PS, Kumar G. Analysis of outcomes of percutaneous coronary intervention in metastatic cancer patients with acute coronary syndrome over a 10-year period. *J Cancer Res Clin Oncol* 2016;**142**:471–479. <https://doi.org/10.1007/s00432-015-2056-5>
375. Yusuf SW, Daraban N, Abbasi N, Lei X, Durand J-B, Daher IN, et al. Treatment and outcomes of acute coronary syndrome in the cancer population. *Clin Cardiol* 2012;**35**:443–450. <https://doi.org/10.1002/clc.22007>
376. Sarkiss MG, Yusuf SW, Warneke CL, Warneke CL, Botz G, Lakkis N, et al. Impact of aspirin therapy in cancer patients with thrombocytopenia and acute coronary syndromes. *Cancer* 2007;**109**:621–627. <https://doi.org/10.1002/cncr.22434>
377. Iannaccone M, D'Ascenzo F, Vadalà P, Wilton SB, Ntoussas P, Colombo F, et al. Prevalence and outcome of patients with cancer and acute coronary syndrome undergoing percutaneous coronary intervention: a BleeMACS substudy. *Eur Heart J Acute Cardiovasc Care* 2018;**7**:631–638. <https://doi.org/10.1177/2048872617706501>
378. Smith CM, Lim Choi Keung SN, Khan MO, Arvanitis TN, Fothergill R, Hartley-Sharpe C, et al. Barriers and facilitators to public access defibrillation in out-of-hospital cardiac arrest: a systematic review. *Eur Heart J Qual Care Clin Outcomes* 2017;**3**:264–273. <https://doi.org/10.1093/ehjccco/qcx023>
379. Ferrante G, Fazzari F, Cozzi O, Maurina M, Bragato R, D'Orazio F, et al. Risk factors for myocardial injury and death in patients with COVID-19: insights from a cohort study with chest computed tomography. *Cardiovasc Res* 2020;**116**:2239–2246. <https://doi.org/10.1093/cvr/cvaa193>
380. Madjid M, Safavi-Naeini P, Solomon SD, Vardeny O. Potential effects of coronaviruses on the cardiovascular system: a review. *JAMA Cardiol* 2020;**5**:831–840. <https://doi.org/10.1001/jamacardio.2020.1286>
381. Choudry FA, Hamshere SM, Rathod KS, Akhtar MM, Archbold RA, Guttmann OP, et al. High thrombus burden in patients with COVID-19 presenting with ST-segment elevation myocardial infarction. *J Am Coll Cardiol* 2020;**76**:1168–1176. <https://doi.org/10.1016/j.jacc.2020.07.022>
382. Solano-López J, Zamorano JL, Sanz AP, Amat-Santos I, Sarnago F, Gutiérrez Ibañes E, et al. Risk factors for in-hospital mortality in patients with acute myocardial infarction during the COVID-19 outbreak. *Rev Esp Cardiol* 2020;**73**:985–993. <https://doi.org/10.1016/j.recresp.2020.07.023>
383. Rodríguez-Leor O, Cid-Álvarez B, Pérez de Prado A, Rossello X, Ojeda S, Serrador A, et al. Impact of COVID-19 on ST-segment elevation myocardial infarction care. The Spanish experience. *Rev Esp Cardiol (Engl Ed)* 2020;**73**:994–1002. <https://doi.org/10.1016/j.rec.2020.08.002>
384. Rodríguez-Leor O, Cid-Álvarez AB, Pérez de Prado A, Rossello X, Ojeda S, Serrador A, et al. In-hospital outcomes of COVID-19 ST-elevation myocardial infarction patients. *Eurointervention* 2021;**16**:1426–1433. <https://doi.org/10.4244/eij-d-20-00935>
385. Kite TA, Ludman PF, Gale CP, Wu J, Caixeta A, Mansourati J, et al. International prospective registry of acute coronary syndromes in patients with COVID-19. *J Am Coll Cardiol* 2021;**77**:2466–2476. <https://doi.org/10.1016/j.jacc.2021.03.309>
386. Hamadeh A, Aldujeli A, Briedis K, Tecson KM, Sanz-Sánchez J, Al dujeli M, et al. Characteristics and outcomes in patients presenting with COVID-19 and ST-segment elevation myocardial infarction. *Am J Cardiol* 2020;**131**:1–6. <https://doi.org/10.1016/j.amjcard.2020.06.063>
387. Sun W, Gholizadeh L, Perry L, Kang K, Heydari M. Factors associated with return to work following myocardial infarction: a systematic review of observational studies. *J Clin Nurs* 2021;**30**:323–340. <https://doi.org/10.1111/jocn.15562>
388. Ambrosetti M, Abreu A, Corrà U, Davos CH, Hansen D, Frederix I, et al. Secondary prevention through comprehensive cardiovascular rehabilitation: from knowledge to implementation. 2020 update. A position paper from the secondary prevention and rehabilitation section of the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2021;**28**:460–495. <https://doi.org/10.1177/2047487320913379>
389. Smith D, Toff W, Joy M, Dowdall N, Johnston R, Clark L, et al. Fitness to fly for passengers with cardiovascular disease. *Heart* 2010;**96**:ii1–ii6. <https://doi.org/10.1136/hrt.2010.203091>
390. Liu C, Chen R, Sera F, Cabrera AM, Guo Y, Tong S, et al. Ambient particulate air pollution and daily mortality in 652 cities. *N Engl J Med* 2019;**381**:705–715. <https://doi.org/10.1056/NEJMoa1817364>
391. Newman JD, Bhatt DL, Rajagopalan S, Balmes JR, Brauer M, Breyse PN, et al. Cardiopulmonary impact of particulate air pollution in high-risk populations: JACC state-of-the-art review. *J Am Coll Cardiol* 2020;**76**:2878–2894. <https://doi.org/10.1016/j.jacc.2020.10.020>
392. Claeys MJ, Rajagopalan S, Nawrot TS, Brook RD. Climate and environmental triggers of acute myocardial infarction. *Eur Heart J* 2017;**38**:955–960. <https://doi.org/10.1093/eurheartj/ehw151>
393. Giorgini P, Rubenfire M, Bard RL, Jackson EA, Ferri C, Brook RD. Air pollution and exercise: a review of the cardiovascular implications for health care professionals. *J Cardiopulm Rehabil Prev* 2016;**36**:84–95. <https://doi.org/10.1097/hcr.0000000000000139>
394. Chen R, Zhao A, Chen H, Zhao Z, Cai J, Wang C, et al. Cardiopulmonary benefits of reducing indoor particles of outdoor origin: a randomized, double-blind crossover trial of air purifiers. *J Am Coll Cardiol* 2015;**65**:2279–2287. <https://doi.org/10.1016/j.jacc.2015.03.553>

395. Allen RW, Barn P. Individual- and household-level interventions to reduce air pollution exposures and health risks: a review of the recent literature. *Curr Environ Health Rep* 2020;**7**:424–440. <https://doi.org/10.1007/s40572-020-00296-z>
396. Münzel T, Schmidt FP, Steven S, Herzog J, Daiber A, Sørensen M. Environmental noise and the cardiovascular system. *J Am Coll Cardiol* 2018;**71**:688–697. <https://doi.org/10.1016/j.jacc.2017.12.015>
397. Fulcher J, O'Connell R, Voysey M, Emberson J, Blackwell L, Mihaylova B, et al. Efficacy and safety of LDL-lowering therapy among men and women: meta-analysis of individual data from 174,000 participants in 27 randomised trials. *Lancet* 2015;**385**:1397–1405. [https://doi.org/10.1016/s0140-6736\(14\)61368-4](https://doi.org/10.1016/s0140-6736(14)61368-4)
398. Collins R, Reith C, Emberson J, Armitage J, Baigent C, Blackwell L, et al. Interpretation of the evidence for the efficacy and safety of statin therapy. *Lancet* 2016;**388**:2532–2561. [https://doi.org/10.1016/s0140-6736\(16\)31357-5](https://doi.org/10.1016/s0140-6736(16)31357-5)
399. Navarese EP, Kowalewski M, Andreotti F, van Wely M, Camaro C, Kolodziejczak M, et al. Meta-analysis of time-related benefits of statin therapy in patients with acute coronary syndrome undergoing percutaneous coronary intervention. *Am J Cardiol* 2014;**113**:1753–1764. <https://doi.org/10.1016/j.amjcard.2014.02.034>
400. Mach F, Baigent C, Catapano AL, Koskinas KC, Casula M, Badimon L, et al. 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J* 2020;**41**:111–188. <https://doi.org/10.1093/eurheartj/ehz455>
401. Cannon CP, Blazing MA, Giugliano RP, McCagg A, White JA, Theroux P, et al. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med* 2015;**372**:2387–2397. <https://doi.org/10.1056/NEJMoa1410489>
402. O'Donoghue ML, Fazio S, Giugliano RP, Stroes ESG, Kanevsky E, Gouni-Berthold I, et al. Lipoprotein(a), PCSK9 inhibition, and cardiovascular risk. *Circulation* 2019;**139**:1483–1492. <https://doi.org/10.1161/circulationaha.118.037184>
403. Sabatine MS, Giugliano RP, Keech AC, Honarpour N, Wiviott SD, Murphy SA, et al. Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med* 2017;**376**:1713–1722. <https://doi.org/10.1056/NEJMoa1615664>
404. Schwartz GG, Steg PG, Szarek M, Bhatt DL, Bittner VA, Diaz R, et al. Alirocumab and cardiovascular outcomes after acute coronary syndrome. *N Engl J Med* 2018;**379**:2097–2107. <https://doi.org/10.1056/NEJMoa1801174>
405. Giugliano RP, Mach F, Zavitz K, Kurtz C, Im K, Kanevsky E, et al. Cognitive function in a randomized trial of evolocumab. *N Engl J Med* 2017;**377**:633–643. <https://doi.org/10.1056/NEJMoa1701131>
406. Ray KK, Colhoun HM, Szarek M, Baccara-Dinet M, Bhatt DL, Bittner VA, et al. Effects of alirocumab on cardiovascular and metabolic outcomes after acute coronary syndrome in patients with or without diabetes: a prespecified analysis of the ODYSSEY OUTCOMES randomised controlled trial. *Lancet Diabetes Endocrinol* 2019;**7**:618–628. [https://doi.org/10.1016/s2213-8587\(19\)30158-5](https://doi.org/10.1016/s2213-8587(19)30158-5)
407. Sabatine MS, De Ferrari GM, Giugliano RP, Huber K, Lewis BS, Ferreira J, et al. Clinical benefit of evolocumab by severity and extent of coronary artery disease: analysis from FOURIER. *Circulation* 2018;**138**:756–766. <https://doi.org/10.1161/circulationaha.118.034309>
408. Bonaca MP, Nault P, Giugliano RP, Keech AC, Pineda AL, Kanevsky E, et al. Low-density lipoprotein cholesterol lowering with evolocumab and outcomes in patients with peripheral artery disease: insights from the FOURIER trial (further cardiovascular outcomes research with PCSK9 inhibition in subjects with elevated risk). *Circulation* 2018;**137**:338–350. <https://doi.org/10.1161/circulationaha.117.032235>
409. Jukema JW, Szarek M, Zijlstra LE, de Silva HA, Bhatt DL, Bittner VA, et al. Alirocumab in patients with polyvascular disease and recent acute coronary syndrome: ODYSSEY OUTCOMES trial. *J Am Coll Cardiol* 2019;**74**:1167–1176. <https://doi.org/10.1016/j.jacc.2019.03.013>
410. Giugliano RP, Pedersen TR, Park JG, De Ferrari GM, Gaciong ZA, Ceska R, et al. Clinical efficacy and safety of achieving very low LDL-cholesterol concentrations with the PCSK9 inhibitor evolocumab: a prespecified secondary analysis of the FOURIER trial. *Lancet* 2017;**390**:1962–1971. [https://doi.org/10.1016/s0140-6736\(17\)32290-0](https://doi.org/10.1016/s0140-6736(17)32290-0)
411. Giugliano RP, Wiviott SD, Blazing MA, De Ferrari GM, Park J-G, Murphy SA, et al. Long-term safety and efficacy of achieving very low levels of low-density lipoprotein cholesterol: a prespecified analysis of the IMPROVE-IT trial. *JAMA Cardiol* 2017;**2**:547–555. <https://doi.org/10.1001/jamacardio.2017.0083>
412. Robinson JG, Rosenson RS, Farnier M, Chaudhari U, Sasiela WJ, Merlet L, et al. Safety of very low low-density lipoprotein cholesterol levels with alirocumab: pooled data from randomized trials. *J Am Coll Cardiol* 2017;**69**:471–482. <https://doi.org/10.1016/j.jacc.2016.11.037>
413. Marston NA, Giugliano RP, Park JG, Ruzza A, Sever PS, Keech AC, et al. Cardiovascular benefit of lowering LDL cholesterol below 40 mg/dl. *Circulation* 2021;**144**:1732–1734. <https://doi.org/10.1161/circulationaha.121.056536>
414. Koskinas KC, Windecker S, Pedrazzini G, Mueller C, Cook S, Matter CM, et al. Evolocumab for early reduction of LDL cholesterol levels in patients with acute coronary syndromes (EVOPACS). *J Am Coll Cardiol* 2019;**74**:2452–2462. <https://doi.org/10.1016/j.jacc.2019.08.010>
415. Bhatt DL, Steg PG, Miller M, Brinton EA, Jacobson TA, Ketchum SB, et al. Cardiovascular risk reduction with icosapent ethyl for hypertriglyceridemia. *N Engl J Med* 2019;**380**:11–22. <https://doi.org/10.1056/NEJMoa1812792>
416. Ray KK, Bays HE, Catapano AL, Lalwani ND, Bloedon LT, Sterling LR, et al. Safety and efficacy of bempedoic acid to reduce LDL cholesterol. *N Engl J Med* 2019;**380**:1022–1032. <https://doi.org/10.1056/NEJMoa1803917>
417. Ballantyne CM, Banach M, Mancini GBJ, Lepor NE, Hanselman JC, Zhao X, et al. Efficacy and safety of bempedoic acid added to ezetimibe in statin-intolerant patients with hypercholesterolemia: a randomized, placebo-controlled study. *Atherosclerosis* 2018;**277**:195–203. <https://doi.org/10.1016/j.atherosclerosis.2018.06.002>
418. Ray KK, Wright RS, Kallend D, Koenig W, Leiter LA, Raal FJ, et al. Two phase 3 trials of inclisiran in patients with elevated LDL cholesterol. *N Engl J Med* 2020;**382**:1507–1519. <https://doi.org/10.1056/NEJMoa1912387>
419. Wright RS, Ray KK, Raal FJ, Kallend DG, Jaros M, Koenig W, et al. Pooled patient-level analysis of inclisiran trials in patients with familial hypercholesterolemia or atherosclerosis. *J Am Coll Cardiol* 2021;**77**:1182–1193. <https://doi.org/10.1016/j.jacc.2020.12.058>
420. Katzmann JL, Packard CJ, Chapman MJ, Katzmann I, Laufs U. Targeting RNA with antisense oligonucleotides and small interfering RNA: JACC state-of-the-art review. *J Am Coll Cardiol* 2020;**76**:563–579. <https://doi.org/10.1016/j.jacc.2020.05.070>
421. Hulley S, Grady D, Bush T, Furberg C, Herrington D, Riggs B, et al. Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. *JAMA* 1998;**280**:605–613. <https://doi.org/10.1001/jama.280.7.605>
422. Manson JE, Hsia J, Johnson KC, Rossouw JE, Assaf AR, Lasser NL, et al. Estrogen plus progestin and the risk of coronary heart disease. *N Engl J Med* 2003;**349**:523–534. <https://doi.org/10.1056/NEJMoa030808>
423. Bretler DM, Hansen PR, Sørensen R, Lindhardtsen J, Ahlehoff O, Andersson C, et al. Discontinuation of hormone replacement therapy after myocardial infarction and short term risk of adverse cardiovascular events: nationwide cohort study. *BMJ* 2012;**344**:e1802. <https://doi.org/10.1136/bmj.e1802>
424. Pines A. Post-myocardial infarction hormone therapy revisited. *Climacteric* 2012;**15**:538–541. <https://doi.org/10.3109/13697137.2012.690127>
425. Paterick TE, Patel N, Tajik AJ, Chandrasekaran K. Improving health outcomes through patient education and partnerships with patients. *Proc (Bayl Univ Med Cent)* 2017;**30**:112–113. <https://doi.org/10.1080/08998280.2017.11929552>
426. Dineen-Griffin S, Garcia-Cardenas V, Williams K, Benrimoj SI. Helping patients help themselves: a systematic review of self-management support strategies in primary health care practice. *PLoS One* 2019;**14**:e0220116. <https://doi.org/10.1371/journal.pone.0220116>
427. Wolf A, Vella R, Fors A. The impact of person-centred care on patients' care experiences in relation to educational level after acute coronary syndrome: secondary outcome analysis of a randomised controlled trial. *Eur J Cardiovasc Nurs* 2019;**18**:299–308. <https://doi.org/10.1177/1474515118821242>
428. Speros CI. Promoting health literacy: a nursing imperative. *Nurs Clin North Am* 2011;**46**:321–333, vi-vii. <https://doi.org/10.1016/j.cnur.2011.05.007>
429. Shah P, Thornton I, Turrin D, Hipkind JE. Informed consent. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing, 2022.
430. Astin F, Stephenson J, Probyn J, Holt J, Marshall K, Conway D, et al. Cardiologists' and patients' views about the informed consent process and their understanding of the anticipated treatment benefits of coronary angioplasty: a survey study. *Eur J Cardiovasc Nurs* 2020;**19**:260–268. <https://doi.org/10.1177/1474515119879050>
431. Sampson F, O'Cathain A, Goodacre S. Feeling fixed and its contribution to patient satisfaction with primary angioplasty: a qualitative study. *Eur J Cardiovasc Nurs* 2009;**8**:85–90. <https://doi.org/10.1016/j.ejcnurse.2008.07.003>
432. Dickert NW, Scicluna VM, Adeoye O, Anigillo DJ, Blankenship JC, Devireddy CM, et al. Emergency consent: patients' and surrogates' perspectives on consent for clinical trials in acute stroke and myocardial infarction. *J Am Heart Assoc* 2019;**8**:e010905. <https://doi.org/10.1161/jaha.118.010905>
433. Baccolini V, Rosso A, Di Paolo C, Isonne C, Salerno C, Migliara G, et al. What is the prevalence of low health literacy in European union member states? A systematic review and meta-analysis. *J Gen Intern Med* 2021;**36**:753–761. <https://doi.org/10.1007/s11606-020-06407-8>
434. Sørensen K, Van den Broucke S, Fullam J, Doyle G, Pelikan J, Slonska Z, et al. Health literacy and public health: a systematic review and integration of definitions and models. *BMC Public Health* 2012;**12**:80. <https://doi.org/10.1186/1471-2458-12-80>
435. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* 2013;**310**:2191–2194. <https://doi.org/10.1001/jama.2013.281053>
436. Olsson A, Ring C, Josefsson J, Eriksson A, Rylance R, Fröbert O, et al. Patient experience of the informed consent process during acute myocardial infarction: a sub-study of the VALIDATE-SWEDEHEART trial. *Trials* 2020;**21**:246. <https://doi.org/10.1186/s13063-020-4147-0>
437. Dickert NW, Miller FG. Involving patients in enrolment decisions for acute myocardial infarction trials. *BMJ* 2015;**351**:h3791. <https://doi.org/10.1136/bmj.h3791>
438. Thompson DR, Ski CF. Patient-reported outcome measures in cardiovascular nursing. *Eur J Cardiovasc Nurs* 2015;**14**:370–371. <https://doi.org/10.1177/1474515115601621>

439. Committee on Quality of Health Care in America, Institute of Medicine. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington (DC): National Academies Press (US), 2001.
440. Hirpa M, Woreta T, Addis H, Kebede S. What matters to patients? A timely question for value-based care. *PLoS One* 2020;**15**:e0227845. <https://doi.org/10.1371/journal.pone.0227845>
441. Rossello X, Massó-van Roessel A, Perelló-Bordoy A, Mas-Lladó C, Ramis-Barceló MF, Vives-Borrás M, et al. Assessment of the ESC quality indicators in patients with acute myocardial infarction: a systematic review. *Eur Heart J Acute Cardiovasc Care* 2021;**10**: 878–889. <https://doi.org/10.1093/ehjacc/zuab042>
442. Fållun N, Fridlund B, Schaufel MA, Schei E, Norekvål TM. Patients' goals, resources, and barriers to future change: a qualitative study of patient reflections at hospital discharge after myocardial infarction. *Eur J Cardiovasc Nurs* 2016;**15**:495–503. <https://doi.org/10.1177/1474515115614712>
443. Goldman JD, Harte FM. Transition of care to prevent recurrence after acute coronary syndrome: the critical role of the primary care provider and pharmacist. *Postgrad Med* 2020;**132**:426–432. <https://doi.org/10.1080/00325481.2020.1740512>
444. Huriani E. Myocardial infarction patients' learning needs: perceptions of patients, family members and nurses. *Int J Nurs Sci* 2019;**6**:294–299. <https://doi.org/10.1016/j.ijnss.2019.05.001>
445. Messerli AW, Deutsch C. Implementation of institutional discharge protocols and transition of care following acute coronary syndrome. *Cardiovasc Revasc Med* 2020;**21**:1180–1188. <https://doi.org/10.1016/j.carrev.2020.02.013>
446. Schiele F, Lemesle G, Angoulvant D, Krempf M, Kownator S, Cheggour S, et al. Proposal for a standardized discharge letter after hospital stay for acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:788–801. <https://doi.org/10.1177/2048872619844444>
447. Wittink H, Oosterhaven J. Patient education and health literacy. *Musculoskelet Sci Pract* 2018;**38**:120–127. <https://doi.org/10.1016/j.msksp.2018.06.004>
448. Miller S, Lattanzio M, Cohen S. 'Teach-back' from a patient's perspective. *Nursing* 2016;**46**:63–64. <https://doi.org/10.1097/01.NURSE.0000476249.18503.f5>
449. Zhao Y, Izadnegahdar M, Lee MK, Kavsak PA, Singer J, Scheuermeyer F, et al. High-sensitivity cardiac troponin-optimizing the diagnosis of acute myocardial infarction/injury in women (CODE-MI): rationale and design for a multicenter, stepped-wedge, cluster-randomized trial. *Am Heart J* 2020;**229**:18–28. <https://doi.org/10.1016/j.ahj.2020.06.013>